

Which treatment classes and combinations are more effective for the management of common tendinopathies? A systematic review and network meta-analysis.

Swinton, P.A.^{1*} Shim, J.¹ Pavlova, A.V.¹ Moss, R.A.¹ Maclean, C.² Brandie, D.³ Mitchell, L.⁴ Tzortziou Brown, V.⁵ Greig, L.¹ Parkinson, E.¹ Morrissey, D.⁶ Alexander, L.¹ Cooper, K.¹

Doi: 10.51224/SRXIV.155

SportRxiv hosted preprint version 1

11/05/2022

PREPRINT - NOT PEER REVIEWED

Institutions

¹ School of Health Sciences, Robert Gordon University, Aberdeen, UK

² Library Services, Robert Gordon University, Aberdeen, UK

³ Sport Scotland, Inverness, UK

⁴ NHS Grampian, Aberdeen, UK

⁵ Wolfson Institute of Population Health, Queen Mary University of London, UK

⁶ Centre for Sports & Exercise Medicine, Barts & the London School of Medicine & Dentistry, London, UK

* Corresponding Author

Dr. Paul Swinton

School of Health Sciences, Robert Gordon University

Garthdee Road

Aberdeen, UK,

AB10 7QG

p.swinton@rgu.ac.uk, +44 (0) 1224 262 3361

Please cite this paper as: Swinton PA, Shim J, Pavlova AV, Moss RA, MacLean C, Brandie D, Mitchell L, Tzortziou Brown V, Greig L, Parkinson E, Morrissey D, Alexander L, Cooper K. Which treatment classes and combinations are more effective for the management of common tendinopathies? A systematic review and network meta-analysis. Pre-print available from SportRxiv. Doi: 10.51224/SRXIV.155.

Abstract

Objective To quantify the comparative effectiveness of treatment classes used for the management of the most common tendinopathies.

Design Network meta-analyses comparing combinations of exercise, non-exercise, and non-active treatments across a range of tendinopathy locations and outcome domains.

Eligibility criteria Randomised and quasi-randomised controlled trials including an exercise arm and persons with a tendinopathy diagnosis at any location and of any severity or duration.

Outcome measures Outcomes assessing disability, function, pain, shoulder range of motion, physical function capacity, or quality of life.

Methods Network meta-analyses of broad (exercise/non-exercise/combined/non-active) and more specific (exercise/biomechanics/injection/electrotherapy/manual-therapy/non-active/surgery) treatment class models were fitted with hierarchical Bayesian models. Results were interpreted using pooled standardised mean difference effect sizes and ranking through Surface Under the Cumulative Ranking curves (SUCRA). Treatment hierarchies were assessed using the GRADE minimally contextualised framework.

Results Two-hundred studies comprising 458 treatments arms were identified. Many comparisons were within the same class reducing data available to assess comparative effectiveness. Data from 85 studies generating 140 pairwise comparisons consistently identified the superiority of combining exercise and non-exercise treatment classes (SUCRA: 0.70 to 0.88). Central estimates indicated that combining exercise and non-exercise treatments increased effect sizes by ~0.1 to 0.3 compared with exercise alone. Analysis of more specific treatment classes identified with low/very low certainty the superiority of combining exercise with either biomechanical (e.g. taping, bracing or splinting; SUCRA: 0.73) or injection therapies (SUCRA: 0.72).

Summary/Conclusion Clinicians should consider as a starting point for tendinopathy management combining exercise and non-exercise therapies. The most effective treatment combinations include exercise with the use of biomechanical or injection therapies.

Keywords: Tendinopathy, Exercise therapy; Physiotherapy; Effect size

Introduction

The clinical management of symptomatic tendinopathy requires complex clinical reasoning with reference to the pathoanatomical diagnosis. Management strategies often vary depending on the stage and location of the tendinopathy, contributing issues within the kinetic chain, and patient factors including activity level, psychosocial factors and comorbidities.¹ Current research supports the role of appropriate load management and progressive rehabilitation that includes strength training as the primary treatment for tendinopathy.² Different loading types, including eccentric, combined, heavy-slow-resistance, and isometric, have each been recommended with similar goals including initiation of structural tendon adaptation, reduction of pain and restoration of function.³ In order to optimise outcomes, which are at best partial and typically slow, management may combine exercise with other treatment categories (classes) including manual-therapy,⁴ electrotherapy,⁵ biomechanical interventions,⁶ injection therapy of various types,⁷ and surgery typically reserved for the most recalcitrant.⁸

Currently, the best choice of treatment class, or combination of classes for the management of tendinopathy remains uncertain. Previous systematic reviews have generally focused on single tendinopathies and resorted to narrative syntheses due to concerns of both statistical and clinical heterogeneity.^{9,10} Where meta-analyses have been conducted, these typically pool data from small numbers of homogenous studies employing conventional pairwise approaches that limit inference regarding comparative effectiveness across a range of treatment classes and subsequent development of treatment hierarchies. A range of contemporary approaches including the use of network meta-analyses (NMA) that can account for multiple outcomes reported in the same study have been developed to better synthesise complex data.¹¹ Recent NMAs investigating the management of tendinopathy, however, have focused on localised site-specific tendinopathies with pain relief and function as the primary outcomes.¹²⁻¹⁶ Four NMAs have investigated comparative effectiveness of treatments in the upper extremity, three of which studied injection treatments in the shoulder¹³ and elbow,^{14,15} while another focused on a range of non-surgical treatments for chronic calcific tendinitis of the shoulder.¹² In an NMA investigating non-surgical treatments for patellar tendinopathy comprising eleven trials, Chen et al.¹⁶ concluded that platelet-rich plasma therapy results in the greatest improvements in pain and function compared with other treatments. The review, however, excluded studies that compared different types of exercise treatments from their analysis.

Two recent NMAs investigating the management of Achilles tendinopathy reported somewhat conflicting findings. In an analysis comprising twenty-nine trials, van der Vlist et al.¹⁷ concluded there was strong evidence that all active treatments were superior to wait-and-see, but no one active treatment could be recommended over another. In contrast, Rhim et al.¹⁸ reported that improvements were highest when Swinton, Shim, Pavlova, Moss, Maclean, Brandie, Mitchell, Tzortziou Brown, Greig, Parkinson, Morrissey, Alexander, Cooper

(2022)

eccentric exercise was combined with high-volume injection and corticosteroid, or combined with extracorporeal shockwave therapy. Confidence in these findings were low, however, as the results from the highest ranked therapies were derived from only two studies. Such comparisons across relatively fine-grained treatment classes from small numbers of homogenous studies are unable to address questions regarding general trends that can be derived from comparisons across many studies and popular treatment classes using data from different tendinopathy locations and outcome domains. Therefore, the purpose of the present systematic review and NMA was to quantify comparative effectiveness of broad treatment classes across the tendinopathy management literature, assessing whether combining exercise and non-exercise treatments were more effective than either in isolation. Following this general overview, the review sought to quantify comparative effectiveness of more specific combinations of treatment classes to inform clinical practice, service redesign, and future research.

Methods

This review was part of a project funded by the National Institute for Health Research (Health Technology Assessment: 129388 Exercise therapy for the treatment of tendinopathies) which examined the exercise therapy evidence base for multiple tendinopathies to make research and practice recommendations. The methods reported here were influenced by the overall project aims, the results of an initial scoping review mapping the exercise and tendinopathy literature,¹⁹ as well as stakeholder workshops. The review was conducted according to the PRISMA extension statement for reporting of systematic reviews incorporating NMAs of health care interventions (Supplementary file 1)²⁰ and the recent Grading of Recommendations Assessment Development and Evaluation (GRADE) approach to drawing conclusions from NMA using a minimally contextualised framework.²¹ An a priori protocol was created and followed for this review.²²

Inclusion criteria

This review included randomized and non-randomized controlled trials comprising at least two trial arms featuring different treatment classes. As exercise therapy was the focus of the overall project, all studies included at least one trial arm that featured exercise components. This review included people of any age or gender with a diagnosis of tendinopathy of any severity or duration and at any anatomical location. Based on the results of our initial reviews^{19,23} and stakeholder workshops we included outcomes that assessed six domains: 1) disability; 2) function; 3) pain (e.g. pain on loading, pain over a specified time, pain without further specification); 4) range of motion for the shoulder joint; 5) physical function capacity (PFC, e.g., objective assessment of hops/stair-climbing/squats); and 6) quality of life (QoL). Definitions of each domain and example tools are presented in supplementary file 2. We also included studies with trial arms whose treatment class comprised: 1) exercise; 2) non-active (e.g. placebo, sham, wait-and-see); 3) injection; 4) electrotherapy; 5) biomechanics; 6) manual-therapy; or 7) surgery. Definitions of each treatment class are presented in supplementary file 3. Inclusion was restricted to studies conducted in primary care, secondary care or community locations in any developed nation (defined as the top 62 countries in the Human Development Index at the time of protocol development).

Search strategy

The search strategy used for this study was part of a larger search conducted to scope the entire exercise therapy for tendinopathy management research base. The search comprised three steps; firstly, a limited search of MEDLINE and CINAHL using initial keywords (MH tendinopathy OR TX tendin* OR TX tendon*) AND (MH exercise OR TX exercis*) was conducted with analysis of the text words in the titles/abstracts and those used to describe articles to develop a full search strategy. Secondly, the full search strategy was adapted to each database and applied systematically to: MEDLINE, CINAHL, AMED, EMBase, SPORTDiscus, Cochrane library (Controlled trials, Systematic reviews), JBI Evidence Synthesis, PEDRo, and Epistemonikos. The following trial registries were also searched: ClinicalTrials.gov, ISRCTN

Registry, The Research Registry, EU-CTR (European Union Clinical trials Registry), ANZCTR (Australia and New Zealand Clinical trials Registry). Finally, the third step involved conducting a search of cited and citing articles using Scopus and hand-searching a total of 130 systematic reviews that were identified to include information relevant to exercise therapy and tendinopathy. No limit was placed on language, with research studies published in languages other than English translated via Google Translate or via international collaborations of the review team members. Searches were initiated from 1998 as (i) the heavy load eccentric calf-training protocol for Achilles tendinosis by Alfredson et al ²⁴ was published in 1998 and may be considered seminal work in the field of tendinopathy, and (ii) there has been a proliferation of research on exercise interventions for tendinopathies post 1998. Search terms and results for MEDLINE are presented in Supplementary file 4 according to the last date of the search which was 21/01/21.

Study selection and Data extraction

Two independent reviewers screened titles and abstracts followed by full-text copies. Conflicts were resolved by a third reviewer with all screening conducted within the Covidence (Melbourne, Australia) platform. Data were extracted independently by eight members of the review team (PS/KC/LA/RM/LG/EP/JS/AP) into pre-piloted excel sheets and coded as described in the codebook presented in the Supplementary file 5. Each entry was then independently checked.

Risk of bias assessment

We used Cochrane's Risk of Bias (RoB) tool ²⁵ to assess six domains: 1) selection bias (random sequence generation & allocation concealment); 2) performance bias (blinding of participants); 3) detection bias (blinding of outcome assessors); 4) attrition bias (incomplete outcome data); 5) reporting bias (selective reporting); and 6) other biases. RoB was recorded for each outcome and time point within each study. The Cochrane's RoB tool ²⁵ was selected as a recent review of popular tools in tendinopathy management highlighted none were superior ²⁶ and Cochrane's RoB tool ²⁵ could be semi-automated with RobotReviewer,²⁷ a machine learning system software. RobotReviewer was used to make initial assessments on selection bias and performance bias domains, with manual validation made on the relevant free texts extracted to support the final selection of low, high, or unclear RoB. This semi-automated process was more efficient and provided an additional element of consistency in the review process.

Statistical analysis

We fitted treatment class (broad and specific) Bayesian NMA models, first categorising broad treatments as exercise only, non-exercise only, non-active, or combined exercise and non-exercise. Second, more specific treatments classes and their combination (e.g. exercise + electrotherapy, or exercise + manual-therapy + injection) were categorised. All outcome measures included in meta-analysis models were continuous with comparative pairwise effect sizes calculated using standardised mean differences

Swinton, Shim, Pavlova, Moss, Maclean, Brandie, Mitchell, Tzortziou Brown, Greig, Parkinson, Morrissey, Alexander, Cooper (2022)

(SMD_{pre}), where the mean difference from one trial arm was subtracted from another and standardised by dividing by the pooled baseline standard deviation. A small sample size correction was made,²⁸ and where required SMD_{pre} values were reflected by multiplying by -1 to ensure that positive values represented an improved clinical effect.

Hierarchical class-level models were used to account for the inclusion of multiple outcomes from the same study. Models were fitted in a Bayesian framework using Markov chain Monte Carlo simulations. Analyses were conducted initially with data pooled across all tendinopathy locations and outcome domains, with subset analyses conducted across the individual levels where sufficient data was available. We planned to conduct moderator analyses investigating the effects of assessment duration after separating outcomes according to short (≤ 12 weeks), medium (13-52 weeks) and long (> 52) time frames. However, data were predominantly collected across short time frames and analyses resulted in estimates that did not converge. Following the initial set of analyses, attempts were made to quantify comparative effectiveness across all unique treatment combinations. Stepwise pruning of the network was conducted by removing the node comprising the lowest number of trial arms until all network parameter estimates converged producing plausible values. Ranking of effectiveness across treatment classes was summarised through calculation of Surface Under the Cumulative Ranking curves (SUCRA) which ranges from zero to one.²⁹ The higher the SUCRA value, the higher the probability that a therapy is in the top ranks of those treatments considered. Analyses were performed using WinBUGS³⁰ and the R package R2WinBUGS.³¹

Following the GRADE approach for presentation and interpretation of results, the most connected node in the network was selected as the reference.²⁵ Using the minimally contextualised framework, a no effect threshold was adopted and any treatment class where the 95% credible interval did not span the threshold was moved above or below the reference accordingly. Second classifications were then made based on comparisons of treatment classes moved relative to the reference. In each of the classifications, treatment classes were separated into moderate to high certainty, and low to very low certainty based on: 1) overall risk of bias ranked as high, low or unclear risk (as identified by the mode rating across all data in the specific analysis); 2) inconsistency assessed comparing the NMA and direct reference pairwise effect sizes; 3) imprecision judged by the number of available data points (studies, trial arms, outcome measures) and width of effect size credible intervals; 4) indirectness identified as low risk for all outcomes based on inclusion criteria from our previous scoping review and stakeholder recommendation; and 5) small-study effects assessed by visual inspection of effect size distribution and sampling variance. Categorisations of overall certainty in evidence began with high certainty in cumulative evidence and were downgraded a level for each domain not judged as low risk.

Results

A flow diagram illustrating study selection with reasons for exclusions is presented in figure 1. A total of 200 studies (Supplementary file 6) were identified comprising 458 trial arms, 326 direct pairwise comparisons, and 11,873 participants. A breakdown of the tendinopathies investigated, the outcome domains measured, and the treatment class combinations are presented in table 1. The most frequently investigated tendinopathy locations included rotator cuff (43.0% of trial arms), elbow (22.5% of trial arms) and Achilles (20.3%). Pain (74.9% of trial arms) and disability (80.1% of trial arms) were the most frequently measured outcome domains across a range of treatment combinations comprising primarily exercise, electrotherapy, and manual-therapy (table 1). Summary risk of bias from each of the 200 included studies are presented in Supplementary file 7.

Figure 1: PRISMA chart illustrating study selection.




Table 1: Tendinopathies and outcome domains investigated across 200 studies and 458 trial arms.

Tendinopathies	Number of trial arms (%)	Outcome Domain	Number of outcomes (%) / Number of trial arms (%)
Rotator cuff	197 (43.0)	Pain	961 (32.2) / 343 (74.9)
Elbow	103 (22.5)	Disability	856 (28.6) / 367 (80.1)
Achilles	93 (20.3)	Physical function capacity	531 (17.8) / 176 (38.4)
Patellar	43 (9.4)	Range of motion	300 (10.0) / 82 (17.9)
Gluteal	9 (2.0)	Function	203 (6.8) / 102 (22.3)
Tibialis posterior	9 (2.0)	Quality of Life	137 (4.6) / 46 (10.0)
Hamstring	2 (0.4)		
Biceps	2 (0.4)		


Treatment Combinations	Number of trial arms (%)	Treatment Combinations	Number of trial arms (%)
Exercise only	180 (39.3)	Exercise + Electrotherapy	63 (13.8)
Non-exercise only	36 (7.9)	Exercise + Biomechanics	30 (6.6)
Exercise + Non-active	30 (6.6)	Exercise + Manual-therapy	27 (5.9)
Exercise + Injection	23 (5.0)	Exercise + Electrotherapy + Manual-therapy	22 (4.8)
Non-active	18 (3.9)	Exercise + Surgery	8 (1.7)
Exercise + Electrotherapy + Biomechanics + Manual-therapy	4 (0.9)	Exercise + Manual-therapy + Non-active	4 (0.9)
Exercise + Electrotherapy + Non-active	3 (0.7)	Exercise + Injection + Manual-therapy	3 (0.7)
Exercise + Injection + Biomechanics	2 (0.4)	Exercise + Biomechanics + Manual-therapy	2 (0.4)
Exercise + Electrotherapy + Injection + Manual-therapy	1 (0.2)	Exercise + Electrotherapy + Biomechanics	1 (0.2)
Injection + Non-active	1 (0.2)		

Non-active treatments comprised placebo, sham or wait-and-see.

The relative frequencies of the broad treatment classes across the 458 trial arms were: combined exercise and non-exercise (48.9% of trial arms); exercise only (39.3% of trial arms); non-exercise only (7.9% of trial arms); and non-active (3.9% of trial arms). The network diagram illustrating the 326 direct pairwise comparisons across the broad treatment classes is presented in figure 2, with most comparisons occurring within the combined exercise and non-exercise class, and a substantive number occurring within the exercise only class. In total, 85 studies provided data comparing different broad treatment classes to generate 140 pairwise comparisons to be included in the primary NMA combining all tendinopathy locations and outcome domains. Results from the primary NMA and subset NMAs across individual tendinopathy locations are presented in Table 2. Combined exercise and non-exercise treatments were ranked most effective (SUCRA: 0.70 to 0.88), and non-active treatments ranked least effective (SUCRA: 0.22 to 0.39) across all analyses. When pooling data across all tendinopathy locations, non-active and exercise-only treatments were identified as inferior with high and low certainty, respectively. Although central estimates indicated that non-exercise only treatments were also inferior, credible intervals were

wide such that it was deemed with low certainty these treatments may be among the most effective (along with combined exercise and non-exercise). When restricting analyses to studies comparing outcomes for rotator cuff tendinopathy, all mean differences and credible intervals indicated combining exercise and non-exercise treatments was superior. Where certainty of evidence was low, this was due to likely bias associated with inconsistency and small-study effects generating large numbers of very large effect sizes. Collectively, central estimates indicated that combining exercise and non-exercise treatments produced on average a ~ 0.1 to 0.3 increase in standardised mean difference effect size (Table 2). Analyses combining results across all tendinopathy locations but separated according to outcome domains are presented in Figure 3. Across all outcome domains SUCRA values were highest for combined exercise and non-exercise treatments (SUCRA: 0.71 to 0.88) and lowest for non-active treatments (SUCRA: 0.07 to 0.30). Frequently, exercise only treatments ranked second (SUCRA: 0.45 to 0.71), followed by non-exercise only treatments (SUCRA: 0.37 to 0.69).

Figure 2: Network diagram illustrating distribution of pairwise comparisons between broad treatment classes. The size of each node is scaled to the number of pairwise comparisons.




The size of each node is scaled to the number of pairwise comparisons. Values on each edge quantifies the number of direct pairwise comparisons between nodes.

Table 2: Application of minimally contextualised GRADE framework to rank effectiveness of broad treatment classes for tendinopathy management. Analyses presented for network meta-analyses conducted with pooled data across all tendinopathy locations, and individual tendinopathies where sufficient data were available.

Analysis	Number of treatment comparisons/ Number of outcomes	Broad treatment class	Comparison with Exercise + Non-Exercise median difference (95% CrI)	Surface under the cumulative ranking
All tendinopathies				
High certainty (moderate to high certainty evidence)				
Category 2: Among the most effective	100/1041	Exercise + Non-exercise	NA	0.86
Category 1: Inferior to the most effective	28/317	Non-active	-0.22 [-0.43 to -0.03]	0.22
Low certainty (low to very low certainty evidence)				
Category 2: Might be among the most effective	50/454	Non-exercise only	-0.12 [-0.46 to 0.22]	0.44
Category 1: Might be inferior to the most effective	103/916	Exercise only	-0.10 [-0.21 to -0.01]	0.49
Rotator cuff				
High certainty (moderate to high certainty evidence)				
Category 2: Among the most effective	50/595	Exercise + Non-exercise	NA	0.88
Category 1: Inferior to the most effective	8/100	Non-active	-0.21 [-0.41 to -0.01]	0.39
Low certainty (low to very low certainty evidence)				
Category 1: Might be inferior to the most effective	14/146	Non-exercise only	-0.38 [-0.55 to -0.21]	0.31
	50/517	Exercise only	-0.19 [-0.36 to -0.03]	0.41
Achilles				
High certainty (moderate to high certainty evidence)				
Category 2: Among the most effective	17/109	Exercise + Non-exercise	NA	0.71
Low certainty (low to very low certainty evidence)				
	15/111	Non-exercise only	-0.16 [-0.71 to 0.38]	0.45
Category 2: Might be among the most effective	21/133	Exercise only	-0.10 [-0.55 to 0.35]	0.55
	5/7	Non-active	-0.30 [-0.93 to 0.33]	0.29
Elbow				
High certainty (moderate to high certainty evidence)				
Category 2: Among the most effective	27/283	Exercise + Non-exercise	NA	0.70
Low certainty (low to very low certainty evidence)				
	14/153	Non-exercise only	-0.09 [-0.65 to 0.50]	0.45
Category 2: Might be among the most effective	21/164	Exercise only	-0.08 [-0.41 to 0.26]	0.46
	12/170	Non-active	-0.11 [-0.42 to 0.22]	0.38

Negative effect sizes favour exercise combined with non-exercise Number of treatment comparisons quantifies the total number of all pairwise comparisons that include the specific treatment class. CrI: Credible interval.


Figure 3: Surface under the cumulative ranking (SUCRA) values for broad treatment classes pooled across all tendinopathy locations for different outcome domains.



PFC: Physical function capacity; **QoL:** Quality of life; **ROM:** Range of motion.

The network-diagram illustrating the pairwise comparisons between exercise and exercise combined with different non-exercise treatments is presented in Figure 4. Most comparisons were within the same class, or comparisons with either exercise-only or exercise plus non-active treatments. As a result, there were limited direct comparisons between combined exercise and active non-exercise treatments (e.g. exercise + electrotherapy; exercise + biomechanics; and exercise + injection). Due to the low number of direct comparisons, initial NMAs failed to converge producing plausible values for pairwise estimates. Pruning of the network was conducted and estimates converged for a NMA pooling data across all tendinopathy locations and outcome domains, which comprised six combined treatment classes (exercise + biomechanics; exercise + injection; exercise + electrotherapy; exercise + manual-therapy; exercise + electrotherapy + manual-therapy; exercise + non-active) and exercise only therapy as a reference (Table 3). Exercise combined with biomechanics interventions (SUCRA:0.73) or injection therapy (SUCRA: 0.72) were identified with low certainty as the most effective treatments, with all other treatments identified with low certainty as inferior (SUCRA: 0.27 to 0.50).

Figure 4: Network diagram illustrating distribution of pairwise comparisons between treatment classes. The size of each node is scaled to the number of pairwise comparisons.



The size of each node is scaled to the number of pairwise comparisons. Values on each edge quantifies the number of direct pairwise comparisons between nodes.

Table 3: Application of minimally contextualised GRADE framework to rank effectiveness of treatment classes for tendinopathy management. Analyses presented for network meta-analysis conducted with pooled data across all tendinopathy locations.

Analysis	Number of treatment comparisons/ Number of outcomes	Exercise and combined treatment classes	Comparison with Exercise only median difference (95% CrI)	Surface under the cumulative ranking
All tendinopathies				
Low certainty (low to very low certainty evidence)				
Category 2: Might be among the most effective	19/182	Exercise + Biomechanics	0.24 [0.03 to 0.41]	0.73
	15/126	Exercise + Injection	0.24 [0.01 to 0.45]	0.72
	5/41	Exercise + Electrotherapy + Manual-therapy	0.08 [-0.29 to 0.46]	0.50
	35/301	Exercise + Electrotherapy	0.07 [-0.32 to 0.48]	0.50
Category 1: Might be inferior to the most effective	35/307	Exercise + Non-active	0.03 [-0.19 to 0.25]	0.40
	53/439	Exercise only	NA	0.37
	18/162	Exercise + Manual-therapy	-0.15 [-0.50 to 0.25]	0.27

Negative effect sizes favour exercise only. Number of treatment comparisons quantifies the total number of all pairwise comparisons that include the specific treatment class. CrI: Credible interval.

Discussion

This systematic review and NMA represents one of the most comprehensive attempts at quantitative evidence synthesis of the effectiveness of relatively broad treatment classes for the management of common tendinopathies. Consistent findings were obtained across different tendinopathy locations and outcome domains that combining exercise and non-exercise treatment classes were superior to either exercise in isolation or to non-active treatments such as placebo, sham or wait-and-see. Central estimates from NMAs indicated that combining exercise and non-exercise treatments may produce on average ~0.1 to 0.3 increases in standardised mean difference effect sizes compared with exercise alone. An overview of all treatment comparisons identified that trials tended to make comparisons within treatment classes rather than between. As a result, the ability to compare across classes and identify clear treatment hierarchies was limited. The results obtained here, however, indicate with low certainty, that exercise combined with biomechanical interventions or injection treatments may be the most effective. Whilst group and individual responses to treatment are likely to be influenced by a wide range of factors, the findings obtained in the present review provide insight into general and consistent patterns that can inform clinical practice, service redesign and future research.

The present review is one of the few evidence synthesis studies to combine information across different tendinopathy locations and found that exercise is best combined with other treatments, with there being a particular need for further assessment of this finding at medium- and long-term follow-up. The relative frequencies of treatment classes identified in the present review are likely influenced by its focus and the associated inclusion criteria that all studies required at least one trial arm to comprise exercise therapy. The focus does, however, reflect the move towards therapeutic exercise as a mainstay of rehabilitation supported by a large volume of evidence in the form of trials and systematic reviews.³²⁻³⁵ The finding that combining exercise and non-exercise treatments may result in additional small to moderate clinical benefit is in line with previous evidence syntheses. In a recent umbrella review evaluating twenty-five systematic reviews of high quality, Irby et al.³⁶ concluded that exercise therapy is the best treatment option for tendinopathy when combined with the use of other therapeutic modalities. Additionally, previous NMAs conducted by van der Vlist et al.¹⁷ and Rhim et al.¹⁸ both ranked exercise combined with shockwave or injection therapies more favourably than exercise alone for Achilles tendinopathy. Rhim et al.¹⁸ further identified the largest sustained improvements at twelve months with treatments combining exercise and shockwave. It seems likely that the consistent recommendation to combined interventions reflects different mechanisms of therapeutic effect alongside the partial efficacy observed for even the best interventions. Further, the additional benefit from combining interventions is consistent with clinical reasoning. In the present review we planned to assess whether comparative effectiveness was influenced by assessment duration. Most outcomes measurements, however, were conducted across short time

frames such that any comparison was unlikely to be stable, indicating that further research with longer follow-up times is required.

Following evidence of superior effectiveness when combining exercise and non-exercise treatments, the review sought to identify potential differences across different treatment combinations. With low certainty and limitations in fitting a suitable NMA model, we identified that combining exercise with either biomechanical interventions or injection treatments may be superior to other combinations such as exercise and electrotherapy and/or manual-therapy. Analysis of the network structure demonstrated that most comparisons were conducted within treatment classes. This may reflect researchers' and clinicians' interest in more specific manipulations to popular treatments including dosing parameters. In addition, where comparisons were made outside of the same treatment class, this was most often with exercise only or exercise combined with non-active treatments acting as control conditions. The lack of comparison across different treatment classes limits the ability to establish general treatment hierarchies as the relatively small number of direct comparisons meant only a single NMA with a restricted range of comparisons could be conducted. The need for more trials comparing combinations of treatments across treatment classes is notable and would ideally follow guidelines on intervention development to ensure the optimal likelihoods of successful comparisons of meaningful interventions.³⁷

In addition to uncertainty due to a lack of direct comparisons, uncertainty of the most effective combination of treatments stems from the fact that the pathophysiology of tendinopathy and the working mechanisms of existing treatments are not completely understood.^{38,39} Some evidence suggests a failed healing response due to degeneration caused by repetitive overloading rather than an inflammatory process,³⁸⁻⁴⁰ while more recent thinking suggest a stronger role for low-grade inflammation.⁴¹ Additionally, the disconnect between observed structural change and reported symptomatic state, and the absence of effective biomarkers of tendinopathy severity limit clinical innovation. The proposed mechanisms for biomechanics interventions depend upon the specific implementation. Taping is purported to act by limiting joint movement (rigid tape), influencing circulatory and neurological systems (kinesiotape), and altering the loading of musculotendinous structures (biomechanical tape).⁴² Counterforce bracing for lateral elbow tendinopathy is reported to have mechanical and neurological effects on elbow proprioception, essentially creating a new tendon origin, allowing for tissue healing.⁴³ There is limited evidence that they may provide short-term pain relief,⁴⁴ especially in younger patients.⁴³ Night splints have been used mainly in Achilles tendinopathy, and are thought to optimize tendon microcirculation.⁴⁵ In contrast, corticosteroid injections reduce pain caused by inflammatory agents but are known to decrease collagen production and therefore, slow the healing process. Given the potential weakening of the tendon matrix structure, there are concerns for the long-term use of corticosteroid injections especially for Achilles tendinopathy.⁴⁶ More recently, the use of regenerative injections has Swinton, Shim, Pavlova, Moss, Maclean, Greig, Parkinson, Morrissey, Alexander, Cooper (2021)

emerged as a potential treatment to facilitate tissue healing and regeneration in tendinopathy with long-term improvements.⁴⁷ Whilst there was less evidence for an additional benefit of combining electrotherapy with exercise, the electrotherapy class comprises a range of different technologies that are widely applied in clinical settings, with shockwave the most commonly used.⁴⁸ Studies have postulated that shockwave therapy may influence nociceptive transmission in the central nervous system, which provoke peripheral, sensory nerve fibres resulting in pain relief.⁴⁹ It is also plausible that shockwave has an indirect influence on neovascularisation which leads to better tissue regeneration in tendinopathies through a better blood supply.⁵⁰ Collectively, improved biomarkers of tendon pathology and clarity of therapeutic mechanisms would enable more targeted treatment, particularly when the innovations afforded by precision medicine are applied to tendinopathy.

The findings of the present review have implications for the design and commissioning of services for the management of tendinopathies. The potential for additional improvements combining biomechanical interventions with exercise compared to exercise alone is important at a time when access to such services is increasingly restricted. In addition, if injections are shown to lead to the same long-term superior results as identified in the short-term when combined with exercise, this will have implications on the need for relevant training and service provision. The inclusion of any of the active treatments investigated in the present analysis require face to face patient assessments while exercise therapy on its own may have comparable effectiveness when delivered remotely – at least in part.

It was outside the remit of this review to compare the cost-effectiveness of different treatment classes but there is a need for such cost-effectiveness analyses in the future to further inform service redesign. In considering the findings and suggested implications of the present review, several limitations are relevant. The focus of the review was on exercise therapy and therefore not all relevant trials comparing non-exercise therapies were included. The present review sought to identify more general trends in the management of tendinopathy, synthesising evidence across a range of outcome domains, tendinopathy locations and broad treatment classes. Whilst this provided the benefit of increasing the available data to synthesise, there is likely to be large variation in expected trial results of any specific treatment class or combination based on interactions of factors such as the specifics of the treatment (e.g. type of electrotherapy and dose), the tendinopathy location, the outcome and the population. Further research is required to identify the extent and pattern with which these factors may influence the comparative effectiveness of different treatments. Treatment heterogeneity effects should also be expected at the individual level, where patient characteristics interact with treatments to determine which are most effective. Therefore, the findings obtained here and the associated low certainty in the evidence, provide a starting point for clinicians in the treatment prescribed to each individual, and that in general combining exercise with other classes such as biomechanics interventions or injection therapy may be most effective. Swinton, Shim, Pavlova, Moss, Maclean, Greig, Parkinson, Morrissey, Alexander, Cooper (2021)

This systematic review and NMA also highlights novel insight into the necessary changes in future trial designs on tendinopathy.

Funding

This project is funded by the National Institute for Health Research (NIHR) [Health Technology Assessment (HTA) 129388 Exercise therapy for the treatment of tendinopathies]. The views expressed are those of the authors and not necessarily those of the NIHR or the Department of Health and Social Care.

Conflicts of interest

The authors declare no conflict of interest.

References

1. Scott A, Docking A, Vicenzio B, Alfredson H, Zwerver J, Lundgreen K, Finlay O, Pollock N, Cook JL, Fearon A, Purdam CR, Hens A, Rees JD, Goetz TJ, Danielson P. Sports and exercise-related tendinopathies: A review of selected topical issues by participants of the second International Scientific Tendinopathy Symposium (ISTS) Vancouver 2012. *Br J Sports Med*. 2013; 47(9): 536-544.
2. Magnusson PS, Langberg H, Kjaer M. The pathogenesis of tendinopathy: Balancing the response to loading. *Nat Rev Rheumatol*. 2010; 6(5): 262-268.
3. Brumitt J, Cuddeford T. Current concepts of muscle and tendon adaptation to strength and conditioning. *Int J Sports Phys Ther*. 2015 Nov; 10(6): 748-759.
4. Jayaseelan DJ, Kecman M, Alcorn D, Sault JD. Manual therapy and eccentric exercise in the management of Achilles tendinopathy. *J Man Manip Ther*. 2017; 25(2): 106-114.
5. Carlisi E, Lisi C, Dall'angelo A, Monteleone S, Nova V, Tinelli C et al. Focused extracorporeal shockwave therapy combined with supervised eccentric training for supraspinatus calcific tendinopathy. *J Phys Med Rehabil Med*. 2018; 54(1): 41-47.
6. de Jonge S, de Vos RJ, Van Schie HT, Verhaar JA, Weir A, Tol JL. One-year follow-up of a randomised controlled trial on added splinting to eccentric exercises in chronic midportion Achilles tendinopathy. *BJSM*. 2010; 44(9): 673-677.
7. Van Ark M, Van den Akker-Scheek I, Meijer LTB, Zwerver J. An exercise-based physical therapy program for patients with patellar tendinopathy after platelet-rich plasma injection. *Phys Ther Sport* 2013; 14(2):124-130.
8. Morrissey D. Guidelines and Pathways for Clinical Practice in Tendinopathy: Their Role and Development. *J Orthop Sports Phys Ther*. 2015; 45(11): 819-822.
9. Lim HY, Wong SH. Effects of isometric, eccentric, or heavy slow resistance exercises on pain and function in individuals with patellar tendinopathy: A systematic review. *Physiother Res Int*. 2018; 23(4): e1721.
10. Page MJ, Green S, Mrocki MA, Surace SJ, Deitch J, McBain B, Lyttle N, Buchbinder R. Electrotherapy modalities for rotator cuff disease. *Cochrane Database Syst Rev*. 2016; 2016(6): CD012225.
11. Lorenc T, Felix L, Petticrew M, Melendez-Torres GJ, Thomas J, Thomas S, O'Mara-Eves A, Richardson M. Meta-analysis, complexity, and heterogeneity: A qualitative interview study of researchers' methodological values and practices. *Syst Rev*. 2016; 5(1): 192.
12. Wu Y, Tsai W, Tu Y, Yu T. Comparative effectiveness of nonoperative treatments for chronic calcific tendinitis of the shoulder: A systematic review and network meta-analysis of randomized controlled trials. *Arch Phys Med Rehabil*. 2017; 98(8): 1678-1692.
13. Lin MT, Chiang CF, Wu CH, Huang YT, Tu YK, Wang TG. Comparative effectiveness of injection therapies in rotator cuff tendinopathy: A systematic review, pairwise and network meta-analysis of randomized controlled trials. *Arch Phys Med Rehabil*. 2019; 100(2): 336-349.
14. Dong W, Goost H, Lin XB, Burger C, Paul C, Wang ZL, Kong FL, Welle K, Jiang ZC, Kabir K. Injection therapies for lateral epicondylalgia: A systematic review and Bayesian network meta-analysis. *Br J Sports Med*. 2016; 50(15): 900-908.

15. Krogh TP, Bartels EM, Ellingsen T, Stengaard-Pedersen K, Buchbinder R, Fredberg U, Bliddal H, Christensen R. Comparative effectiveness of injection therapies in lateral epicondylitis: A systematic review and network meta-analysis of randomized controlled trials. *Am J Sports Med.* 2013; 41(6): 1435-1446.
16. Chen PC, Wu KT, Chou WY, Huang YC, Wang LY, Yang TH, Siu KK, Tu YK. Comparative effectiveness of different nonsurgical treatments for patellar tendinopathy: A systematic review and network meta-analysis. *Arthroscopy.* 2019; 35(11): 3117-3131
17. van der Vlist AC, Winters M, Weir A, Arden CL, Welton NJ, Caldwell DM, Verhaar JAN, de Vos RJ. Which treatment is most effective for patients with Achilles tendinopathy? A living systematic review with network meta-analysis of 29 randomised controlled trials. *Br J Sports Med.* 2021; 55(5): 249-256.
18. Rhim HC, Kim MS, Choi S, Tenforde AS. Comparative efficacy and tolerability of nonsurgical therapies for the treatment of midportion Achilles tendinopathy. A systematic review with network meta-analysis. *Orthop J Sports Med.* 2020; 8(7): 2325967120930567.
- 19 Alexander, L.A., Harrison, I., Moss, R.A., Greig, L., Shim, J., Pavlova, A.V., Parkinson, E., Maclean, C., Morrissey, D., Swinton P.A., Brandie, D., Mitchell, L., Brown, V.T. & Cooper, K. Exercise therapy for tendinopathy: A scoping review mapping interventions and outcomes. Pre-print available from SportRxiv. 2021.
20. Hutton B, Salanti G, Caldwell DM, Chaimani A, Schmid CH, Cameron C, Ioannidis JPA, Straus S, Thorlund K, Jansen JP, Mulrow C, Catalá-López F, Gøtzsche PC, Dickersin K, Boutron I, Altman DG, Moher D. The PRISMA extension statement for reporting of systematic reviews incorporating network meta-analyses of health care interventions: Checklists and explanations. *Ann Intern Med.* 2015; 26(11): 777-784.
21. Brignardello-Petersen R, Florez ID, Izcovich A, Santesso N, Hazelwood G, Alhazanni W, Yepes-Nuñez JJ, Tomlinson G, Schünemann HJ, Guyatt GH. Grade approach to drawing conclusions from a network meta-analysis using a minimally contextualised framework. *BMJ.* 2020; 371:m3900.
22. Swinton, P.A. Shim, J. Pavlova, A.V. Moss, R.A. MacLean, C. Brandie, D. Mitchell, L. Greig, L. Parkinson, E. Brown, V.T. Morrissey, D. Alexander, L. Cooper, K. Which treatments are most effective for common tendinopathies? A systematic review and network meta-analysis protocol. Pre-print available from SportRxiv. 2021. Doi: 10.31236/osf.io/mx5pv
23. Swinton, P.A. Shim, J. Pavlova, A.V. Moss, R.A. MacLean, C. Brandie, D. Mitchell, L. Greig, L. Parkinson, E. Brown, V.T. Morrissey, D. Alexander, L. Cooper, K. Empirically derived guidelines for interpreting the effectiveness of exercise therapy for tendinopathies: A meta-analysis. Pre-print available from SportRxiv. 2022. Doi: 10.51224/SRXIV.111
24. Alfredson H, Pietilä T, Jonsson P, Lorentzon R. Heavy-load eccentric calf muscle training for the treatment of chronic Achilles tendinosis. *Am J Sports Med.* 1998; 26(3): 360-366.
25. Higgins JP, Altman DG, Gøtzsche PC, Jüni P, Moher D, Oxman AD, Savović J, Schulz KF, Sterne JAC. The Cochrane collaboration's tool for assessing risk of bias in randomised trials. *BMJ.* 2011; 343:d5928.
26. Challoumas D, Millar NL. Risk of bias in systematic reviews of tendinopathy management: Are we comparing apples with oranges? *Transl Sports Med.* 2021; 4(1): 21-37.
27. Marshall IJ, Kuiper J, Wallace, BC. RobotReviewer: evaluation of a system for automatically assessing bias in clinical trials. *J Am Med Inform Assoc.* 2016; 23(1): 193-201.
28. Morris S. Estimating effect sizes from pretest-posttest-control group designs. *Organ Res Methods.* 2008; 11(2): 364-386.

29. Salanti G, Ades AE, Ioannidis JP. Graphical methods and numerical summaries for presenting results from multiple-treatment meta-analysis: an overview and tutorial. *J Clin Epidemiol*. 2011; 64(2): 163–171.
 30. Lunn DJ, Thomas A, Best N, Spiegelhalter D. WinBUGS—a Bayesian modelling framework: concepts, structure, and extensibility. *Stat Comput*. 2000; 10(1): 325–337.
 31. Sturtz S, Ligges U, Gelman AE. R2WinBUGS: a package for running WinBUGS from R. *J Stat Softw*. 2005; 12(3): 1-16.
 32. Busch AJ, Webber SC, Richards RS, Bidonde J, Schachter CL, Schafer LA, Danyliw A, Sawant A, Bello-Haas VD, Rader T, Overend TJ. Resistance exercise training for fibromyalgia. *Cochrane Database Syst Rev*. 2013; (12):CD010884.
 33. Sogaard K, Jull G. Therapeutic exercise for prevention, treatment and rehabilitation of musculoskeletal pain and function. *Man Ther*. 2015; 20(5): 631-632.
 34. National Institute for Health Research Dissemination Centre. Moving forward: Physiotherapy for musculoskeletal health and wellbeing. 2018; London, UK.
 35. National Institute for Health and Care Excellence. Chronic pain (primary and secondary) in over 16s: assessment of all chronic pain and management of chronic primary pain. 2021; NG193.
 36. Irby A, Gutierrez J, Chamberlin C, Thomas SJ, Rosen AB. Clinical management of tendinopathy: A systematic review of systematic reviews evaluating the effectiveness of tendinopathy treatments. *Scand J Med Sci Sports*. 2020; 30(10): 1810-1826.
 37. Skivington K, Matthews L, Simpson SA, Craig P, Baird J, Blazeby JM, Boyd KA, Craig N, French DP, McIntosh R, Petticrew M, Rycroft-Malone J, White M, Moore L. A new framework for developing and evaluating complex interventions: Update of Medical Research Council guidance. *BMJ*. 2021; 374: n2061.
 38. Cook JL, Khan KM, Purdam C. Achilles tendinopathy. *Man Ther*. 2002; 7(3): 121-130.
 39. Kannus P. Etiology and pathophysiology of chronic tendon disorders in sports. *Scand J Med Sci Sports* 1997; 7(2): 78-85.
 40. Alfredson H. The chronic painful achilles and patellar tendon: Research on basic biology and treatment. *Scand J Med Sci Sports* 2005; 15(4): 252-259.
 41. Rees JD. The role of inflammatory cells in tendinopathy: is the picture getting any clearer? *Br J Sports Med*. 2016; 50(4): 201-202.
 42. Hill CE, Stanton R, Heales LJ, Kean CO. Therapeutic tape use for lateral elbow tendinopathy: A survey of Australian healthcare practitioners. *Musculoskelet Sci Pract*. 2020; 48: 102160.
 43. Shahabi S, Lankarani KB, Heydari ST, Jalali M, Ghahramani S, Kamyab B, Tabrizi R, Hosseinabadi M. The effects of counterforce brace on pain in subjects with lateral elbow tendinopathy: A systematic review and meta-analysis of randomized controlled trials. *Prosthet Orthot Int*. 2020; 44(5):341-354.
 44. Heales LJ, McClintock SR, Maynard S, Lems CJ, Rose JA, Hill C, Kean CO, Obst SO. Evaluating the immediate effect of forearm and wrist orthoses on pain and function in individuals with lateral elbow tendinopathy: A systematic review. *Musculoskelet Sci Pract*. 2020; 47:102147.
 45. Knobloch K, Schreibmueller L, Longo UG, Vogt PM. 2008. Eccentric exercises for the management of tendinopathy of the main body of the Achilles tendon with or without an AirHeel™ Brace: A randomized controlled trial. B: effects of compliance. *Disabil Rehabil*. 2008; 30(20-22): 1692–1696.
- Swinton, Shim, Pavlova, Moss, Maclean, Greig, Parkinson, Morrissey, Alexander, Cooper (2021)

46. Fredberg U, Bolvig L, Pfeiffer-Jensen M, Clemmensen D, Jakobsen BW, Stengaard-Pedersen K. Ultrasonography as a tool for diagnosis, guidance of local steroid injection and, together with pressure algometry, monitoring of the treatment of athletes with chronic jumper's knee and Achilles tendinitis: A randomized, double-blind, placebo-controlled study. *Scand J Rheumatol.* 2004; 33(2): 94-101.
47. Barnett J, Bernacki MN, Kainer JL, Smith HN, Zaharoff AM, Subramanian SK. The effects of regenerative injection therapy compared to corticosteroids for the treatment of lateral Epicondylitis: a systematic review and meta-analysis. *Arch Physiother.* 2019; 9:12.
48. Korakakis V, Whiteley R, Tzavara A, Malliaropoulos N. The effectiveness of extracorporeal shockwave therapy in common lower limb conditions: A systematic review including quantification of patient-rated pain reduction. *Br J Sports Med.* 2018; 52(6): 387-407.
49. Wang CJ, Wang FS, Yang KD, Weng LH, Hsu CC, Huang CS, Yang LC. Shock wave therapy induces neovascularization at the tendon-bone junction. A study in rabbits. *J Orthop Res.* 2003; 21(6): 984-989.
50. Chung B, Wiley PJ. Extracorporeal shockwave therapy: A review. *Sports Med.* 2002; 32(13): 851-865.

Supplementary file 1: PRISMA NMA checklist of items to include when reporting a systematic review involving a network meta-analysis

Section/Topic	Item #	Checklist Item	Reported on Page #
TITLE			
Title	1	Identify the report as a systematic review <i>incorporating a network meta-analysis (or related form of meta-analysis)</i> .	1
ABSTRACT			
Structured summary	2	Provide a structured summary including, as applicable: Background: main objectives Methods: data sources; study eligibility criteria, participants, and interventions; study appraisal; and <i>synthesis methods, such as network meta-analysis</i> . Results: number of studies and participants identified; summary estimates with corresponding confidence/credible intervals; <i>treatment rankings may also be discussed. Authors may choose to summarize pairwise comparisons against a chosen treatment included in their analyses for brevity.</i> Discussion/Conclusions: limitations; conclusions and implications of findings. Other: primary source of funding; systematic review registration number with registry name.	2
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of what is already known, <i>including mention of why a network meta-analysis has been conducted</i> .	3-4
Objectives	4	Provide an explicit statement of questions being addressed, with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	5
METHODS			
Protocol and registration	5	Indicate whether a review protocol exists and if and where it can be accessed (e.g., Web address); and, if available, provide registration information, including registration number.	5
Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale. <i>Clearly describe eligible treatments included in the treatment network, and note whether any have been clustered or merged into the same node (with justification).</i>	6-7
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	6-7
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	SF4
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	5
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.	5

Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made.	SF5
Geometry of the network	S1	Describe methods used to explore the geometry of the treatment network under study and potential biases related to it. This should include how the evidence base has been graphically summarized for presentation, and what characteristics were compiled and used to describe the evidence base to readers.	7
Risk of bias within individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	7
Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means). <i>Also describe the use of additional summary measures assessed, such as treatment rankings and surface under the cumulative ranking curve (SUCRA) values, as well as modified approaches used to present summary findings from meta-analyses.</i>	6-7
Planned methods of analysis	14	Describe the methods of handling data and combining results of studies for each network meta-analysis. This should include, but not be limited to: <ul style="list-style-type: none"> • <i>Handling of multi-arm trials;</i> • <i>Selection of variance structure;</i> • <i>Selection of prior distributions in Bayesian analyses; and</i> • <i>Assessment of model fit.</i> 	7
Assessment of Inconsistency	S2	Describe the statistical methods used to evaluate the agreement of direct and indirect evidence in the treatment network(s) studied. Describe efforts taken to address its presence when found.	7
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).	7
Additional analyses	16	Describe methods of additional analyses if done, indicating which were pre-specified. This may include, but not be limited to, the following: <ul style="list-style-type: none"> • Sensitivity or subgroup analyses; • Meta-regression analyses; • <i>Alternative formulations of the treatment network; and</i> • <i>Use of alternative prior distributions for Bayesian analyses (if applicable).</i> 	7

RESULTS†

Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	8
Presentation of network structure	S3	Provide a network graph of the included studies to enable visualization of the geometry of the treatment network.	10,13
Summary of network geometry	S4	Provide a brief overview of characteristics of the treatment network. This may include commentary on the abundance of trials and randomized patients for the different interventions and pairwise comparisons in the network, gaps of evidence in the treatment network, and potential biases reflected by the network structure.	11

Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.	11
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment.	SF7
Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: 1) simple summary data for each intervention group, and 2) effect estimates and confidence intervals. <i>Modified approaches may be needed to deal with information from larger networks.</i>	NI
Synthesis of results	21	Present results of each meta-analysis done, including confidence/credible intervals. <i>In larger networks, authors may focus on comparisons versus a particular comparator (e.g. placebo or standard care), with full findings presented in an appendix. League tables and forest plots may be considered to summarize pairwise comparisons.</i> If additional summary measures were explored (such as treatment rankings), these should also be presented.	11,14
Exploration for inconsistency	S5	Describe results from investigations of inconsistency. This may include such information as measures of model fit to compare consistency and inconsistency models, <i>P</i> values from statistical tests, or summary of inconsistency estimates from different parts of the treatment network.	11,14
Risk of bias across studies	22	Present results of any assessment of risk of bias across studies for the evidence base being studied.	11,14
Results of additional analyses	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression analyses, <i>alternative network geometries studied, alternative choice of prior distributions for Bayesian analyses, and so forth</i>).	NA
DISCUSSION			
Summary of evidence	24	Summarize the main findings, including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy-makers).	14
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review level (e.g., incomplete retrieval of identified research, reporting bias). <i>Comment on the validity of the assumptions, such as transitivity and consistency. Comment on any concerns regarding network geometry (e.g., avoidance of certain comparisons).</i>	16
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	16
FUNDING			
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review. This should also include information regarding whether funding has been received from manufacturers of treatments in the network and/or whether some of the authors are content experts with professional conflicts of interest that could affect use of treatments in the network.	19

PICOS = population, intervention, comparators, outcomes, study design.

* Text in italics indicateS wording specific to reporting of network meta-analyses that has been added to guidance from the PRISMA statement.

† Authors may wish to plan for use of appendices to present all relevant information in full detail for items in this section.

Supplementary File 2: Outcome domains and example outcomes included in review.

Domain	ICON Definition	Example Tools
Disability	Composite scores of a mix of patient-rated pain & disability due to the pain, usually relating to tendon-specific activities/tasks	VISA scales; DASH; quick DASH; SPADI; Patient-rated tennis-elbow evaluation questionnaire; Constant Murley Score; WORC (Western Ontario Rotator Cuff Index); AOFAS (American Orthopaedic Foot & Ankle Society); Roles and Maudsley score; ASES (American Shoulder & Elbow Surgeons Index; Tegner activity score; Lysholm knee scale; Pain free function questionnaire; Ankle activity score; Subjective elbow Value (SEV); Placzek score; Shoulder disability questionnaire; International Knee Documentation Committee form (IKDC); Penn Shoulder score (university of Pennsylvania shoulder score) (PSS); Brief pain inventory (BPI); UCLA Shoulder Rating Scale; FILLA - functional index of leg and lower limb; Neer Shoulder Score; Nirschl phase rating scale; American Shoulder and Elbow Surgeon's (MASES) questionnaire; Mayo Elbow Performance Score (MEPS); Shoulder rating questionnaire (SRQ)
Function	Patient-rated level of function (and not referring to the intensity of their pain).	Patient-specific functional scale
Pain	Pain on loading/activity: Patient reported intensity of pain performing a task that loads the tendon.	VAS; NRS; Pain experience scale
	Pain over a specified time: Patient-reported pain intensity over period of time e.g. morning/night/24-hours/1-week.	VAS; NRS Painful days in 3 months
	Pain without further specification: Patient asked about pain levels without reference to activity or timeframe.	VAS; NRS; Borg CR10 Scale; Pain status
Physical function capacity	Quantitative measures of physical tasks (e.g. hops, times walk, single leg squat) includes muscle strength	Counter movement jump; One-leg triple hop; Single-leg decline squat; Muscle strength measured by dynamometry (hand-held, isokinetic); Repetition maximum; Manual muscle testing.
Quality of Life	General wellbeing	EQ5D; EQ3D; SF-36 or SF-12; Assessment of Quality of Life (AQoL); Nottingham Health Profile; Gothenburg QoL Instrument
Range of Motion (Shoulder only)	Active or passive range of motion in specified plane, measured in degrees.	Hand-held goniometer; inclinometer

Supplementary File 3: Definitions used to define broad and more specific treatment classes.

Broad treatment class	Definition	More specific treatment class	Definition
Exercise only	Exercise therapy is defined as a regimen or program of physical activities specifically designed and prescribed to correct impairments, restore musculoskeletal function, and/or maintain a state of wellbeing.	Same as broad treatment class	Same as broad treatment class
Non-active (placebo, sham, wait and see)	Includes any appropriate inactive treatment such as waiting list control, sham shockwave, sham laser, sham taping or true placebo.	Same as broad treatment class	Same as broad treatment class
Non-exercise only	Active treatments used to treat tendinopathy that do not meet the criteria to be considered exercise.	Electrotherapy	Modality that delivers therapeutic levels of physical energy into a biologic system e.g. soft tissue. Includes shockwave, laser and other systems.
		Biomechanics	Treatment using external devices that immobilises (e.g. splinting) or alters the kinematics/kinetics of the limb (e.g. taping, bracing and orthotics).
		Manual-therapy	Manual therapy is the skilled application of “hands-on” techniques to treat soft tissues and joint structures for the purpose of improving pain, increasing range of motion, stimulating tissue repair response, and/or improving function.
		Injection therapy	Injection therapy for tendinopathy typically involves direct administration of a pharmacologically active drug, or combination of drugs using a syringe and needle or equivalent. It may or may not be image-guided. Includes Autologous, drug, and volumetric types.
		Surgery	Any relevant surgical intervention for tendinopathy including minimally invasive peritendinous and open intra-tendinous.

Exercise and non-exercise	Treatment comprising multiple components which collectively meet both exercise and non-exercise criteria	Same as broad treatment class	Same as broad treatment class
---------------------------	--	-------------------------------	-------------------------------

Supplementary file 4: Search terms and results for MEDLINE search

MEDLINE (EBSCoHost)

Search conducted on 21/01/21.

Search	Query	Records retrieved
#1	MH exercise OR AB exercis* OR MH “isometric contraction” OR MH rehabilitation OR TX eccentric OR TX concentric OR TX “heavy slow resistance” OR TX isokinetic	362,722
#2	MH tendinopathy OR MH “shoulder injuries” OR MH tendons OR MH “tendon injuries” OR TX tendin* OR TX tendon* OR MH bursitis OR AB bursitis OR MH “posterior tibial tendon dysfunction” OR MH “shoulder impingement syndrome” OR AB “greater trochanteric pain syndrome”	96,490
#3	#1 AND #2	4,363

Limited to 1998 to present

Supplementary file 5: Extraction codebook

Column	Heading	Description
Study Details	A	Initials Reviewer
	B	Covidence Identifier
	C	Author
	D	Year
	E	Title
	F	Country
	G	Journal
	H	Aims/Purpose
	I	Tendinopathy type
	J	Study Design
	K	Age Mean
	L	Age SD
	M	Baseline Total N
	N	Training Status Description
	O	Training Status Code
	P	Sex
	Q	BMI Mean
	R	BMI SD
	S	Symptom Severity Mean
	T	Symptom Severity SD
	U	Symptom Duration Mean (Months)
	V	Symptom Duration SD (Months)
	W	Population Comments
Outcomes	X	Outcome Category
	Y	Outcome Tool
	Z	Reflection
	AA	Measurement Time (Weeks)
Intervention	AB	Dominant Broad Treatment Class
	AC	Total Broad Treatment class
	AD	Dominant Specific Treatment Class
	AE	Total Specific Treatment Class
	AF	Intervention N
	AG	Intervention Total Duration
	AH	Intervention Adherence %
	AI	Intervention Location
	AJ	Intervention Volume
	AK	Intervention Volume Category

	AL	Intervention Volume Comments	Any additional information relevant.
	AM	Intervention Intensity	Numerical value describing intensity
	AN	Intervention Intensity Category	1 = Absolute; 2 = Relative
	AO	Intervention Frequency	Number of sessions per week. Where there is progression, average value is to be entered.
	AP	Intervention Frequency Comments	Any additional information relevant.
	AQ	Intervention Progression	Multiple themes to be selected as required 1 = No progression; 2 = NR; 3 = Progression volume; 4 = Progression intensity; 5 = Progression frequency; 6 = Progression specificity; 7 = Progression capacity; 8 = Other
	AR	Intervention Progression Comments	Any additional information relevant.
Data	AS	Intervention Baseline Mean	Baseline mean for exercise therapy
	AT	Intervention Baseline SD	Baseline standard deviation for exercise therapy
	AU	Intervention Measurement Mean	Mean of outcome for exercise therapy at stated time point
	AV	Intervention Measurement SD	Standard deviation of outcome for exercise therapy at stated time point
	AW	Control Baseline Mean	Baseline mean for control
	AX	Control Baseline SD	Baseline standard deviation for control
	AY	Control Measurement Mean	Mean of outcome for control at stated time point
	AZ	Control Measurement SD	Standard deviation of outcome for control at stated time point
	BA	Measurement Comments	State if a different value has been entered for means (e.g. median), a different value for standard deviations (e.g. standard error, IQR, percentiles, distance from mean to upper bound). Provide the relevant statistic (width of CI's, width of percentiles). Also state if data has extracted by digitization

Supplementary file 6: Reference list of included studies

1. Abat F, Sanchez-Sanchez J, Martin-Nogueras A, Calvo-Arenillas J, Yajeya J, Mendez-Sanchez R, et al. Randomized controlled trial comparing the effectiveness of the ultrasound-guided galvanic electrolysis technique (USGET) versus conventional electro-physiotherapeutic treatment on patellar tendinopathy. 2016;3(1).
2. Aceituno-Gómez J, Avendaño-Coy J, Gómez-Soriano J, García-Madero VM, Ávila-Martín G, Serrano-Muñoz D, et al. Efficacy of high-intensity laser therapy in subacromial impingement syndrome: a three-month follow-up controlled clinical trial. *Clinical rehabilitation*. 2019;33(5):894-903.
3. Akkaya N, Akkaya S, Gungor HR, Yaşar G, Atalay NS, Sahin F. Effects of weighted and un-weighted pendulum exercises on ultrasonographic acromiohumeral distance in patients with subacromial impingement syndrome. *Journal of back and musculoskeletal rehabilitation*. 2017;30(2):221-8.
4. Aktas I, Akgun K, Cakmak B. Therapeutic effect of pulsed electromagnetic field in conservative treatment of subacromial impingement syndrome. *Clinical rheumatology*. 2007;26(8):1234-9.
5. Akyol Y, Ulus Y, Durmus D, Canturk F, Bilgici A, Kuru O, et al. Effectiveness of microwave diathermy on pain, functional capacity, muscle strength, quality of life, and depression in patients with subacromial impingement syndrome: a randomized placebo-controlled clinical study. *Rheumatology international*. 2012;32(10):3007-16.
6. Alfredson H, Pietilä T, Jonsson P, Lorentzon R. Heavy-load eccentric calf muscle training for the treatment of chronic Achilles tendinosis. *The American Journal of Sports Medicine*. 1998;26(3):360-6.
7. Alfredson H, Nordström P, Pietilä T, Lorentzon R. Bone mass in the calcaneus after heavy loaded eccentric calf-muscle training in recreational athletes with chronic achilles tendinosis. *Calcified tissue international*. 1999;64(5):450-5.
8. Arias-Buría JL, Truyols-Domínguez S, Valero-Alcaide R, Salom-Moreno J, Atín-Arratibel MA, Fernández-de-las-Peñas C. Ultrasound-guided percutaneous electrolysis and eccentric exercises for subacromial pain syndrome: a randomized clinical trial. *Evidence-Based Complementary and Alternative Medicine*. 2015;2015.
9. Arias-Buría JL, Fernández-de-Las-Peñas C, Palacios-Ceña M, Koppenhaver SL, Salom-Moreno J. Exercises and dry needling for subacromial pain syndrome: A randomized parallel-group trial. *The Journal of Pain*. 2017;18(1):11-8.
10. Bae YH, Lee GC, Shin WS, Kim TH, Lee SM. Effect of motor control and strengthening exercises on pain, function, strength and the range of motion of patients with shoulder impingement, syndrome. *Journal of Physical Therapy Science*. 2011;23(4):687-92.
11. Bagcier F, Yilmaz N. The impact of extracorporeal shock wave therapy and dry needling combination on the Pain, Grip strength and functionality in patients diagnosed with lateral epicondylitis. *Turk Osteoporoz Dergisi*. 2019;25(2):65-71.
12. Bahr R, Fossan B, Løken S, Engebretsen L. Surgical treatment compared with eccentric training for patellar tendinopathy (jumper's knee): a randomized, controlled trial. *JBJS*. 2006;88(8):1689-98.
13. Balias R, Álvarez G, Baró F, Jiménez F, Pedret C, Costa E, et al. A 3-arm randomized trial for Achilles tendinopathy: eccentric training, eccentric training plus a dietary supplement containing mucopolysaccharides, or passive stretching plus a dietary supplement containing mucopolysaccharides. *Current Therapeutic Research*. 2016;78:1-7.
14. Bang MD, Deyle GD. Comparison of supervised exercise with and without manual physical therapy for patients with shoulder impingement syndrome. *Journal of Orthopaedic & Sports Physical Therapy*. 2000;30(3):126-37.
15. Başkurt Z, Başkurt F, Gelecek N, Özkan MH. The effectiveness of scapular stabilization exercise in the patients with subacromial impingement syndrome. *Journal of back and musculoskeletal rehabilitation*. 2011;24(3):173-9.
16. Beaudreuil J, Lasbleiz S, Yelnik A, Bardin T, Orcel P. Effect of dynamic humeral centering on painful active elevation of the arm in subacromial impingement syndrome: a randomized trial. *Annals of Physical and Rehabilitation Medicine*. 2012;55:e158-e61.

17. Bek N, Simşek IE, Erel S, Yakut Y, Uygur F. Home-based general versus center-based selective rehabilitation in patients with posterior tibial tendon dysfunction. *Acta orthopaedica et traumatologica turcica*. 2012;46(4):286-92.
18. Bell KJ, Fulcher ML, Rowlands DS, Kerse N. Impact of autologous blood injections in treatment of mid-portion Achilles tendinopathy: double blind randomised controlled trial. *BMJ (Clinical research ed)*. 2013;346.
19. Bennell K, Wee E, Coburn S, Green S, Harris A, Staples M, et al. Efficacy of standardised manual therapy and home exercise programme for chronic rotator cuff disease: randomised placebo controlled trial. *British medical journal*. 2010;(7763)(7763).
20. Beyer R, Kongsgaard M, Hougs Kjær B, Øhlenschläger T, Kjær M, Magnusson SP. Heavy slow resistance versus eccentric training as treatment for Achilles tendinopathy: a randomized controlled trial. *The American Journal of Sports Medicine*. 2015;43(7):1704-11.
21. Blume C, Wang-Price S, Trudelle-Jackson E, Ortiz A. Comparison of eccentric and concentric exercise interventions in adults with subacromial impingement syndrome. *International journal of sports physical therapy*. 2015;10(4):441-55.
22. Boesen AP, Hansen R, Boesen MI, Malliaras P, Langberg H. Effect of High-Volume Injection, Platelet-Rich Plasma, and Sham Treatment in Chronic Midportion Achilles Tendinopathy: a Randomized Double-Blinded Prospective Study. *The American Journal of Sports Medicine*. 2017;45(9).
23. Boesen AP, Langberg H, Hansen R, Malliaras P, Boesen MI. High volume injection with and without corticosteroid in chronic midportion achilles tendinopathy. *Scandinavian Journal of Medicine & Science in Sports*. 2019;29(8):1223-31.
24. Boudreau N, Gaudreault N, Roy JS, Bédard S, Balg F. The Addition of Glenohumeral Adductor Coactivation to a Rotator Cuff Exercise Program for Rotator Cuff Tendinopathy: a Single-Blind Randomized Controlled Trial. *The Journal of orthopaedic and sports physical therapy*. 2019;49(3):126-35.
25. Breda SJ, Oei EHG, Zwerver J, Visser E, Waarsing E, Krestin GP, et al. Effectiveness of progressive tendon-loading exercise therapy in patients with patellar tendinopathy: a randomised clinical trial. *British journal of sports medicine*. 2020.
26. Brox JI, Gjengedal E, Uppheim G, Bøhmer AS, Brevik JI, Ljunggren AE, et al. Arthroscopic surgery versus supervised exercises in patients with rotator cuff disease (stage II impingement syndrome): a prospective, randomized, controlled study in 125 patients with a 2 1/2-year follow-up. *Journal of shoulder and elbow surgery*. 1999;8(2):102-11.
27. Buyuksireci DE, Turk AC. Evaluation of the effectiveness of dexamethasone iontophoresis in patients with subacromial impingement syndrome. *Journal of Orthopaedic Science*. 2021;26(5):786-91.
28. Cacchio A, Rompe JD, Furia JP, Susi P, Santilli V, De Paulis F. Shockwave therapy for the treatment of chronic proximal hamstring tendinopathy in professional athletes. *The American Journal of Sports Medicine*. 2011;39(1).
29. Calis HT, Berberoglu N, Calis M. Are ultrasound, laser and exercise superior to each other in the treatment of subacromial impingement syndrome? A randomized clinical trial. *European journal of physical and rehabilitation medicine*. 2011;47(3):375-80.
30. Celik D, Akyüz G, Yeldan I. [Comparison of the effects of two different exercise programs on pain in subacromial impingement syndrome]. Turkey: Turkish Association of Orthopaedics and Traumatology with publishing services by Elsevier B.V; 2009. p. 504-9.
31. Celik D, Atalar AC, Güçlü A, Demirhan M. [The contribution of subacromial injection to the conservative treatment of impingement syndrome]. *Acta orthopaedica et traumatologica turcica*. 2009;43(4):331-5.
32. Celik D, Anaforoglu Kulunkoglu B. Photobiomodulation Therapy Versus Extracorporeal Shock Wave Therapy in the Treatment of Lateral Epicondylitis. *Photobiomodulation, photomedicine, and laser surgery*. 2019;37(5):269-75.
33. Cha J-Y, Kim J-H, Hong J, Choi Y-T, Kim M-H, Cho J-H, et al. A 12-week rehabilitation program improves body composition, pain sensation, and internal/external torques of baseball pitchers with shoulder impingement symptom. *Journal of Exercise Rehabilitation*. 2014;10(1):35-44.

34. Chaconas EJ, Kolber MJ, Hanney WJ, Daugherty ML, Wilson SH, Sheets C. Shoulder external rotator eccentric training versus general shoulder exercise for subacromial pain syndrome: a randomized controlled trial. *International Journal of Sports Physical Therapy*. 2017;12(7):1121-33.
35. Chary-Valckenaere I, Loeuille D, Jay N, Kohler F, Tamisier JN, Roques CF, et al. Spa therapy together with supervised self-mobilisation improves pain, function and quality of life in patients with chronic shoulder pain: a single-blind randomised controlled trial. *International journal of biometeorology*. 2018;62(6):1003-14.
36. Chen TW, Huei Su J, Lin TY, Lin CW, Chou PS. Effects of Eccentric Exercise and Extracorporeal Shock Wave Therapy on Rehabilitation of Patients with Noncalcific Rotator Cuff Tendinopathy. *Clin Res Foot Ankle*. 2017;5(222):2.
37. Cheng AS, Hung L. Randomized controlled trial of workplace-based rehabilitation for work-related rotator cuff disorder. *Journal of Occupational Rehabilitation*. 2007;17(3):487-503.
38. Cho S-I, Shin Y-A. Effect of rehabilitation and prolotherapy on pain and functional performance in patients with chronic patellar tendinopathy. 2017;176(6).
39. Chung B, Wiley JP. Effectiveness of extracorporeal shock wave therapy in the treatment of previously untreated lateral epicondylitis: A randomized controlled trial. *American Journal of Sports Medicine*. 2004;32(7):1660-7.
40. Clarke AW, Ahmad M, Curtis M, Connell DA. Lateral elbow tendinopathy: correlation of ultrasound findings with pain and functional disability. *The American Journal of Sports Medicine*. 2010;38(6):1209-14.
41. Conroy DE, Hayes KW. The effect of joint mobilization as a component of comprehensive treatment for primary shoulder impingement syndrome. *The Journal of orthopaedic and sports physical therapy*. 1998;28(1):3-14.
42. Coombes BK, Bisset L, Brooks P, Khan A, Vicenzino B. Effect of Corticosteroid Injection, Physiotherapy, or Both on Clinical Outcomes in Patients With Unilateral Lateral Epicondylalgia: A Randomized Controlled Trial. *JAMA : the journal of the American Medical Association*. 2013;309(5):461-9.
43. Croisier JL, Foidart-Dessalle M, Tinant F, Crielaard JM, Forthomme B. An isokinetic eccentric programme for the management of chronic lateral epicondylar tendinopathy. *British journal of sports medicine*. 2007;41(4).
44. de Jonge S, Tol JL, de Vos R-J, van Schie HTM, Verhaar JAN. One-year follow-up of a randomised controlled trial on added splinting to eccentric exercises in chronic midportion Achilles tendinopathy. *bjsm.bmj.com*. 2008.
45. de Miguel Valtierra L, Salom Moreno J, Fernández-de-Las-Peñas C, Cleland JA, Arias-Burúa JL. Ultrasound-Guided Application of Percutaneous Electrolysis as an Adjunct to Exercise and Manual Therapy for Subacromial Pain Syndrome: A Randomized Clinical Trial. *The journal of pain : official journal of the American Pain Society*. 2018;19(10):1201-10.
46. Oliveira L. It is not always what it appears. *Clinical Journal of Sport Medicine*. 2022;32(2):e204-e5.
47. De Vos RJ, Weir A, Visser RJA, de Winter T, Tol JL. The additional value of a night splint to eccentric exercises in chronic midportion Achilles tendinopathy: a randomised controlled trial. *British journal of sports medicine*. 2007;41(7).
48. de Vos RJ, Weir A, van Schie HT, Bier-Zeinstra S, Verhaar JA, Weinans H, et al. Platelet-rich plasma injection for chronic Achilles tendinopathy. *Journal - American Medical Association*. 2010;303(2):144-9.
49. DeJaco B, Habets B, van Loon C, van Grinsven S, van Cingel R. Eccentric versus conventional exercise therapy in patients with rotator cuff tendinopathy: a randomized, single blinded, clinical trial. *Knee Surgery, Sports Traumatology, Arthroscopy*. 2017;25(7):2051-9.
50. Devereaux M, Velanoski KQ, Pennings A, Elmaraghy A. Short-Term Effectiveness of Precut Kinesiology Tape Versus an NSAID as Adjuvant Treatment to Exercise for Subacromial Impingement: A Randomized Controlled Trial. *Clinical journal of sport medicine : official journal of the Canadian Academy of Sport Medicine*. 2016;26(1):24-32.
51. Dimitrios S, Pantelis M, Kalliopi S. Comparing the effects of eccentric training with eccentric training and static stretching exercises in the treatment of patellar tendinopathy. A controlled clinical trial. *Clinical rehabilitation*. 2012;26(5):423-30.

52. Stasinopoulos D, Manias P. Comparing two eccentric exercise programmes for the management of Achilles tendinopathy. A pilot trial. *Journal of Bodywork and Movement Therapies*. 2013;17(3):309-15.
53. Dogan SK, Ay S, Evcik D. The effectiveness of low laser therapy in subacromial impingement syndrome: a randomized placebo controlled double-blind prospective study. *Clinics (Sao Paulo, Brazil)*. 2010;65(10):1019-22.
54. Dupuis F, Barrett E, Dubé M-O, McCreesh KM, Lewis JS, Roy J-S. Cryotherapy or gradual reloading exercises in acute presentations of rotator cuff tendinopathy: a randomised controlled trial. *BMJ open sport & exercise medicine*. 2018;4(1).
55. Eken Gedik D, Dost Surucu G, Yildirim A, Karabiber M. Effectiveness of autologous blood injection in the treatment of lateral epicondylitis: Randomized clinical study. *Duzce Medical Journal*. 2016;18(1):1-7.
56. Ellegaard K, Christensen R, Rosager S, Bartholdy C, Torp-Pedersen S, Bandholm T, et al. Exercise therapy after ultrasound-guided corticosteroid injections in patients with subacromial pain syndrome: a randomized controlled trial. *Arthritis research & therapy*. 2016;18(1):129.
57. Engebretsen K, Grotle M, Bautz-Holter E, Sandvik L, Juel NG, Ekeberg OM, et al. Radial extracorporeal shockwave treatment compared with supervised exercises in patients with subacromial pain syndrome: single blind randomised study. *BMJ (Clinical research ed)*. 2009;339.
58. Engebretsen K, Grotle M, Bautz-Holter E, Ekeberg OM, Juel NG, Brox JI. Supervised exercises compared with radial extracorporeal shock-wave therapy for subacromial shoulder pain: 1-year results of a single-blind randomized controlled trial. *Physical Therapy*. 2011;91(1):37-47.
59. Faria APGS, Martinho KO, Silva EB, Dantas EHM. The Effects of Different Resources Fotottherapeutic on the Pain in Individuals Bearers of Syndrome of the Impact of the Shoulder. *Fitness & Performance Journal (Online Edition)*. 2006;5(6):54-61.
60. Belley AF, Mercier C, Bastien M, Léonard G, Gaudreault N, Roy J-S. Anodal Transcranial Direct-Current Stimulation to Enhance Rehabilitation in Individuals With Rotator Cuff Tendinopathy: A Triple-Blind Randomized Controlled Trial. *Journal of Orthopaedic & Sports Physical Therapy*. 2018;48(7):541-51.
61. Ganderton C, Semciw A, Cook J, Moreira E, Pizzari T. Gluteal loading versus sham exercises to improve pain and dysfunction in postmenopausal women with greater trochanteric pain syndrome: a randomized controlled trial. *Journal of Women's Health*. 2018;27(6):815-29.
62. García I, Lobo C, López E, Serván JL, Tenías JM. Comparative effectiveness of ultrasonophoresis and iontophoresis in impingement syndrome: a double-blind, randomized, placebo controlled trial. *Clinical rehabilitation*. 2016;30(4):347-58.
63. Gatz M, Betsch M, Dirrichs T, Schrading S, Tingart M, Michalik R, et al. Eccentric and Isometric Exercises in Achilles Tendinopathy Evaluated by the VISA-A Score and Shear Wave Elastography. *Sports health*. 2020;1941738119893996.
64. Genc E, Duymaz T. Effectiveness of kinesio taping in bicipital tendinitis treatment: A randomized controlled trial kinesio taping in bicipital tendinitis. *Annals of Clinical and Analytical Medicine*. 2020;11(5):498-503.
65. Giray E, Karali-Bingul D, Akyuz G. The Effectiveness of Kinesiotaping, Sham Taping or Exercises Only in Lateral Epicondylitis Treatment: A Randomized Controlled Study. *Pm&r*. 2019;11(7):681-93.
66. Granviken F, Vasseljen O. Home exercises and supervised exercises are similarly effective for people with subacromial impingement: a randomised trial. *Journal of physiotherapy*. 2015;61(3):135-41.
67. Gürsel YK, Ulus Y, Bilgiç A, Dinçer G, van der Heijden GJM. Adding ultrasound in the management of soft tissue disorders of the shoulder: a randomized placebo-controlled trial. *Physical Therapy*. 2004;84(4):336-43.
68. Hallgren HC, Adolfsson LE, Johansson K, Öberg B, Peterson A, Holmgren TM. Specific exercises for subacromial pain. *Acta orthopaedica*. 2017;88(6):600-5.
69. Hallgren HC, Holmgren T, Öberg B, Johansson K, Adolfsson LE. A specific exercise strategy reduced the need for surgery in subacromial pain patients. *Br J Sports Med*. 2014(1473-0480 (Electronic)).
70. Heron SR, Woby SR, Thompson DP. Comparison of three types of exercise in the treatment of rotator cuff tendinopathy/shoulder impingement syndrome:a randomised control trial assessing. *Physiotherapy* 2017 Jun;103(2):167-173. 2017.

71. Holmgren T, Hallgren HB, Öberg B, Adolfsson L, Johansson K. Effect of specific exercise strategy on need for surgery in patients with subacromial impingement syndrome: randomised controlled study. *Bmj*. 2012;344.
72. Hotta GH, Gomes de Assis Couto A, Cools AM, McQuade KJ, Siriani de Oliveira A. Effects of adding scapular stabilization exercises to a periscapular strengthening exercise program in patients with subacromial pain syndrome: A randomized controlled trial. *Musculoskeletal Science and Practice*. 2020;49.
73. Houck J, Neville C, Tome J, Flemister A. Randomized Controlled Trial Comparing Orthosis Augmented by Either Stretching or Stretching and Strengthening for Stage II Tibialis Posterior Tendon Dysfunction. *Foot and Ankle International*. 2015(1944-7876 (Electronic)).
74. İlhanlı I, Guder N, Gul M. Platelet-Rich Plasma Treatment With Physical Therapy in Chronic Partial Supraspinatus Tears. *Iran Red Crescent Med J*. 2015;17(9):e23732-e.
75. Ingwersen KG, Jensen SL, Sørensen L, Jørgensen HR, Christensen R, Søgaard K, et al. Three months of progressive high-load versus traditional low-load strength training among patients with rotator cuff tendinopathy: primary results from the double-blind randomized controlled RoCTEx trial. *Orthopaedic journal of sports medicine*. 2017;5(8):2325967117723292.
76. Jasnauskaitė-Gedrimė D, Gedrimas D, Karpavičienė A, Skurvydas A. Effect of Visual and Auditory Feedback Exercises on Shoulder Function in Rotator Cuff Tendonitis Patients. *Baltic Journal of Sport & Health Sciences*. 2018;109(2):20-6.
77. Jeong TH, Jk O, Lee HJ. The effect of the combined stretching and strengthening exercise on the clinical symptoms in posterior tibial tendon dysfunction patient. *J Korean Foot Ankle Soc*. 2008;12:47-54.
78. Johansson KM, Adolfsson LE, Foldevi MOM. Effects of acupuncture versus ultrasound in patients with impingement syndrome: randomized clinical trial. *Physical Therapy*. 2005;85(6):490-501.
79. Jonsson P, Alfredson H. Superior results with eccentric compared to concentric quadriceps training in patients with jumper's knee: a prospective randomised study. *Br J Sports Med*. 2005;39.
80. Jonsson P. Eccentric training in the treatment of tendinopathy. Department of Surgical and Perioperative Sciences, Sports Medicine; 2009.
81. Juul-Kristensen B, Larsen CM, Eshoj H, Clemmensen T, Hansen A, Bo Jensen P, et al. Positive effects of neuromuscular shoulder exercises with or without EMG-biofeedback, on pain and function in participants with subacromial pain syndrome - A randomised controlled trial. *Journal of electromyography and kinesiology : official journal of the International Society of Electrophysiological Kinesiology*. 2019;48:161-8.
82. Kachanathu SJ, Alenazi AM, Hafez AR, Algarni AD, Alsubiheen AM. Comparison of the effects of short-duration wrist joint splinting combined with physical therapy and physical therapy alone on the management of patients with lateral epicondylitis. *European journal of physical and rehabilitation medicine*. 2019;55(4):488-93.
83. Kang F-J, Chiu Y-C, Wu S-C, Wang T-G, Yang J-I, Lin J-J. Kinesiology taping with exercise does not provide additional improvement in round shoulder subjects with impingement syndrome: A single-blinded randomized controlled trial. *Physical Therapy in Sport*. 2019;40:99-106.
84. Kedia M, Williams M, Jain L, Barron M, Bird N, Blackwell B, et al. The effects of conventional physical therapy and eccentric strengthening for insertional Achilles tendinopathy. *International journal of sports physical therapy*. 2014;9(4):488.
85. Ketola S, Lehtinen J, Arnala I, Nissinen M, Westenius H, Sintonen H, et al. Does arthroscopic acromioplasty provide any additional value in the treatment of shoulder impingement syndrome?: a two-year randomised controlled trial. *The Journal of bone and joint surgery British volume*. 2009;91(10):1326-34.
86. Ketola S, Lehtinen J, Rousi T, Nissinen M, Huhtala H, Kontinen YT, et al. No evidence of long-term benefits of arthroscopic acromioplasty in the treatment of shoulder impingement syndrome. *Bone & Joint Research*. 2013;2(7):132-9.
87. Kim J, Shin D, Song C. Visual feedback to improve the effects of scapular stabilization exercises on pain intensity, range of motion, strength, and disability in patients with shoulder impingement syndrome. *Medical Science Technology*. 2017;58(1):42-8.

88. Kim SY, Dvir Z, Oh JS. The application of the Neurac technique vs. manual therapy in patients during the acute phase of subacromial impingement syndrome: A randomized single-blinded controlled trial. *Journal of Back and Musculoskeletal Rehabilitation*. 2020;33(4):645-53.
89. Knobloch K, Schreibmueller L, Kraemer R, Jagodzinski M, Richter M, Zeichen J, et al. Eccentric training and an Achilles wrap reduce Achilles tendon capillary blood flow and capillary venous filling pressures and increase tendon oxygen saturation in insertional and midportion tendinopathy: A randomized trial. *American Journal of Sports Medicine*. 2007;35(2):189-96.
90. Knobloch K, Kraemer R, Jagodzinski M, Zeichen J, Melier R, Vogt PM. Eccentric training decreases paratendon capillary blood flow and preserves paratendon oxygen saturation in chronic achilles tendinopathy. *Journal of Orthopaedic and Sports Physical Therapy*. 2007;37(5):269-76.
91. Knobloch K, Schreibmueller L, Longo UG, Vogt PM. Eccentric exercises for the management of tendinopathy of the main body of the Achilles tendon with or without an AirHeel Brace. A randomized controlled trial. B: effects of compliance. *Disability and rehabilitation*. 2008;30(20).
92. Koç C, Kurt EE, Koçak FA, Erdem HR, Konar NM. Does balneotherapy provide additive effects to physical therapy in patients with subacute supraspinatus tendinopathy? A randomized, controlled, single-blind study. *International journal of biometeorology*. 2021;65(2):301-10.
93. Kongsgaard M, Kovanen V, Aagaard P, Doessing S, Hansen P, Laursen AH, et al. Corticosteroid injections, eccentric decline squat training and heavy slow resistance training in patellar tendinopathy. *Scandinavian Journal of Medicine & Science in Sports*. 2009;19(6):790-802.
94. Kromer TO, de Bie RA, Bastiaenen CHG. Effectiveness of physiotherapy and costs in patients with clinical signs of shoulder impingement syndrome: One-year follow-up of a randomized controlled trial. *Journal of Rehabilitation Medicine*. 2014;46(10):1029-36.
95. Kromer TO, de Bie RA, Bastiaenen CHG. Physiotherapy in patients with clinical signs of shoulder impingement syndrome: a randomized controlled trial. *Journal of Rehabilitation Medicine*. 2013;45(5):488-97.
96. Kulig K, Reischl SF, Pomrantz AB, Burnfield JM, Mais-Requejo S, Thordarson DB, et al. Nonsurgical management of posterior tibial tendon dysfunction with orthoses and resistive exercise: a randomized controlled trial. *Physical Therapy*. 2009;89(1):26-37.
97. Land H, Gordon S, Watt K. Effect of manual physiotherapy in homogeneous individuals with subacromial shoulder impingement: A randomized controlled trial. *Physiotherapy Research International : The Journal for Researchers and Clinicians in Physical Therapy*. 2019;24(2).
98. Lee W-C, Ng GY-F, Zhang Z-J, Malliaras P, Masci L, Fu S-N. Changes on Tendon Stiffness and Clinical Outcomes in Athletes Are Associated With Patellar Tendinopathy After Eccentric Exercise. *Clinical journal of sport medicine : official journal of the Canadian Academy of Sport Medicine*. 2017.
99. Lee S, Ko Y, Lee W. Changes in pain, dysfunction, and grip strength of patients with acute lateral epicondylitis caused by frequency of physical therapy: A randomized controlled trial. *Journal of Physical Therapy Science*. 2014;26(7):1037-40.
100. Lee WC. The mechanical, physiological and therapeutic effects of eccentric exercise combined with extracorporeal shockwave therapy in athletes with patellar tendinopathy. 2017.
101. Littlewood C, Bateman M, Brown K, Bury J, Mawson S, May S, et al. A self-managed single exercise programme versus usual physiotherapy treatment for rotator cuff tendinopathy: A randomised controlled trial (the SELF study). *Clinical rehabilitation*. 2016;30(7):686-96.
102. Ludewig PM, Borstad JD. Effects of a home exercise programme on shoulder pain and functional status in construction workers. *Occupational and environmental medicine*. 2003;60(11):841-9.
103. Luginbuhl R, Brunner F, Schneeberger AG. No effect of forearm band and extensor strengthening exercises for the treatment of tennis elbow: a prospective randomised study. *La Chirurgia degli organi di movimento*. 2008;91(1):35-40.
104. Maenhout AG, Mahieu NN, De Muyenck M, De Wilde LF, Cools AM. Does adding heavy load eccentric training to rehabilitation of patients with unilateral subacromial impingement result in better outcome? A randomized, clinical trial. *Knee surgery, sports traumatology, arthroscopy*. 2013;21(5):1158-67.

105. Mafi N, Lorentzon R, Alfredson H. Superior short-term results with eccentric calf muscle training compared to concentric training in a randomized prospective multicenter study on patients with chronic Achilles tendinosis. 2001;9(1).
106. Manias P, Stasinopoulos D. A controlled clinical pilot trial to study the effectiveness of ice as a supplement to the exercise programme for the management of lateral elbow tendinopathy. *British journal of sports medicine*. 2006;40(1):81-5.
107. Martinez-Silvestrini J, Newcomer KL, Gay RE, Schaefer MP, Kortebein P, Arendt KW. Chronic lateral epicondylitis: comparative effectiveness of a home exercise program including stretching alone versus stretching supplemented with eccentric or concentric strengthening. *Journal of Hand Therapy*. 2005;18(4):411-20.
108. Marzetti E, Rabini A, Piccinini G, Piazzini DB, Vulpiani MC, Vetrano M, et al. Neurocognitive therapeutic exercise improves pain and function in patients with shoulder impingement syndrome: a single-blind randomized controlled clinical trial. *European journal of physical and rehabilitation medicine*. 2014;50(3):255-64.
109. Mayer F, Hirschmüller A, Müller S, Schuberth M, Baur H. Effects of short-term treatment strategies over 4 weeks in Achilles tendinopathy. *British journal of sports medicine*. 2007;41(7).
110. McCormack JR, Underwood FB, Slaven EJ, Cappaert TA. Eccentric Exercise Versus Eccentric Exercise and Soft Tissue Treatment (Astym) in the Management of Insertional Achilles Tendinopathy. *Sports health*. 2016(1941-0921 (Electronic)).
111. McGee C, Kersting E, Palmer-McLean K, Davies GJ. Standard rehabilitation vs standard plus closed kinetic chain rehabilitation for patients with shoulder impingement: A rehabilitation outcomes study. *Physical Therapy*. 1999;79.
112. McQueen KS, Powell RK, Keener T, Whalley R, Calfee RP. Role of strengthening during nonoperative treatment of lateral epicondyle tendinopathy. *Journal of hand therapy*. 2021;34(4):619-26.
113. Melegati G, Tornese D, Bandi M. Effectiveness of extracorporeal shock wave therapy associated with kinesitherapy in the treatment of subacromial impingement: A randomised, controlled study [Efficacia della terapia con onde d'urto extracorporee associata a chinesiaterapia nel trattamento della sindrome da conflitto subacromiale: Studio randomizzato controllato]. *Journal of Sports Traumatology and Related Research*. 2000;22(2):58-64.
114. Menek B, Algun C, Tarakci D, editors. Effectiveness of mulligan mobilization on range of motion and function in individuals with subacromial impingement syndrome. *Osteoporosis International Conference 18th World Conference on Osteoporosis, Degenerative Disease and Musculoskeletal Disorders, WCO-IOF-ESCEO; 2018 2018; Poland: Springer London*.
115. Mulligan EP, Huang M, Dickson T, Khazzam M. The Effect of Axioscapular and Rotator Cuff Exercise Training Sequence in Patients with Subacromial Impingement Syndrome: a Randomized Crossover Trial. *International Journal of Sports Physical Therapy*. 2016;11(1):94-107.
116. Nazligul T, Akpinar P, Aktas I, Unlu Ozkan F, Cagliyan Hartevioglu H. The effect of interferential current therapy on patients with subacromial impingement syndrome: a randomized, double-blind, sham-controlled study. *European journal of physical and rehabilitation medicine*. 2018;54(3):351-7.
117. Nishizuka T, Iwatsuki K, Kurimoto S, Yamamoto M, Hirata H. Efficacy of a forearm band in addition to exercises compared with exercises alone for lateral epicondylitis: A multicenter, randomized, controlled trial. *Journal of Orthopaedic Science*. 2017;22(2):289-94.
118. Nørregaard J, Larsen CC, Bieler T, Langberg H. Eccentric exercise in treatment of Achilles tendinopathy. *Scandinavian Journal of Medicine & Science in Sports*. 2007;17(2):133-8.
119. Notarnicola A, Maccagnano G, Tafuri S, Forcignanò MI, Panella A, Moretti B. CHELT therapy in the treatment of chronic insertional Achilles tendinopathy. 2014;29(3).
120. Nowotny J, El-Zayat B, Goronzy J, Biewener A, Bausenhardt F, Greiner S, et al. Prospective randomized controlled trial in the treatment of lateral epicondylitis with a new dynamic wrist orthosis. *European journal of medical research*. 2018;23(1):43.

121. Olausson M, Holmedal Ø, Mdala I, Brage S, Lindbæk M. Corticosteroid or placebo injection combined with deep transverse friction massage, Mills manipulation, stretching and eccentric exercise for acute lateral epicondylitis: a randomised, controlled trial. *BMC musculoskeletal disorders*. 2015;16(1):122.
122. Østerås H, Torstensen TA, Haugerud L, Østerås BS. Dose-response effects of graded therapeutic exercises in patients with long-standing subacromial pain. *Advances in Physiotherapy*. 2009;11(4):199-209.
123. Østerås H, Torstensen TA, Østerås B. High-dosage medical exercise therapy in patients with long-term subacromial shoulder pain: a randomized controlled trial. *Physiotherapy Research International : The Journal for Researchers and Clinicians in Physical Therapy*. 2010;15(4):232-42.
124. Paavola M, Malmivaara A, Taimela S, Kanto K, Inkinen J, Kalske J, et al. Subacromial decompression versus diagnostic arthroscopy for shoulder impingement: randomised, placebo surgery controlled clinical trial. *BMJ (Clinical research ed)*. 2018;362.
125. Park JY, Park HK, Choi JH, Moon ES, Kim BS, Kim WS, et al. Prospective evaluation of the effectiveness of a home-based program of isometric strengthening exercises: 12-month follow-up. *Clinics in orthopedic surgery*. 2010;2(3):173-8.
126. Pearson J, Rowlands D, Highet R. Autologous blood injection to treat achilles tendinopathy? A randomized controlled trial. *Journal of Sport Rehabilitation*. 2012;21(3).
127. Pearson SJ, Stadler S, Menz H, Morrissey D, Scott I, Munteanu S, et al. Immediate and Short-Term Effects of Short-and Long-Duration Isometric Contractions in Patellar Tendinopathy. *Clinical journal of sport medicine: official journal of the Canadian Academy of Sport Medicine*. 2018.
128. Pekgoz F, TaAkiran H, Mutlu EK, Atalay A, Celiker R. Comparison of mobilization with supervised exercise for patients with subacromial impingement syndrome. *Turkish Journal of Physical Medicine and Rehabilitation*. 2020;66(2):184-92.
129. Pekiavas NO, Baltaci G. Short-term effects of high-intensity laser therapy, manual therapy, and Kinesio taping in patients with subacromial impingement syndrome. *Lasers in medical science*. 2016;31(6):1133-41.
130. Pérez-Merino L, Casajuana MC, Bernal G, Faba J, Astilleros AE, González R, et al. Evaluation of the effectiveness of three physiotherapeutic treatments for subacromial impingement syndrome: a randomised clinical trial. *Physiotherapy*. 2016;102(1):57-63.
131. Petersen W, Welp R, Rosenbaum D. Chronic Achilles tendinopathy: a prospective randomized study comparing the therapeutic effect of eccentric training, the AirHeel brace, and a combination of both. *The American Journal of Sports Medicine*. 2007;35(10):1659-67.
132. Peterson M, Butler S, Eriksson M, Svärdsudd K. A randomized controlled trial of exercise versus wait-list in chronic tennis elbow (lateral epicondylitis). *Uppsala journal of medical sciences*. 2011;116(4):269-79.
133. Peterson M, Butler S, Eriksson M, Svärdsudd K. A randomized controlled trial of eccentric vs. concentric graded exercise in chronic tennis elbow (lateral elbow tendinopathy). *Clinical rehabilitation*. 2014;28(9):862-72.
134. Polimeni V, Panuccio A, Furfari P, Crupi D, Barreca G, Forgione C, et al. Preliminary study on the efficacy of various rehabilitation therapies for shoulder pain. *Europa Medicophysica*. 2003;39(1):59-63.
135. Praet SF, Purdam C, Welvaert M, Vlahovich N, Lovell G, Burke LM, et al. Oral supplementation of specific collagen peptides combined with calf-strengthening exercises enhances function and reduces pain in achilles tendinopathy patients. 2019;11(1).
136. Rabusin CL, Menz HB, McClelland JA, Evans AM, Malliaras P, Docking SI, et al. Efficacy of heel lifts versus calf muscle eccentric exercise for mid-portion Achilles tendinopathy (HEALTHY): a randomised trial. *British journal of sports medicine*. 2020.
137. Ramon S, Russo S, Santoboni F, Lucenteforte G, Di Luise C, de Unzurrunzaga R, et al. Focused Shockwave Treatment for Greater Trochanteric Pain Syndrome: A Multicenter, Randomized, Controlled Clinical Trial. *The Journal of bone and joint surgery American volume*. 2020;102(15):1305-11.
138. Reyhan AC, Sindel D, Dereli EE. The effects of Mulligan's mobilization with movement technique in patients with lateral epicondylitis. *Journal of Back and Musculoskeletal Rehabilitation*. 2020;33(1):99-107.

139. Rhon DI, Boyles RB, Cleland JA. One-year outcome of subacromial corticosteroid injection compared with manual physical therapy for the management of the unilateral shoulder impingement syndrome: A pragmatic randomized trial. *Annals of Internal Medicine*. 2014;161(3):161-9.
140. Rio E, Purdam C, Girdwood M, Cook J. Isometric Exercise to Reduce Pain in Patellar Tendinopathy In-Season; Is It Effective "on the Road?". *Clinical journal of sport medicine : official journal of the Canadian Academy of Sport Medicine*. 2017.
141. Rodriguez-Huguet M, Gongora-Rodriguez J, Lomas-Vega R, Martin-Valero R, Diaz-Fernandez A, Obrero-Gaitan E, et al. Percutaneous electrolysis in the treatment of lateral epicondylalgia: A single-blind randomized controlled trial. *Journal of Clinical Medicine*. 2020;9(7):1-11.
142. Rodriguez-Huguet M, Gongora-Rodriguez J, Rodriguez-Huguet P, Ibanez-Vera AJ, Rodriguez-Almagro D, Martin-Valero R, et al. Effectiveness of percutaneous electrolysis in supraspinatus tendinopathy: a single-blinded randomized controlled trial. *Journal of clinical medicine*. 2020;9(6):1-13.
143. Røe C, Ødegaard TT, Hilde F, Maehlum S, Halvorsen T. [No effect of supplement of essential fatty acids on lateral epicondylitis]. *Tidsskrift for den Norske lægeforening : tidsskrift for praktisk medicin, ny række*. 2005;125(19):2615-8.
144. Romero-Morales C, MartinLlantino PJ, CalvoLobo C, San AntolinGil M, LopezLopez D, Pedro MBD, et al. Vibration increases multifidus cross-sectional area versus cryotherapy added to chronic non-insertional Achilles tendinopathy eccentric exercise. *Physical Therapy in Sport*. 2020;42:61-7.
145. Rompe JD, Nafe B, Furia JP, Maffulli N. Eccentric loading, shock-wave treatment, or a wait-and-see policy for tendinopathy of the main body of tendo Achillis: a randomized controlled trial. *The American Journal of Sports Medicine*. 2007;35(3):374-83.
146. Rompe JD, Furia J, Maffulli N. Eccentric loading compared with shock wave treatment for chronic insertional achilles tendinopathy: A randomized, controlled trial. *Journal of Bone and Joint Surgery - Series A*. 2008;90(1):52-61.
147. Rompe JD, Segal NA, Cacchio A, Furia JP, Morral A, Maffulli N. Home training, local corticosteroid injection, or radial shock wave therapy for greater trochanter pain syndrome. *American Journal of Sports Medicine*. 2009;37(10):1981-90.
148. Rompe JD, Furia J, Maffulli N. Eccentric loading versus eccentric loading plus shock-wave treatment for midportion achilles tendinopathy: a randomized controlled trial. *The American Journal of Sports Medicine*. 2009;37(3):463-70.
149. Roos EM, Engstrom M, Lagerquist A, Soderberg B. Clinical improvement after 6 weeks of eccentric exercise in patients with mid-portion Achilles tendinopathy -- a randomized trial with 1-year follow-up [with consumer summary]. *Scandinavian Journal of Medicine & Science in Sports* 2004 Oct;14(5):286-295. 2004.
150. Rosety-Rodriguez M, Ordonez-Munoz F, Huesa-Jimenez F, Rosety Rodriguez J, Gomez-Rodriguez F, Rosety-Plaza M. Eccentric training programs for infrapatellar tendinopathy: new strategies for an old problem. 2006;4(2).
151. Santamato A, Panza F, Notarnicola A, Cassatella G, Fortunato F, De SJL, et al. Is Extracorporeal Shockwave Therapy Combined With Isokinetic Exercise More Effective Than Extracorporeal Shockwave Therapy Alone for Subacromial Impingement Syndrome? A Randomized Clinical Trial. *Journal of Orthopaedic & Sports Physical Therapy*. 2016;46(9):714-25.
152. Sawada T, Okawara H, Nakashima D, Iwabuchi S, Matsumoto M, Nakamura M, et al. Effects of alternating heat and cold stimulation using a wearable thermo-device on subjective and objective shoulder stiffness. *Journal of physiological anthropology*. 2022;41(1):1.
153. Şenbursa G, Baltacı G, Atay ÖA. The effectiveness of manual therapy in supraspinatus tendinopathy. *Acta orthopaedica et traumatologica turcica*. 2011;45(3):162-7.
154. Seven MM, Ersen O, Akpınar S, Özkan H, Türkkan S, Yıldız Y, et al. Effectiveness of prolotherapy in the treatment of chronic rotator cuff lesions. *Orthopaedics and Traumatology: Surgery and Research*. 2017;103(3):427-33.
155. Sevier TL, Stegink-Jansen C. Astymtreatment vs. eccentric exercise for lateral elbow tendinopathy: a randomized controlled clinical trial. 2015;2015(5).

156. Silbernagel KG, Thomeé R, Thomeé P, Karlsson J. Eccentric overload training for patients with chronic Achilles tendon pain--a randomised controlled study with reliability testing of the evaluation methods. *Scandinavian Journal of Medicine & Science in Sports*. 2001;11(4):197-206.
157. Silbernagel KG, Thomeé R, Eriksson BI, Karlsson J. Continued sports activity, using a pain-monitoring model, during rehabilitation in patients with Achilles tendinopathy: a randomized controlled study. *The American Journal of Sports Medicine*. 2007;35(6):897-906.
158. Şimşek HH, Balki S, Keklik SS, Öztürk H, Elden H. Does Kinesio taping in addition to exercise therapy improve the outcomes in subacromial impingement syndrome? A randomized, double-blind, controlled clinical trial. *Acta orthopaedica et traumatologica turcica*. 2013;47(2):104-10.
159. Silder A, Sherry MA, Sanfilippo J, Tuite MJ, Hetzel SJ, Heiderscheit BC. Clinical and morphological changes following 2 rehabilitation programs for acute hamstring strain injuries: A randomized clinical trial. *Journal of Orthopaedic and Sports Physical Therapy*. 2013;43(5):284-99.
160. Smidt N, van der Windt DAWM, Assendelft WJJ, Devillé WLJM, Korthals-de Bos IBC, Bouter LM. Corticosteroid injections, physiotherapy, or a wait-and-see policy for lateral epicondylitis: a randomised controlled trial. *The Lancet*. 2002;359(9307):657-62.
161. Söderberg J, Grooten WJ, Äng BO. Effects of eccentric training on hand strength in subjects with lateral epicondylalgia: a randomized-controlled trial. *Scandinavian Journal of Medicine & Science in Sports*. 2012;22(6):797-803.
162. Stasinopoulos D, Manias P. Comparing two exercise programmes for the management of lateral elbow tendinopathy (tennis elbow/lateral epicondylitis)-A controlled clinical trial. *The Open Access Journal of Science and Technology*. 2013;1.
163. Stasinopoulos D, Stasinopoulos I. Comparison of effects of Cyriax physiotherapy, a supervised exercise programme and polarized polychromatic non-coherent light (Biopton light) for the treatment of lateral epicondylitis. *Clinical rehabilitation*. 2006;20(1):12-23.
164. Stasinopoulos D, Stasinopoulos I, Pantelis M, Stasinopoulou K. Comparing the effects of exercise program and low-level laser therapy with exercise program and polarized polychromatic non-coherent light (bioptron light) on the treatment of lateral elbow tendinopathy. *Photomedicine and laser surgery*. 2009;27(3):513-20.
165. Stasinopoulos D, Stasinopoulos I, Pantelis M, Stasinopoulou K. Comparison of effects of a home exercise programme and a supervised exercise programme for the management of lateral elbow tendinopathy. *British journal of sports medicine*. 2010;44(8):579-83.
166. Stasinopoulos D, Stasinopoulos I. Comparison of effects of eccentric training, eccentric-concentric training, and eccentric-concentric training combined with isometric contraction in the treatment of lateral elbow tendinopathy. *Journal of Hand Therapy*. 2017;30(1):13-9.
167. Stefansson SH, Brandsson S, Langberg H, Arnason A. Using Pressure Massage for Achilles Tendinopathy: A Single-Blind, Randomized Controlled Trial Comparing a Novel Treatment Versus an Eccentric Exercise Protocol. *Orthopaedic journal of sports medicine*. 2019;7(3):2325967119834284.
168. Stergioulas A. Effects of low-level laser and plyometric exercises in the treatment of lateral epicondylitis. 2007;25(3).
169. Stergioulas A, Stergioula M, Aarskog R, Lopes-Martins R, Bjordal JM. Effects of low-level laser therapy and eccentric exercises in the treatment of recreational athletes with chronic achilles tendinopathy. *The American Journal of Sports Medicine*. 2008;36(5):881-7.
170. Steunebrink M, Zwerver J, Brandsema R, Groenenboom P, van den Akker-Scheek I, Weir A. Topical glyceryl trinitrate treatment of chronic patellar tendinopathy: a randomised, double-blind, placebo-controlled clinical trial. *British journal of sports medicine*. 2013;47(1):34-9.
171. Stevens M, Tan C-W. Effectiveness of the Alfredson protocol compared with a lower repetition-volume protocol for midportion Achilles tendinopathy: a randomized controlled trial. *journal of orthopaedic & sports physical therapy*. 2014;44(2):59-67.
172. Struijs PAA, Kerkhoffs GMMJ, Assendelft WJJ, van Dijk CN. Conservative Treatment of Lateral Epicondylitis: Brace Versus Physical Therapy or a Combination of Both—A Randomized Clinical Trial. *The American Journal of Sports Medicine*. 2004;32(2):462-9.

173. Svernlöv B, Adolfsson L. Non-operative treatment regime including eccentric training for lateral humeral epicondylalgia. *Scandinavian Journal of Medicine & Science in Sports*. 2001;11(6):328-34.
174. Tahrán Ö, Yeşilyaprak SS. Effects of Modified Posterior Shoulder Stretching Exercises on Shoulder Mobility, Pain, and Dysfunction in Patients With Subacromial Impingement Syndrome. *Sports health*. 2020;12(2):139-48.
175. Thijs KM, Zwerver J, Backx FJG, Steeneken V, Rayer S, Groenenboom P, et al. Effectiveness of shockwave treatment combined with eccentric training for patellar tendinopathy: a double-blinded randomized study. *Clinical journal of sport medicine*. 2017;27(2):89-96.
176. Tonks JH. Evaluation of short-term conservative treatment in patients with tennis elbow (lateral epicondylitis): A prospective randomised, assessor-blinded trial. University of Central Lancashire; 2012.
177. Tonks JH, Pai SK, Murali SR. Steroid injection therapy is the best conservative treatment for lateral epicondylitis: A prospective randomised controlled trial. *International journal of clinical practice*. 2007;61(2):240-6.
178. Tumilty S, McDonough S, Hurley DA, Baxter GD. Clinical effectiveness of low-level laser therapy as an adjunct to eccentric exercise for the treatment of Achilles' tendinopathy: a randomized controlled trial. *Archives of Physical Medicine and Rehabilitation*. 2012;93(5):733-9.
179. Tumilty S, Mani R, Baxter GD. Photobiomodulation and eccentric exercise for Achilles tendinopathy: a randomized controlled trial. *Lasers in Medical Science*. 2016;31(1):127-35.
180. Turgut E, Duzgun I, Baltaci G. Effects of Scapular Stabilization Exercise Training on Scapular Kinematics, Disability, and Pain in Subacromial Impingement: A Randomized Controlled Trial. *Archives of Physical Medicine and Rehabilitation*. 2017;98(10):1915.
181. Tyler TF, Thomas GC, Nicholas SJ, McHugh MP. Addition of isolated wrist extensor eccentric exercise to standard treatment for chronic lateral epicondylitis: a prospective randomized trial. *Journal of Shoulder and Elbow surgery*. 2010;19(6):917-22.
182. Valles-Carrascosa E, Gallego-Izquierdo T, Jimenez-Rejano J, Plaza-Manzano G, Pecos-Martin D, Hita-Contreras F, et al. Pain, motion and function comparison of two exercise protocols for the rotator cuff and scapular stabilizers in patients with subacromial syndrome [with consumer summary]. *Journal of Hand Therapy* 2018 Apr-Jun;31(2):227-237. 2018.
183. Van Der Vlist AC, Van Oosterom RF, Van Veldhoven PLJ, BiermaZeinstra SMA, Waarsing JH, Verhaar JAN, et al. Effectiveness of a high volume injection as treatment for chronic Achilles tendinopathy: Randomised controlled trial. *The BMJ*. 2020;370.
184. Van Ark M, Cook JL, Docking SI, Zwerver J, Gaida JE, Van Den Akker-Scheek I, et al. Do isometric and isotonic exercise programs reduce pain in athletes with patellar tendinopathy in-season? A randomised clinical trial. *Journal of science and medicine in sport*. 2016;19(9):702-6.
185. Vinuesa-Montoya S, Aguilar-Ferrández ME, Matarán-Peñarocha GA, Fernández-Sánchez M, Fernández-Espinar EM, Castro-Sánchez AM. A Preliminary Randomized Clinical Trial on the Effect of Cervicothoracic Manipulation Plus Supervised Exercises vs a Home Exercise Program for the Treatment of Shoulder Impingement. *J Chiropr Med*. 2017;16(2):85-93.
186. Visnes H, Hoksrud A, Cook J, Bahr R. No effect of eccentric training on jumper's knee in volleyball players during the competitive season: a randomized clinical trial. 2005;15(4).
187. Vuvan V, Vicenzino B, Mellor R, Heales LJ, Coombes BK. Unsupervised Isometric Exercise versus Wait-and-See for Lateral Elbow Tendinopathy. *Medicine and science in sports and exercise*. 2020;52(2):287-95.
188. Walther M, Werner A, Stahlschmidt T, Woelfel R, Gohlke F. The subacromial impingement syndrome of the shoulder treated by conventional physiotherapy, self-training, and a shoulder brace: Results of a prospective, randomized study. *Journal of Shoulder and Elbow Surgery*. 2004;13(4):417-23.
189. Warden SJ, Metcalf BR, Kiss ZS, Cook JL, Purdam CR, Bennell KL, et al. Low-intensity pulsed ultrasound for chronic patellar tendinopathy: a randomized, double-blind, placebo-controlled trial. *Rheumatology (Oxford, England)*. 2008;47(4).
190. Wegener RL, Brown T, O'Brien L. A randomized controlled trial of comparative effectiveness of elastic therapeutic tape, sham tape or eccentric exercises alone for lateral elbow tendinosis. 2016;21(4).

191. Wen DY, Schultz BJ, Schaal B, Graham ST, Kim BS. Eccentric strengthening for chronic lateral epicondylitis: a prospective randomized study. *Sports health*. 2011;3(6):500-3.
192. Werner A, Walther M, Ilg A, Stahlschmidt T, Gohlke F. Self-training versus conventional physiotherapy in subacromial impingement syndrome. *Zeitschrift für Orthopädie und ihre Grenzgebiete*. 2002;140(4):375-80.
193. Wiedmann M, Mauch F, Huth J, Burkhardt P, Drews BH. Treatment of mid-portion Achilles tendinopathy with eccentric training and its effect on neovascularization. *Sports Orthopaedics and Traumatology*. 2017;33(3):278-85.
194. Yazmalar L, Sarıyıldız MA, Batmaz İ, Alpaycı M, Burkan YK, Özkan Y, et al. Efficiency of therapeutic ultrasound on pain, disability, anxiety, depression, sleep and quality of life in patients with subacromial impingement syndrome: A randomized controlled study. *Journal of back and musculoskeletal rehabilitation*. 2016;29(4):801-7.
195. Yeldan I, Cetin E, Ozdinciler AR. The effectiveness of low-level laser therapy on shoulder function in subacromial impingement syndrome. *Disability and rehabilitation*. 2009;31(11):935-40.
196. Yelland MJ, Sweeting KR, Lyftogt JA, Ng SK, Scuffham PA, Evans KA. Prolotherapy injections and eccentric loading exercises for painful Achilles tendinosis: a randomised trial. *Br J Sports Med*. 2011;45(5):421-8.
197. Yerlikaya M, Talay Çalış H, Tomruk Sütbeyaz S, Sayan H, Ibiş N, Koç A, et al. Comparison of Effects of Leukocyte-Rich and Leukocyte-Poor Platelet-Rich Plasma on Pain and Functionality in Patients With Lateral Epicondylitis. *Archives of rheumatology*. 2017;33(1):73-9.
198. Young MA, Cook JL, Purdam CR, Kiss ZS, Alfredson H. Eccentric decline squat protocol offers superior results at 12 months compared with traditional eccentric protocol for patellar tendinopathy in volleyball players [with consumer summary]. *British Journal of Sports Medicine* 2005 Feb;39(2):102-105. 2005.
199. Yu J, Park D, Lee G. Effect of eccentric strengthening on pain, muscle strength, endurance, and functional fitness factors in male patients with achilles tendinopathy. *American journal of physical medicine & rehabilitation*. 2013;92(1):68-76.
200. Yuruk ZO, Kirdi N, editors. The effects of radial extracorporeal shock wave therapy on subjective pain and functionality in patients with lateral epicondylitis. *Fizyoterapi Rehabilitasyon Conference: 15th Congress of Advances in Physiotherapy*; 2014; Ankara Turkey: Turkish Physical Therapy Association.

Supplementary file 7: Summary risk of bias of included studies

Risk of bias assessments were made for each outcome and time point in a study. The results presented here represent a summary, with the mode value selected.

Author, Year	Random sequence generation	Allocation concealment	Blinding of participants /personnel	Blinding of outcome assessment	Incomplete outcome bias	Selective reporting	Other bias
Abat et al 2016 ¹	Not applicable (quasi)	Not applicable (quasi)	Unclear	Unclear	Low risk	Unclear	High risk
Aceituno-Gómez et al 2019 ²	Low risk	Unclear	Low risk	Low risk	Low risk	Low risk	High risk
Akkaya et al 2016 ³	Low risk	Unclear	High risk	High risk	Low risk	Unclear	Low risk
Aktas et al 2007 ⁴	Low risk	Unclear	Low risk	Low risk	Low risk	Unclear	Low risk
Akyol et al 2012 ⁵	Low risk	Low risk	Low risk	Low risk	Low risk	Unclear	High risk
Alfredson et al 1998 ⁶	High risk	Unclear	High risk	Unclear	Low risk	Unclear	High risk
Alfredson et al 1999 ⁷	Not applicable (quasi)	Not applicable (quasi)	High risk	High risk	Low risk	Unclear	High risk
Arias-Buría et al 2015 ⁸	Low risk	Low risk	High risk	Low risk	Low risk	Unclear	Low risk
Arias--Buría et al 2017 ⁹	Low risk	Low risk	Unclear	Low risk	Unclear	Low risk	High risk
Bae et al 2011 ¹⁰	Not applicable (quasi)	Not applicable (quasi)	Unclear	Unclear	Unclear	Unclear	High risk
Bagcier et al 2019 ¹¹	Not applicable (quasi)	Not applicable (quasi)	Unclear	Low risk	Unclear	Unclear	High risk
Bahr et al 2006 ¹²	Low risk	Low risk	High risk	Low risk	Low risk	Unclear	Low risk
Balius et al 2016 ¹³	Low risk	Low risk	Unclear	Low risk	Low risk	Unclear	Low risk
Bang et al 2000 ¹⁴	Low risk	Unclear	Unclear	Low risk	Low risk	Unclear	Low risk
Başkurt et al 2011 ¹⁵	Low risk	Unclear	High risk	Unclear	Low risk	Unclear	Low risk
Beaudreuil et al 2012 ¹⁶	Unclear	Low risk	Low risk	Low risk	Low risk	Unclear	Low risk

Swinton, Shim, Pavlova, Moss, Maclean, Greig, Parkinson, Morrissey, Alexander, Cooper (2021)

Author, Year	Random sequence generation	Allocation concealment	Blinding of participants /personnel	Blinding of outcome assessment	Incomplete outcome bias	Selective reporting	Other bias
Bek et al 2012 ¹⁷	Low risk	Unclear	High risk	High risk	Low risk	Low risk	Low risk
Bell et al 2013 ¹⁸	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk
Bennell et al 2010 ¹⁹	Bennell <i>et al.</i>	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk
Beyer et al 2015 ²⁰	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	High risk
Blume et al 2015 ²¹	Unclear	Low risk	Low risk	Low risk	Low risk	Unclear	Low risk
Boesen et al 2017 ²²	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk
Boesen et al 2019 ²³	Low risk	Low risk	Low risk	Low risk	Low risk	Unclear	Low risk
Boudreau et al 2019 ²⁴	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk
Breda et al 2020 ²⁵	Low risk	Low risk	High risk	Low risk	Low risk	High risk	High risk
Brox et al 1999 ²⁶	High risk	High risk	High risk	High risk	No Data	No Data	No Data
Buyuksireci et al 2020 ²⁷	Low risk	Unclear	High risk	Unclear	Low risk	Unclear	High risk
Cacchio et al 2011 ²⁸	Low risk	Low risk	High risk	Low risk	Unclear	Unclear	Low risk
Calis et al 2011 ²⁹	Low risk	Low risk	Unclear	Unclear	Low risk	Unclear	Unclear
Celik et al 2009 ³⁰	Unclear	Unclear	Low risk	Low risk	Low risk	Low risk	Low risk
Celik et al 2009 ³¹	Unclear	Unclear	Unclear	Unclear	Low risk	Low risk	High risk
Celik et al 2019 ³²	Low risk	Low risk	Unclear	Unclear	Low risk	Unclear	Unclear
Cha et al 2014 ³³	Unclear	Unclear	Unclear	Unclear	Low risk	Unclear	Unclear
Chaconas et al 2017 ³⁴	Low risk	Unclear	Unclear	Low risk	High risk	Unclear	High risk
Chary-Valckenaere et al 2018 ³⁵	Low risk	Low risk	Unclear	Low risk	High risk	Low risk	Low risk
Chen et al 2017 ³⁶	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	High risk
Cheng et al 2007 ³⁷	High risk	High risk	Unclear	Unclear	Unclear	Unclear	High risk
Cho et al 2017 ³⁸	High risk	High risk	Unclear	Unclear	Low risk	Low risk	Unclear
Chung et al 2004 ³⁹	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk

Swinton, Shim, Pavlova, Moss, Maclean, Greig, Parkinson, Morrissey, Alexander, Cooper (2021)

Author, Year	Random sequence generation	Allocation concealment	Blinding of participants /personnel	Blinding of outcome assessment	Incomplete outcome bias	Selective reporting	Other bias
Clarke et al 2010 ⁴⁰	Low risk	Low risk	Low risk	Low risk	Low risk	Unclear	Low risk
Conroy et al 1998 ⁴¹	Unclear	Unclear	Unclear	Low risk	Low risk	Unclear	Low risk
Coombes et al 2013 ⁴²	Low risk	Low risk	High risk	Low risk	Low risk	Low risk	Low risk
Croisier et al 2007 ⁴³	Not applicable (quasi)	Not applicable (quasi)	Unclear	Low risk	Unclear	Unclear	High risk
de Jonge et al 2008 ⁴⁴	Unclear	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk
de Miguel Valtierra et al 2018 ⁴⁵	Low risk	Low risk	High risk	Low risk	Low risk	Low risk	High risk
De Oliveira et al 2020 ⁴⁶	Low risk	Low risk	Low risk	Low risk	Low risk	High risk	High risk
de Vos et al 2007 ⁴⁷	Low risk	Unclear	Low risk	Low risk	Unclear	Low risk	High risk
de Vos et al 2010 ⁴⁸	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk
Dejaco et al 2017 ⁴⁹	Low risk	Low risk	Low risk	Low risk	Low risk	High risk	Low risk
Devereaux et al 2016 ⁵⁰	Low risk	High risk	High risk	High risk	High risk	Unclear	Low risk
Dimitrios et al 2012 ⁵¹	Not applicable (quasi)	Not applicable (quasi)	Low risk	Low risk	Low risk	Unclear	High risk
Dimitrios et al 2013 ⁵²	Not applicable (quasi)	Not applicable (quasi)	Low risk	Low risk	Low risk	Unclear	High risk
Dogan et al 2010 ⁵³	Low risk	Low risk	Low risk	Low risk	Low risk	Unclear	Low risk
Dupuis et al ⁵⁴	Low risk	Low risk	High risk	Low risk	Low risk	Low risk	High risk
EkenGedik et al 2016 ⁵⁵	Low risk	Low risk	Unclear	Low risk	Unclear	Unclear	High risk
Ellegaard et al 2016 ⁵⁶	Low risk	Low risk	High risk	Low risk	Low risk	Unclear	Low risk
Engebretsen et al 2009 ⁵⁷	Low risk	Low risk	High risk	Low risk	Low risk	Unclear	Low risk

Author, Year	Random sequence generation	Allocation concealment	Blinding of participants /personnel	Blinding of outcome assessment	Incomplete outcome bias	Selective reporting	Other bias
Engebretsen et al 2011 ⁵⁸	Low risk	Low risk	High risk	Low risk	Low risk	Unclear	Low risk
Eraslan et al 2018 ⁵⁶	Eraslan <i>et al.</i>	Low risk	Unclear	Unclear	Unclear	Low risk	Unclear
Faria et al 2006 ⁵⁹	High risk	High risk	Unclear	Unclear	Unclear	Unclear	Unclear
Fournier Belley et al 2018 ⁶⁰	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk
Ganderton et al 2018 ⁶¹	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	High risk
García et al 2016 ⁶²	Low risk	Low risk	Low risk	Low risk	Low risk	Unclear	High risk
Gatz et al 2020 ⁶³	Low risk	Low risk	Low risk	Low risk	Unclear	Unclear	High risk
Genc et al 2020 ⁶⁴	Low risk	Low risk	High risk	High risk	Unclear	Unclear	High risk
Giray et al 2019 ⁶⁵	Low risk	Low risk	High risk	Low risk	Low risk	Low risk	High risk
Granviken et al 2015 ⁶⁶	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk
Gürsel et al 2004 ⁶⁷	Low risk	Low risk	Low risk	Unclear	Unclear	Unclear	High risk
Hallgren 2017 ⁶⁸	Unclear	Unclear	Low risk	Low risk	Low risk	Low risk	Low risk
Hallgren et al 2014 ⁶⁹	High risk	Low risk	High risk	Low risk	Unclear	Low risk	Low risk
Heron et al 2017 ⁷⁰	Low risk	Low risk	Low risk	Low risk	High risk	High risk	Low risk
Holmgren et al 2012 ⁷¹	Low risk	Low risk	Low risk	Low risk	Unclear	Low risk	High risk
Hotta et al 2020 ⁷²	Low risk	Low risk	High risk	Low risk	Low risk	Low risk	High risk
Houck et al 2015 ⁷³	Low risk	Unclear	Unclear	High risk	Low risk	Low risk	Low risk
Ilhanli et al. 2015 ⁷⁴	Low risk	Low risk	Unclear	Low risk	Unclear	Unclear	Low risk
Ingwersen et al 2017 ⁷⁵	Low risk	Low risk	Low risk	Low risk	Low risk	High risk	Low risk

Author, Year	Random sequence generation	Allocation concealment	Blinding of participants /personnel	Blinding of outcome assessment	Incomplete outcome bias	Selective reporting	Other bias
Jasnauskaitė-Gedrimė et al 2018 ⁷⁶	Unclear	Unclear	Unclear	Unclear	Low risk	Unclear	High risk
Jeong et al 2008 ⁷⁷	Unclear	Unclear	Unclear	Unclear	Low risk	Unclear	Unclear
Johansson et al 2005 ⁷⁸	Low risk	Unclear	Unclear	Low risk	Low risk	Unclear	High risk
Jonsson et al 2005 ⁷⁹	Unclear	Unclear	Low risk	Unclear	High risk	Unclear	High risk
Jonsson 2009 ⁸⁰	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear
Juul-Kristensen et al 2019 ⁸¹	Low risk	Low risk	Low risk	Low risk	Low risk	Unclear	Low risk
Kachanathu et al 2019 ⁸²	Low risk	Low risk	Unclear	Low risk	Low risk	Unclear	High risk
Kang et al 2019 ⁸³	Low risk	Low risk	High risk	Low risk	High risk	High risk	High risk
Kedia et al 2014 ⁸⁴	Low risk	Low risk	Low risk	Low risk	Low risk	Unclear	Low risk
Ketola et al 2009 ⁸⁵	Low risk	Low risk	Unclear	Low risk	Low risk	Unclear	High risk
Ketola et al 2013 ⁸⁶	Low risk	Low risk	High risk	Low risk	Low risk	Unclear	Low risk
Kim et al 2017 ⁸⁷	Low risk	Unclear	Unclear	Low risk	Unclear	Low risk	Low risk
Kim et al 2020 ⁸⁸	Low risk	Low risk	High risk	High risk	Low risk	Unclear	High risk
Knobloch et al 2007 ⁸⁹	Low risk	Low risk	Unclear	Unclear	High risk	Unclear	High risk
Knobloch et al 2007 ⁹⁰	Unclear	Unclear	High risk	Low risk	Unclear	Unclear	High risk
Knobloch et al 2008 ⁹¹	Unclear	Low risk	Unclear	Unclear	Unclear	Unclear	High risk
Koç et al 2020 ⁹²	Low risk	Unclear	High risk	Low risk	Low risk	Unclear	Low risk
Kongsgaard et al 2009 ⁹³	Low risk	Low risk	High risk	Low risk	Low risk	Unclear	Low risk

Author, Year	Random sequence generation	Allocation concealment	Blinding of participants /personnel	Blinding of outcome assessment	Incomplete outcome bias	Selective reporting	Other bias
Kromer et al 2014 ⁹⁴	Low risk	Low risk	High risk	High risk	Low risk	Low risk	Low risk
Kromer et al 2013 ⁹⁵	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk
Kulig et al 2009 ⁹⁶	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	High risk
Land et al 2019 ⁹⁷	Low risk	Low risk	Low risk	Low risk	High risk	Unclear	Low risk
Lee et al 2017 ⁹⁸	Low risk	Unclear	Unclear	Unclear	Low risk	Unclear	High risk
Lee et al 2014 ⁹⁹	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	High risk
Lee et al 2017 ¹⁰⁰	No Data	No Data	No Data	No Data	No Data	No Data	No Data
Littlewood et al 2016 ¹⁰¹	Low risk	Low risk	High risk	High risk	Unclear	Unclear	High risk
Ludewig et al 2003 ¹⁰²	Low risk	Unclear	High risk	Unclear	Low risk	Unclear	Low risk
Luginbuhl et al 2008 ¹⁰³	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	High risk
Maenhout et al 2013 ¹⁰⁴	Unclear	High risk	High risk	High risk	Low risk	Unclear	Low risk
Mafi et al 2001 ¹⁰⁵	Low risk	Unclear	Unclear	Unclear	Unclear	Unclear	High risk
Manias et al 2006 ¹⁰⁶	High risk	High risk	High risk	High risk	Low risk	Unclear	Unclear
Martinez-Silvestrini et al 2005 ¹⁰⁷	Unclear	Unclear	Unclear	Unclear	Low risk	Unclear	High risk
Marzetti et al 2014 ¹⁰⁸	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk
Mayer et al 2007 ¹⁰⁹	Unclear	Unclear	High risk	High risk	Unclear	Unclear	High risk
McCormack et al 2016 ¹¹⁰	Low risk	Low risk	Unclear	Unclear	Low risk	Low risk	Low risk
McGee et al 1999 ¹¹¹	Unclear	Unclear	Unclear	Unclear	High risk	Unclear	High risk

Author, Year	Random sequence generation	Allocation concealment	Blinding of participants /personnel	Blinding of outcome assessment	Incomplete outcome bias	Selective reporting	Other bias
McQueen et al 2020 ¹¹²	High risk	Unclear	Unclear	Unclear	High risk	Unclear	High risk
Melegati et al 2000 ¹¹³	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	High risk
Menek et al 2019 ¹¹⁴	Low risk	Low risk	Unclear	Unclear	Low risk	Unclear	Unclear
Mulligan et al 2016 ¹¹⁵	Low risk	Low risk	Low risk	Low risk	Low risk	Unclear	High risk
Nazligul et al 2018 ¹¹⁶	Low risk	Low risk	Low risk	Unclear	Low risk	Unclear	Low risk
Nishizuka et al 2017 ¹¹⁷	Low risk	Low risk	High risk	Unclear	Low risk	Unclear	High risk
Nørregaard et al 2007 ¹¹⁸	Low risk	Low risk	Unclear	Unclear	Unclear	Unclear	High risk
Notarnicola et al 2014 ¹¹⁹	Not applicable (quasi)	Not applicable (quasi)	High risk	Unclear	Unclear	Unclear	High risk
Nowotny et al 2018 ¹²⁰	Low risk	Unclear	Low risk	Low risk	High risk	Unclear	High risk
Olaussen et al 2015 ¹²¹	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk
Østerås et al 2009 ¹²²	Low risk	Low risk	Unclear	High risk	Low risk	Unclear	Low risk
Østerås et al 2010 ¹²³	Low risk	Low risk	High risk	High risk	Low risk	Unclear	High risk
Paavola et al 2018 ¹²⁴	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk
Park et al 2010 ¹²⁵	Low risk	Unclear	Unclear	Unclear	Unclear	Unclear	High risk
Pearson et al 2012 ¹²⁶	Unclear	Unclear	High risk	Unclear	Low risk	Unclear	High risk

Author, Year	Random sequence generation	Allocation concealment	Blinding of participants /personnel	Blinding of outcome assessment	Incomplete outcome bias	Selective reporting	Other bias
Pearson et al 2018 ¹²⁷	Low risk	Low risk	High risk	Low risk	Low risk	Low risk	High risk
Pekgöz et al 2020 ¹²⁸	Low risk	Low risk	Low risk	Low risk	Low risk	Unclear	Low risk
Pekyavas et al 2016 ¹²⁹	Low risk	Low risk	High risk	Low risk	Unclear	Unclear	Low risk
Pérez-Merino et al 2016 ¹³⁰	Low risk	Unclear	Low risk	Low risk	Unclear	Low risk	Low risk
Petersen et al 2007 ¹³¹	Low risk	Unclear	Unclear	Unclear	Unclear	Unclear	High risk
Peterson et al 2011 ¹³²	Low risk	Low risk	Unclear	High risk	Low risk	Low risk	Low risk
Peterson et al 2014 ¹³³	Low risk	Unclear	Low risk	High risk	Low risk	Low risk	Low risk
Polimeni et al 2003 ¹³⁴	Unclear	Unclear	High risk	Low risk	Unclear	Unclear	Unclear
Praet et al 2019 ¹³⁵	Low risk	Low risk	Low risk	Low risk	Low risk	Unclear	Low risk
Rabusin et al 2020 ¹³⁶	Low risk	Low risk	High risk	High risk	Low risk	Low risk	High risk
Ramon et al 2020 ¹³⁷	Unclear	Unclear	Low risk	Low risk	Low risk	Unclear	High risk
Reyhan et al 2020 ¹³⁸	Low risk	Low risk	Unclear	Unclear	Low risk	Unclear	High risk
Rhon et al 2014 ¹³⁹	Low risk	Low risk	High risk	Low risk	Low risk	Low risk	Low risk
Rio et al 2017 ¹⁴⁰	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	High risk
Rodríguez-Huguet et al 2020 ¹⁴¹	Low risk	Low risk	Unclear	Low risk	Low risk	Low risk	Low risk
Rodríguez-Huguet et al 2020 ¹⁴²	Low risk	Low risk	Unclear	Low risk	Low risk	Low risk	Low risk

Author, Year	Random sequence generation	Allocation concealment	Blinding of participants /personnel	Blinding of outcome assessment	Incomplete outcome bias	Selective reporting	Other bias
Røe et al 2005 ¹⁴³	No Data	No Data	No Data	No Data	No Data	No Data	No Data
Romero-Morales et al 2020 ¹⁴⁴	Unclear	Unclear	Unclear	Unclear	Low risk	Low risk	High risk
Rompe et al 2007 ¹⁴⁵	Low risk	Low risk	Unclear	Low risk	Low risk	Unclear	Low risk
Rompe et al 2008 ¹⁴⁶	Low risk	Low risk	Unclear	Low risk	Low risk	Unclear	Unclear
Rompe et al 2009 ¹⁴⁷	Unclear	Low risk	High risk	High risk	Low risk	Unclear	Low risk
Rompe et al 2009 ¹⁴⁸	Low risk	Low risk	High risk	Low risk	Low risk	Unclear	Low risk
Roos et al 2004 ¹⁴⁹	Low risk	Unclear	Unclear	Low risk	Low risk	Unclear	Low risk
Rosety-Rodriguez et al 2006 ¹⁵⁰	Low risk	Unclear	Unclear	Unclear	Unclear	Unclear	High risk
Santamato et al 2016 ¹⁵¹	Low risk	Low risk	High risk	Low risk	Low risk	Unclear	High risk
Scott et al 2019 ¹⁵²	Low risk	Low risk	High risk	Low risk	Low risk	Low risk	Low risk
Şenbursa et al 2011 ¹⁵³	Low risk	Unclear	Unclear	Unclear	Low risk	Unclear	Low risk
Seven et al 2017 ¹⁵⁴	Low risk	Low risk	High risk	Low risk	High risk	Unclear	Low risk
Sevier et al 2015 ¹⁵⁵	Low risk	Unclear	High risk	High risk	High risk	Unclear	High risk
Silbernagel et al 2001 ¹⁵⁶	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	High risk
Silbernagel et al 2007 ¹⁵⁷	Low risk	Low risk	High risk	High risk	Low risk	Unclear	Low risk
Şimşek et al 2013 ¹⁵⁸	Unclear	Unclear	Unclear	Low risk	Unclear	Unclear	Unclear
Slider et al 2013 ¹⁵⁹	Low risk	Unclear	Low risk	Low risk	Low risk	Unclear	Unclear
Smidt et al 2002 ¹⁶⁰	Low risk	Low risk	High risk	High risk	Low risk	Unclear	Low risk

Author, Year	Random sequence generation	Allocation concealment	Blinding of participants /personnel	Blinding of outcome assessment	Incomplete outcome bias	Selective reporting	Other bias
Söderberg et al 2012 ¹⁶¹	Unclear	Low risk	Low risk	Low risk	Low risk	Unclear	Low risk
Stasinopoulos 2013 ¹⁶²	High risk	High risk	High risk	Low risk	Low risk	Unclear	High risk
Stasinopoulos et al 2006 ¹⁶³	Not applicable (quasi)	Not applicable (quasi)	Unclear	Low risk	Low risk	Unclear	High risk
Stasinopoulos et al 2009 ¹⁶⁴	Not applicable (quasi)	Low risk	High risk	Low risk	Unclear	Unclear	High risk
Stasinopoulos et al 2010 ¹⁶⁵	Not applicable (quasi)	Not applicable (quasi)	Low risk	Low risk	Low risk	Unclear	High risk
Stasinopoulos et al 2017 ¹⁶⁶	Low risk	Unclear	Low risk	Low risk	Low risk	Unclear	High risk
Stefansson et al 2019 ¹⁶⁷	Low risk	Unclear	High risk	Low risk	High risk	Unclear	Low risk
Stergioulas et al 2007 ¹⁶⁸	Low risk	Low risk	High risk	Low risk	Unclear	Unclear	Unclear
Stergioulas et al 2008 ¹⁶⁹	Low risk	Low risk	Low risk	Low risk	Low risk	Unclear	Low risk
Steunebrink et al 2013 ¹⁷⁰	Low risk	Low risk	Low risk	Low risk	Low risk	Unclear	Low risk
Stevens et al 2014 ¹⁷¹	Unclear	Unclear	High risk	High risk	Unclear	Unclear	High risk
Struijs et al 2004 ¹⁷²	Low risk	Low risk	High risk	Low risk	Low risk	Unclear	High risk
Svernlöv et al 2001 ¹⁷³	Not applicable (quasi)	Not applicable (quasi)	Unclear	Unclear	Unclear	Unclear	High risk
Tahrán et al 2020 ¹⁷⁴	Low risk	Unclear	Low risk	Low risk	Low risk	Unclear	Low risk
Thijs et al 2017 ¹⁷⁵	Low risk	Low risk	Low risk	Low risk	Low risk	Unclear	Low risk
Tonks 2012 ¹⁷⁶	No Data	No Data	No Data	No Data	No Data	No Data	No Data

Author, Year	Random sequence generation	Allocation concealment	Blinding of participants /personnel	Blinding of outcome assessment	Incomplete outcome bias	Selective reporting	Other bias
Tonks et al 2007 ¹⁷⁷	Low risk	Low risk	Low risk	High risk	High risk	Low risk	Low risk
Tumilty et al 2012 ¹⁷⁸	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk
Tumilty et al 2016 ¹⁷⁹	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk
Turgut et al 2017 ¹⁸⁰	Low risk	Unclear	Unclear	Unclear	High risk	Unclear	Low risk
Tyler et al 2010 ¹⁸¹	Unclear	Unclear	Unclear	Low risk	Unclear	Unclear	High risk
Vallés-Carrascosa et al 2018 ¹⁸²	Low risk	Low risk	Low risk	High risk	Low risk	Low risk	High risk
van Der Vlist 2020 ¹⁸³	Low risk	Low risk	High risk	Low risk	Low risk	Low risk	Low risk
vanArk et al 2016 ¹⁸⁴	Low risk	Low risk	Low risk	Unclear	Unclear	Low risk	Low risk
Vinuesa-Montoya et al 2017 ¹⁸⁵	Low risk	Low risk	High risk	Low risk	Low risk	Low risk	Low risk
Visnes et al 2005 ¹⁸⁶	Low risk	Low risk	High risk	Low risk	Unclear	Unclear	Unclear
Vuvan et al 2019 ¹⁸⁷	Low risk	Low risk	High risk	High risk	Low risk	Low risk	Low risk
Walther et al 2004 ¹⁸⁸	Unclear	Unclear	Unclear	Unclear	Low risk	Unclear	Unclear
Warden et al 2008 ¹⁸⁹	Low risk	Low risk	Low risk	Low risk	Low risk	Unclear	Low risk
Wegener et al 2016 ¹⁹⁰	Low risk	Low risk	High risk	Low risk	Low risk	Unclear	High risk
Wen et al 2011 ¹⁹¹	Unclear	Unclear	Low risk	Unclear	High risk	Unclear	High risk
Werner et al 2002 ¹⁹²	Low risk	Unclear	Unclear	Unclear	Unclear	Unclear	High risk
Wiedmann et al 2017 ¹⁹³	Low risk	Low risk	Unclear	Unclear	Unclear	Unclear	High risk

Author, Year	Random sequence generation	Allocation concealment	Blinding of participants /personnel	Blinding of outcome assessment	Incomplete outcome bias	Selective reporting	Other bias
Yazmalar et al 2016 ¹⁹⁴	Low risk	Unclear	Unclear	Unclear	Unclear	Unclear	High risk
Yeldan et al 2009 ¹⁹⁵	Low risk	Unclear	Unclear	Unclear	Unclear	Low risk	Low risk
Yelland et al 2011 ¹⁹⁶	Low risk	Low risk	High risk	Low risk	Low risk	Unclear	Low risk
Yerlikaya et al 2018 ¹⁹⁷	Low risk	Unclear	Low risk	Low risk	Unclear	Unclear	High risk
Young et al 2005 ¹⁹⁸	Unclear	Unclear	High risk	Low risk	High risk	Unclear	High risk
Yu et al 2013 ¹⁹⁹	Low risk	Low risk	Low risk	Unclear	Low risk	Unclear	Unclear
Yuruk et al 2014 ²⁰⁰	Low risk	Low risk	Unclear	Unclear	Low risk	Unclear	Unclear