



Exploring the Dose-Response Relationship Between Estimated Resistance Training Proximity to Failure, Strength Gain, and Muscle Hypertrophy: A Series of Meta-Regressions

Supplementary materials:
<https://osf.io/7knsj/>
For correspondence:
zrobinson2019@fau.edu

Zac P. Robinson¹, Joshua C. Pelland¹, Jacob F. Remmert¹,
Martin C. Refalo², Ivan Jukic³, James Steele⁴, Michael C. Zourdos¹

¹ Department of Exercise Science and Health Promotion, Florida Atlantic University

² Institute for Physical Activity and Nutrition (IPAN), School of Exercise and Nutrition Sciences,
Deakin University, Geelong, VIC, Australia

³ Sport Performance Research Institute New Zealand (SPRINZ), Auckland University
of Technology, Auckland, New Zealand

⁴ Faculty of Sport, Health, and Social Sciences, Solent University, Southampton, UK

Please cite as: Robinson ZP, Pelland JC, Remmert JF, Refalo MC, Jukic I, Steele J, and Zourdos MC, (2023). Exploring the Dose-Response Relationship Between Estimated Resistance Training Proximity to Failure, Strength Gain, and Muscle Hypertrophy: A Series of Meta-Regressions.

All authors have read and approved this version of the manuscript. This article was last modified in May 2023.

Abstract

Background: The proximity to failure in which sets are terminated has gained attention in the scientific literature as a potentially key resistance training variable. Multiple meta-analyses have directly (i.e., failure versus not to failure) or indirectly (e.g., velocity loss, alternative set structures) evaluated the effect of proximity to failure on strength and muscle hypertrophy outcomes categorically; however, the dose response effects of proximity to failure have not been analyzed collectively in a continuous manner. **Objective:** To meta-analyze the aforementioned areas of relevant research, proximity to failure was quantified as the number of repetitions in reserve (RIR). Importantly, the RIR associated with each effect in the analysis was estimated based the available descriptions of the training interventions in each study. Data were extracted and a series of exploratory multi-level meta-regressions were performed for outcomes related to both strength and muscle hypertrophy. A range of other sensitivity analyses were also performed. All models were adjusted for the effects of load, method of volume equating, duration of intervention, and training status. **Results:** Linear-log models demonstrated the best fit for both strength and muscle hypertrophy outcomes. In all of best-fit models for strength, the confidence intervals of the marginal slopes for estimated RIR contained a null point estimate, indicating a negligible relationship with strength gains. However, in all of the best-fit models for muscle hypertrophy, the marginal slopes for estimated RIR were negative and their confidence intervals did not contain a null point estimate, indicating that changes in muscle size increased non-linearly as sets were terminated closer to failure. **Conclusions:** The dose-response relationship between proximity to failure and strength gain appears to differ from the relationship with muscle hypertrophy, with only the latter being meaningfully influenced by RIR. Strength gains were similar across a wide range of RIR, while muscle hypertrophy improves non-linearly as sets are terminated closer to failure. Researchers and practitioners should be aware that optimal proximity to failure may differ between strength and muscle hypertrophy outcomes, but caution is warranted when interpreting the present analysis as proximity to failure was estimated. Future studies deliberately designed to explore the continuous nature of the dose-response effects of proximity to failure in large samples should be considered.

Introduction

It is well established that the configuration of resistance training variables, such as set volume [1,2], load (percentage of one-repetition maximum (% of 1RM)) [3,4], and frequency [6] can impact resistance training outcomes. The impact of another variable, proximity to failure, operationally defined as the number of repetitions in reserve (RIR) prior to momentary failure (i.e., the inability to complete the concentric phase of a repetition despite maximal effort to do so) following set termination has gained recent attention in the scientific literature [7,8].

Despite the recent attention on this topic, the specific proximity to failure that maximizes various resistance training outcomes (i.e., muscular hypertrophy and strength gain) remains unclear. Indeed, recent meta-analyses from Grgic et al. [9] and Vieira et al. [10] reported no significant difference for muscle hypertrophy [SMD: 0.15 ($p = 0.237$); SMD: 0.59 ($p = 0.239$)], and strength gain [SMD: 0.01 ($p = 0.860$); SMD: 0.16 ($p = 0.566$)] when comparing volume-equated resistance training interventions in which sets were performed to or not to failure. Moreover, meta-analyses from Jukic et al. [11] and Davies et al. [12] show equivocal outcomes for muscle hypertrophy [SMD: -0.03 ($p = 0.708$); SMD: -0.05 ($p = 0.73$)] and strength gain [SMD: -0.06 ($p = 0.291$); SMD: -0.05 ($p = 0.56$)] when comparing the effects of traditional and alternative set structures; which alter proximity to failure via the manipulation of intra-set rest intervals [13].

Although the data from these meta-analyses [9–12] provide valuable insight into the influence of proximity to failure on resistance training outcomes, they investigate proximity to failure in a categorical (e.g., to failure versus not to failure; traditional versus alternative set structures) fashion despite its continuous nature. Additionally, there is ambiguity in the criteria for set termination [14], further limiting the applicability of these categorical comparisons. Fisher et al. [15] recently suggested investigating proximity to failure in a categorical fashion fails to inform the overall dose-response relationship between proximity to failure and resistance training outcomes. In other words, previous meta-analyses [9–12] have not attempted to ascertain the relationship between the number of RIR per set with muscle hypertrophy and strength gain, which limits the ability to make practical recommendations regarding proximity to failure. Specifically, if proximity to failure in resistance training is only examined in a categorical fashion, it cannot answer the question *“how far from failure should someone train to optimize muscle hypertrophy and strength gain?”*. To our knowledge, no meta-analysis to date has

examined proximity to failure as a continuous variable on muscle hypertrophy and strength gain.

The most commonly used method that aims to rectify the limitations of categorically examining proximity to failure is intra-set velocity loss (VL). Intra-set VL controls for proximity to failure by prescribing set termination once the concentric velocity of a repetition has declined by a predetermined percentage from a set's fastest (usually first) repetition (e.g., 70% of 1RM to 20%VL). In this way, higher VL thresholds terminate sets closer to failure, while lower VL thresholds terminate sets with a greater number of RIR. Recent meta-analyses [16–20], have indeed included meta-regressions examining the dose-response relationship between intra-set VL and various resistance training outcomes. However, while different VL thresholds lead to a different number of RIR, this method of set termination does not control for the number of repetitions performed in a set or the relative volume (repetitions x sets x % of 1RM) [21], which limits conclusions that can be made since training volume is related to muscle hypertrophy and strength gain [1,2]. For example, Pareja-Blanco et al. [22] compared the effects of training the Smith-machine squat to either a 20% or 40% VL threshold for eight weeks with the total number of sets and load equated. These intra-set VL thresholds resulted in the 20% and 40% groups training at an average mean propulsive velocity of 0.69 and 0.58 m·s⁻¹, respectively, suggesting the groups trained at different proximities to failure. However, the 40% VL group performed an average of 67% more repetitions throughout the study (40% VL: 310.5 ± 42 vs. 20% VL: 185.9 ± 22.2 repetitions); thus, the relative volume load was not equated. Studies exist [23,24] that have compared different VL thresholds and controlled for relative volume load but have only been included in one meta-analysis to date [16].

Therefore, this exploratory meta-analysis aimed to investigate the effect of resistance training proximity to failure on muscle hypertrophy and maximal strength gain by combining all of the aforementioned areas of research. To best investigate the effect of proximity to failure continuously and explore potential dose-response relationships, RIR was estimated for the training intervention associated with each effect.

Methods

This exploratory meta-analysis was performed without a systematic search and was not preregistered. First, all studies from the existing relevant meta-analyses [9–12,16–20] were

collected. Then, any additional studies we were aware of, or were discovered during data extraction, that met the inclusion criteria were also collected. To be included in this meta-analysis, studies had to: (1) be published in English and was either published in a peer-reviewed journal, on a pre-print repository, or as a MSc or PhD thesis; (2) ensure participants had no known medical conditions or injuries; (3) include either set and/or repetition volume-equated conditions; (4) include load-equated conditions (i.e., $\pm 5\%$ of 1RM); (5) compare at least two different proximities to failure; (6) include measurement(s) of maximal strength (i.e., isometric, isotonic, or isokinetic) and/or direct measurements of muscle hypertrophy (i.e., ultrasound, magnetic resonance imaging [MRI], etc.). Studies that were initially gathered but did not meet these inclusion criteria were excluded. This process was performed by ZR.

Data describing the population studied, specifications of the training intervention, and the outcomes of interest (e.g., muscle hypertrophy and strength) were extracted from studies found to be eligible. In the case that any necessary data were not reported, ZR emailed the authors of the manuscripts requesting the raw or mean values. If the authors did not reply within three months, we resorted to calculating the desired data based on the figures and tables (data was digitized using WebPlotDigitizer; v4.3, Ankit Rohatgi; <https://apps.automeris.io/wpd/>).

RIR Estimation

To operationally define proximity to failure as a continuous variable, the primary predictor in this meta-analysis was the estimated RIR to which training was performed by each group. Because self-reported RIR was not reported in almost all studies, specific procedures were followed to estimate RIR for every group included in the analysis. A detailed breakdown of the estimation of each study included in the analysis and all estimation equations utilized can be found in supplementary file 1. All initial estimations were performed by ZR, and these estimations were subsequently verified by JP, MZ, MR, and IJ. All conflicting predictions were due to manual error and were resolved upon re-estimation. The estimation process categorized studies into one of the following five subgroups: 1) groups training to failure, 2) groups reporting velocity, 3) groups reporting RIR, 4) groups reporting load and repetitions performed, and 5) groups training with alternative set structures.

Groups Training to Failure

The first category of studies consisted of those that included a group training to failure [25–52]. Groups that trained to failure were estimated, by definition, to have trained to 0 RIR. However, due to the ambiguity of the terminology utilized to describe “failure” [7], studies that provided a clear definition (e.g., momentary failure, concentric failure, etc.) and did not indicate participants terminated sets upon their own volition (i.e., volitional failure) were separately coded with an RIR of “-1” for the linear-spline meta-regression model to potentially tease out differences associated with failure definition. All other groups with RIR estimates of 0, including volitional failure, were coded as “0.”

In studies that included a group training to failure, if a group did not train to failure but performed sets at an equivalent load, subtraction of the repetitions performed per set was used to estimate RIR. For example, if the group training to failure (i.e., 0 RIR) performed 10 repetitions with 75% of 1RM and the group not training to failure performed 5 repetitions with 75% of 1RM, the number of RIR was estimated to be 5. This was also applied for groups training with a repetition maximum (e.g., 10RM). For example, if a group performed sets of 8 repetitions with a 10RM load, then the estimated RIR was 2.

Groups Reporting Velocity

The second category of studies reported velocity data that allowed for RIR estimations [22–24,53–65]. Utilizing the most representative citations available [66–69], equations were utilized or created to predict the maximum possible number of repetitions at a given load based on the repetitions performed and the intra-set VL. These equations were matched by exercise, loading range, training status, sex, and concentric intended velocity as closely as possible. After the maximum possible number of repetitions was estimated, subtraction could be performed from the repetitions per set provided.

For example, Pareja-Blanco and colleagues [22] reported a mean velocity loss of $41.9 \pm 1.9\%$ in the VL40 group. Using the reported average number of repetitions per set (6.5 ± 0.9) and load used (75% of 1RM), the maximum possible number of repetitions was predicted using the regression equations for the Smith-machine squat published by Rodriguez-Rosell et al. [66]. Specifically, at 70% of 1RM, the predicted maximum possible number of repetitions was 8.31. When subtracting the average number of repetitions performed per set reported by Pareja-Blanco et al. [22], the estimated RIR was 1.81. However, because the average load reported

was 75% of 1RM, we averaged the estimated RIR created from the equations for 70% and 80% of 1RM from Rodriguez-Rosell [66], resulting in the final estimated RIR for the VL40 group of 1.67. The prediction equations that were used for each velocity loss study can be found in supplementary file 1.

Groups Reporting RIR

The third category of studies directly reported RIR, and these values were utilized verbatim for the RIR estimations [37,70–72]. For example, if a group reported the average RIR throughout the study was 2.3, the estimated RIR was also 2.3. If a study reported RIR but also fell into one of the two prior categories (i.e., training to failure or provided velocity data), the other method was used to predict RIR due to its objectivity.

Groups Reporting Load and Repetitions Performed

The next category of studies reported the load and repetitions performed from which RIR estimations were derived [28,29,73–79]. Once again, prediction equations were utilized from the most representative citations available [66–68,80–82] to establish the maximum possible number of repetitions at a given load, to which subtraction of the repetitions performed per set could estimate RIR. These equations were matched by exercise, loading range, training status, sex, and concentric intended velocity as closely as possible. For example, Carneiro et al. [77] had untrained women perform sets of 4 repetitions at 90% of 1RM on the bilateral knee extension. Using the prediction equation created from the data of Hoeger et al. [83], it was estimated that approximately 4.65 repetitions are possible at 90% of 1RM. Thus, subtracting 4 repetitions per set from that estimation led to an estimated RIR of 0.65.

Alternatively, some studies reported load as a percentage of a repetition maximum (RM) other than a 1RM. In this case, the load relative to a 1RM (i.e., % of 1RM), was first predicted. Next, utilizing the predicted percentage of 1RM, the same steps could be followed as previously outlined. For example, Drinkwater and colleagues [73] had trained men perform sets of bench press with percentages of 6RM loads. Therefore, using the prediction equation from Chapman et al. [68] the percentage of 1RM associated with 100% of 6RM was predicted, which equated to 89% of 1RM. Then, the percentages of this value used in Drinkwater et al. [73] were applied to calculate the load per set (e.g., 70% of 89% (6RM) = 62.3% of 1RM). The same steps outlined in the previous paragraph were then followed to predict the maximum possible repetitions at

the given load. Finally, the number of repetitions performed per set was subtracted from the predicted maximum possible number of repetitions to estimate RIR.

Groups Training with Alternative Set Structures

For groups training with alternative set structures [26,28–31,33,35,38,40,41,50,64,65,73–79,84], a determination needed to be made as to what constitutes a “set” from which proximity to failure could be evaluated. We decided to treat each group of repetitions performed with any intra-set or inter-set rest between them as an individual set. For rest-pause groups, each of these sets was performed to failure, making the RIR estimation simple (i.e., 0 RIR) [76]. For cluster and rest redistribution groups, this RIR estimation method was chosen because of these set structures’ ability to maintain repetition performance, and thus RIR, compared to traditional set structures.

For example, Iglesias-Soler et al. [85] compared the total number of repetitions performed with a 4RM load on the Smith-machine squat between traditional and cluster set structures in a crossover design. The traditional set condition performed 3 sets to failure, whereas the cluster set condition performed as many repetitions as possible with an individualized inter-repetition rest period (i.e., single-repetition sets). The authors observed that participants were able to increase the total number of repetitions performed by approximately 5-fold with a cluster set structure (Cluster: 45.5 ± 32 repetitions; Traditional: 9.33 ± 1.87 repetitions), indicating RIR was likely maintained for many of the early repetitions. Due to this evidence, along with variability in the initial proximity to failure of each set, intra- and inter-set rest intervals, the load used, sex of the participants, and exercise selection, we determined this estimating RIR for cluster and rest redistribution groups in this manner to be the best course of action.

To illustrate, Farinas et al. [40] investigated the effects of set structure within a unilateral knee extension training program. Participants in the traditional group performed 4 sets of 8 repetitions with a 10RM load and 3-minute inter-set rest periods. Using an equivalent load (i.e., 10RM), the alternative set structure group instead performed 1 repetition every 17.4 seconds until 32 total repetitions were completed. The estimated RIR for each group was calculated by subtracting the number of repetitions performed in each single-repetition set from the maximum possible number of repetitions. Thus, the RIR estimation for the traditional group would be 2 and the alternative set structure group would be 9.

Special Cases

Throughout the estimation process, special case rules were applied to a few groups. Specifically, in these cases, even though a group fell into one of the aforementioned categories, the estimation approach led to subjectively implausible RIR values. These alternative approaches are outlined in supplementary file 1.

Additional Details

For each group in the analysis, the estimated RIR was used to represent the average proximity to failure at which sets were terminated. Specifically, averages were developed for both the “muscle” and the “exercise.” For the “muscle” RIR, if a study included multiple exercises, those that primarily trained the muscle site measured or were prime movers in the exercise or joint action tested counted towards the estimation. For example, if a lower body training program included the barbell back squat, knee extension machine, and knee flexion machine, only the RIR estimations of the barbell back squat and knee extension machine would be averaged for quadriceps specific outcomes (i.e., strength or muscle hypertrophy). Alternatively, for the “exercise” RIR, only sets performed on the same exercise that was tested were counted towards the estimation; thus, this is only relevant for dynamic strength outcomes. For example, in the same lower body training program that included the barbell back squat, knee extension machine, and knee flexion machine, only the RIR estimations of the barbell back squat would be used for the exercise RIR. When applicable, the RIR estimations of multiple sets performed by a single group were averaged for the final estimation. Finally, if a RIR estimate was negative, it was recorded as “0.”

Statistical Analysis

This exploratory meta-analysis was performed using the *‘metafor’* package [86] in R language and environment for statistical computing (v 4.0.2; R Core Team, <https://www.r-project.org/>). The extracted dataset, analysis scripts, models summaries, and supplementary materials are available on the Open Science Framework (<https://osf.io/7knsj/>). Given the goal of this analysis, we have opted to avoid dichotomizing our findings and therefore did not employ traditional null hypothesis significance testing [87]. Rather, we took an estimation-based approach in which effect estimates and their precision were interpreted cautiously and probabilistically [88]. As the included studies often had multiple groups and reported effects within these groups for multiple outcomes, we opted to calculate effect sizes in a nested structure.

Therefore, multi-level mixed-effects models [89] with cluster-robust variance estimation [90] were performed with study, group, and observation included as nested random effects in the model (i.e., observations were nested within groups which were nested within studies). Effects were weighted by inverse sampling variance to account for the observation-level, within-study, and between-study variance. Models were constructed with effect sizes, and variances thereof, calculated as both standardized mean change and response ratio using the 'escalc' function [91,92]. Specifically, standardized mean changes were calculated as the difference between post-test and pre-test means, divided by the pre-test standard deviation with an adjustment (i.e., C) for a small sample bias. Alternatively, response ratios were calculated as the natural logarithm of the ratio of post-test and pre-test means, which were later exponentiated (i.e., e^x) and thereby converted to percentage change scores to aid practical interpretation. Formulas for each effect size and their variances can be seen below:

$$SMC = C \left(\frac{M_{post} - M_{pre}}{SD_{pre}} \right); C = 1 - \frac{3}{4(n-1) - 1}$$

$$var(SMC) = \frac{2(1-r)}{n} + \frac{(SMC)^2}{2 \cdot n}$$

$$RR = \ln \left(\frac{M_{post}}{M_{pre}} \right); var(RR) = \frac{(SD_{post})^2}{n \cdot M_{post}} + \frac{(SD_{pre})^2}{n \cdot M_{pre}} + \frac{2r \cdot SD_{post} \cdot SD_{pre}}{M_{post} \cdot M_{pre} \cdot n}$$

$$RR_{exp} = (e^{RR} - 1) * 100$$

No studies reported the pre-intervention to post-intervention correlations required to determine the variance. Therefore, the available data were used to retroactively calculate pre-to-post correlations if possible [93]. Then, we meta-analyzed these approximated correlation coefficients (i.e., Fishers r-to-z transformed correlation coefficient) and imputed this estimate for the studies where we were unable to obtain the required data. Since all meta-analytic models included moderators, statistical heterogeneity was evaluated using I², which represents the remaining variance that is not already accounted by the moderators [94]. This heterogeneity was then partitioned across the three levels of the nested random effects (i.e., study, group, and observation). Additionally, marginal and conditional R² were calculated to

quantify the proportion of variance explained by only the fixed effects and the sum of the fixed and random effects, respectively [95]. Both I^2 and R^2 were calculated using the '*orchaRd*' package for multi-level models [96]. To account for potential non-linear dose-response relationships between proximity to failure and resistance training outcomes the following models were fit and subsequently compared with the '*performance*' package [97], utilizing the composite rank to describe the best fitting model for each outcome (i.e., strength and muscle hypertrophy):

- 1) Linear
- 2) Linear Spline (knot at 0 RIR)
- 3) Linear-log
- 4) Quadratic

All models included the following fixed effects and their interactions: 1) RIR, 2) Load per set, 3) Method of volume equating (set, repetition, or both), while the duration of the training intervention (continuous) and training status of the participants (binary categorical) were included as covariates¹. Estimated marginal means (and their slopes) with 95% compatibility intervals (confidence and prediction) were extracted for both main effects (adjusted for all predictors except RIR) and interactions effects of interest between RIR, load per set, and the method of volume equating using the '*emmeans*' package [98]. For RIR, estimates were extracted at 0 to 23 RIR to represent the range of values observed in the dataset. For load per set, estimates were extracted at 40, 60, and 80% of 1RM to represent "low", "moderate", and "high" loads, respectively. For the method of volume equating, estimates were extracted for "set" and "repetition" equated effects. Finally, after identifying a potentially meaningful interaction with load, estimates were extracted in increments of 2.5% of 1RM to identify thresholds where the marginal slopes changed in directionality (e.g., positive to negative) and magnitude (e.g., when the confidence interval of the estimate no longer contained to a null value) compared to the main effect.

Following the determination of the best fit models for each outcome (i.e., strength or muscle hypertrophy) and effect size (i.e., SMC or RR), additional moderator analyses were performed

¹ Upon fitting all of the linear spline models, the three-way interaction of RIR, load, and volume could not be fit. Thus, these models were fit with only the corresponding two-way interactions specified.

to evaluate the influence of a variety of factors related to study design and participant characteristics (supplementary file 2). Specifically, separate models were fit for each moderator that maintained the same structure as the best fit models from the main analysis, but also included an interaction term between estimated RIR and the moderator of interest. Finally, after inspecting the data, it appeared that some studies with very high RIR estimations may have been disproportionately influencing the models. Thus, a sensitivity analysis was performed where all models were refit only with effect sizes from groups that were estimated to train with less than or equal to 10 RIR.

Results

Study Characteristics

A breakdown of all 55 studies included in this analysis can be found in supplementary file 0. On average, training interventions lasted 8.28 ± 2.36 weeks and participants were 27.85 ± 12.87 years old. A visual summary of the training interventions from the included studies can be seen in Figure 1. Additionally, tables summarizing study characteristics can be seen in supplementary file 2. The most frequently occurring (i.e., mode) values of the primary training variables for effects included in the strength models were: volume- 6 sets per week; load- 75% of 1RM; and frequency- 2 sessions per week. The average values for these metrics were 9.58 ± 4.49 sets per week, $72.03 \pm 13.29\%$ of 1RM, and 2.23 ± 0.48 sessions per week. For the effects in the muscle hypertrophy models, the most frequently occurring values were 6 sets per week, 85% of 1RM, and 2 sessions per week. The average values for these metrics were 9.69 ± 4.62 sets per week, $72.22 \pm 14.57\%$ of 1RM, and 2.07 ± 0.38 sessions per week.

Visual Summaries of Training Interventions Included in Meta-regression Models




Figure 1: Visual summary of training interventions included in the analysis. Data are presented as raincloud plots with each data point representing an effect. Adjusted sets per week = Number of sets performed per week while alternative set structure groups were set to the same number of sets as the traditional set structure group in the same study.

Primary Analysis

The following sections will present the results of all meta-regression models. Specifically, for each model, we will evaluate the marginal slope for the main effect of estimated RIR (i.e., the slope after adjusting for load, method of volume equating and the other covariates in the model) and the marginal slope of any relevant interactions (i.e., two-way or three-way). Finally, for interactions between estimated RIR and load, we will identify thresholds where the directionality (i.e., positive or negative) and magnitude (i.e., does the confidence interval contain a null point estimate?) of the relationship between estimated RIR and the outcome variable change. Full model summary tables, comparisons thereof, and all extracted estimates are located at <https://osf.io/7knsj/>.

Strength Outcomes

The multi-level meta-regression models for strength included 242 total effects from 55 studies. Model comparisons revealed that the linear-log model was the best fit with effects as a standardized mean change (Figure 2: $R^2_{\text{marginal}} = 14.05\%$; $R^2_{\text{conditional}} = 79.26\%$; $I^2_{\text{study}} = 57\%$; $I^2_{\text{study/group}} = 8.8\%$; $I^2_{\text{study/group/effect}} = 20.93\%$).

Meta-regressions from Linear-log Multilevel Model for Maximal Strength (Standardized Mean Change)




Figure 2: Best fit (linear-log) multi-level meta-regression for strength outcomes analyzed as a standardized mean change. Data are presented as estimated marginal means (solid line) with 95% compatibility intervals (dark band = confidence, light band = prediction) after adjusting for load, method of volume equating, intervention duration, and training status of the participants. Colored circles represent the effect size of each observation included in the analysis, with the size of each circle representing its weight determined by inverse variance weighting. The main effect for estimated RIR is presented at the mean of continuous fixed effects (i.e., load and intervention duration) and averaged across categorical fixed effects (i.e., method of volume equating and training status). Additionally, effects extracted at multiple levels of load (i.e., 40, 60, and 80% of 1RM) and method of volume equating (i.e., set and repetition volume equated) can be seen in the rightward panels to visualize potential interactions with estimated RIR.

When adjusted for load and the method in which volume was equated, the marginal slope for estimated RIR was positive and the confidence interval contained a null point estimate ($\beta = 0.01318$ [95% CI: -0.00068, 0.02703; 95% PI: -0.67391, 0.70026]). The slope indicates that strength gains improve negligibly in a non-linear fashion as sets are terminated farther from failure. Moreover, there was evidence of a three-way interaction between estimated RIR, load, and the method in which volume was equated, as the confidence interval of the marginal slope did not contain a null point estimate ($\beta = 0.0012$ [95% CI: 0.00004, 0.00237; 95% PI: -0.68575, 0.68815]). The positive slope indicates that when sets are equated and load increases, better

strength gains are expected farther from failure; however when repetitions are equated, and load increases, worse strength gains are expected farther from failure.

With sets equated, loads <52.5% of 1RM resulted in negative slopes, while heavier loads resulted in positive slopes. All slopes contained a null point estimate within the confidence interval. These findings indicate that training farther from failure resulted in more favorable strength outcomes with heavy loads when set volume is equated. Alternatively, with repetitions equated, loads >75% of 1RM exhibited negative slopes, while lighter loads resulted in positive slopes. All slopes contained a null point estimate within the confidence interval. Therefore, training farther from failure resulted in more favorable strength outcomes with light loads when repetition volume is equated.

Model comparisons revealed that the linear-log model remained the best fit with effects as a response ratio (Figure 3: $R^2_{marginal} = 28.8\%$; $R^2_{conditional} = 58.77\%$; $I^2_{study} = 38.22\%$; $I^2_{study/group} = 1.06\%$; $I^2_{study/group/effect} = 54.03\%$).

Meta-regressions from Linear-log Multilevel Model for Maximal Strength (Exponentiated Response Ratio)




Figure 3: Best fit (linear-log) multi-level meta-regression for strength outcomes analyzed as a response ratio. Data are presented as exponentiated estimated marginal means (solid line) with 95% compatibility intervals (dark band = confidence, light band = prediction) after adjusting for load, method of volume equating, intervention duration, and training status of the participants. Colored circles represent the effect size of each observation included in the analysis, with the size of each circle representing its weight determined by inverse variance weighting. The main effect for estimated RIR is presented at the mean of continuous fixed effects (i.e., load and intervention duration) and averaged across categorical fixed effects (i.e., method of volume equating and training status). Additionally, effects extracted at multiple levels of load (i.e., 40, 60, and 80% of 1RM) and method of volume equating (i.e., set and repetition volume equated) can be seen in the rightward panels to visualize potential interactions with estimated RIR.

When adjusted for load and the method in which volume was equated, the marginal slope for estimated RIR was positive and the confidence interval contained a null point estimate ($\beta_{exp} = 0.12736$ [95% CI: -0.1478, 0.40327; 95% PI: -12.90797, 15.11372]). The slope indicates that strength gains improve negligibly in a non-linear fashion as sets are terminated farther from failure. There was no convincing evidence of a three-way interaction between estimated RIR, load, and the method in which volume was equated ($\beta_{exp} = 0.01249$ [95% CI: -0.00883, 0.03382; 95% PI: -13.00553, 14.97856]) or a two-way interaction between estimated RIR and the method in which volume was equated ($\beta_{exp} = -0.31246$ [95% CI: -0.90091, 0.27948; 95% PI:

-13.29907, 14.61936]), as the confidence intervals of the marginal slopes contained a null point estimate.

However, there was evidence of a two-way interaction between estimated RIR and load, as the confidence interval of the marginal slope did not contain a null point estimate ($\beta_{exp} = -0.07149$ [95% CI: -0.12114, -0.02181; 95% PI: -13.07865, 14.8821]). The negative slope indicates that training farther from failure results in greater strength gains as load decreases. Specifically, loads >72.5% of 1RM exhibited negative slopes but only loads >80% of 1RM did not contain a null point estimate within the confidence interval. Lighter loads exhibited positive slopes but only loads <70% of 1RM did not contain a null point estimate within the confidence interval. Therefore, training farther from failure resulted in favorable strength gains for lighter rather than heavier loads.

Muscle Hypertrophy Outcomes

The multi-level meta-regression models for muscle hypertrophy included 139 total effects from 26 studies. Model comparisons revealed that the linear-log model was the best fit with effects as a standardized mean change (Figure 4: $R^2_{marginal} = 19.39\%$; $R^2_{conditional} = 77.2\%$; $I^2_{study} = 62.2\%$; $I^2_{study/group} = 1.77\%$; $I^2_{study/group/effect} = 25.23\%$).

Meta-regressions from Linear-log Multilevel Model for Muscle Hypertrophy (Standardized Mean Change)




Figure 4: Best fit (linear-log) multi-level meta-regression for muscle hypertrophy outcomes analyzed as a standardized mean change. Data are presented as estimated marginal means (solid line) with 95% compatibility intervals (dark band = confidence, light band = prediction) after adjusting for load, method of volume equating, intervention duration, and training status of the participants. Colored circles represent the effect size of each observation included in the analysis, with the size of each circle representing its weight determined by inverse variance weighting. The main effect for estimated RIR is presented at the mean of continuous fixed effects (i.e., load and intervention duration) and averaged across categorical fixed effects (i.e., method of volume equating and training status). Additionally, effects extracted at multiple levels of load (i.e., 40, 60, and 80% of 1RM) and method of volume equating (i.e., set and repetition volume equated) can be seen in the rightward panels to visualize potential interactions with estimated RIR.

When adjusted for load and the method in which volume was equated, the marginal slope for estimated RIR was negative and the confidence interval did not contain a null point estimate ($\beta = -0.02641$ [95% CI: -0.04075, -0.01208; 95% PI: -0.60151, 0.54869]). The negative slope indicates that muscle hypertrophy improves in a non-linear fashion as sets are terminated closer to failure. There was no convincing evidence of a three-way interaction between estimated RIR, load, and the method in which volume was equated ($\beta = -0.00019$ [95% CI: -0.0014, 0.00102; 95% PI: -0.57511, 0.57473]) or a two-way interaction between estimated RIR and the method in which volume was equated ($\beta = 0.00652$ [95% CI: -0.03497, 0.04802; 95% PI:

-0.56989, 0.58294]), as the confidence intervals of the marginal slopes contained a null point estimate.

However, there was evidence of a two-way interaction between estimated RIR and load, as the confidence interval of the marginal slope did not contain a null point estimate ($\beta = 0.00413$ [95% CI: 0.00283, 0.00543; 95% PI: -0.57079, 0.57905]). The positive slope indicates that training closer to failure has less of an impact on muscle hypertrophy when training with heavy loads. Specifically, loads >77.5% of 1RM exhibited positive slopes but only loads >82.5% of 1RM did not contain a null point estimate within the confidence interval. All loads <75% of 1RM exhibited negative slopes and did not contain a null point estimate within the confidence interval. Therefore, training closer to failure is more beneficial for muscle hypertrophy with lighter rather than heavier loads.

Model comparisons revealed that the linear-log model remained the best fit with effects as a response ratio (Figure 5: $R_{marginal}^2 = 26.97\%$; $R_{conditional}^2 = 68.49\%$; $I_{study}^2 = 46.51\%$; $I_{study/group}^2 = 4.5\%$; $I_{study/group/effect}^2 = 38.7\%$)

Meta-regressions from Linear-log Multilevel Model for Muscle Hypertrophy (Exponentiated Response Ratio)




Figure 5: Best fit (linear-log) multi-level meta-regression for muscle hypertrophy outcomes analyzed as a response ratio. Data are presented as exponentiated estimated marginal means (solid line) with 95% compatibility intervals (dark band = confidence, light band = prediction) after adjusting for load, method of volume equating, intervention duration, and training status of the participants. Colored circles represent the effect size of each observation included in the analysis, with the size of each circle representing its weight determined by inverse variance weighting. The main effect for estimated RIR is presented at the mean of continuous fixed effects (i.e., load and intervention duration) and averaged across categorical fixed effects (i.e., method of volume equating and training status). Additionally, effects extracted at multiple levels of load (i.e., 40, 60, and 80% of 1RM) and method of volume equating (i.e., set and repetition volume equated) can be seen in the rightward panels to visualize potential interactions with estimated RIR.

When adjusted for load and the method in which volume was equated, the marginal slope for estimated RIR was negative and the confidence interval did not contain a null point estimate ($\beta_{exp} = -0.54654$ [95% CI: -0.80344, -0.28898; 95% PI: -10.62288, 10.6658]). The negative slope indicates that muscle hypertrophy improves in a non-linear fashion as sets are terminated closer to failure. There was no convincing evidence of a three-way interaction between estimated RIR, load, and the method in which volume was equated ($\beta_{exp} = 0.00263$ [95% CI: -0.02292, 0.02818; 95% PI: -10.12656, 11.27343]) or a two-way interaction between estimated RIR and the method in which volume was equated ($\beta_{exp} = 0.22383$ [95% CI: -0.46749, 0.91994;

95% PI: -9.94792, 11.54452]), as the confidence intervals of the marginal slopes contained a null point estimate.

However, there was evidence of a two-way interaction between estimated RIR and load, as the confidence interval of the marginal slope did not contain a null point estimate ($\beta_{exp} = 0.10289$ [95% CI: 0.0672, 0.13859; 95% PI: -10.03648, 11.38502]). The positive slope indicates that training closer to failure has less of an impact on muscle hypertrophy when training with heavy loads. Specifically, loads >77.5% of 1RM exhibited positive slopes but only loads >80% of 1RM did not contain a null point estimate within the confidence interval. All loads <75% of 1RM exhibited negative slopes and did not contain a null point estimate within the confidence interval. Therefore, training closer to failure is more beneficial for muscle hypertrophy with lighter rather than heavier loads.

Sensitivity Analysis

Strength Outcomes

Upon refitting the multi-level meta-regression models for strength only with the effects with an RIR estimate of <10, they included 231 total effects from 54 studies. Model comparisons revealed that the linear-log model was the best fit with effects as a standardized mean change (Figure 6: $R^2_{marginal} = 15.14\%$; $R^2_{conditional} = 76.36\%$; $I^2_{study} = 57.78\%$; $I^2_{study/group} = 4.7\%$; $I^2_{study/group/effect} = 24.13\%$).

Meta-regressions from Linear-log Multilevel Model for Maximal Strength (Standardized Mean Change)




Figure 6: Best fit (linear-log) multi-level meta-regression for strength outcomes analyzed as a standardized mean change after effects with >10 estimated RIR were removed. Data are presented as estimated marginal means (solid line) with 95% compatibility intervals (dark band = confidence, light band = prediction) after adjusting for load, method of volume equating, intervention duration, and training status of the participants. Colored circles represent the effect size of each observation included in the analysis, with the size of each circle representing its weight determined by inverse variance weighting. The main effect for estimated RIR is presented at the mean of continuous fixed effects (i.e., load and intervention duration) and averaged across categorical fixed effects (i.e., method of volume equating and training status). Additionally, effects extracted at multiple levels of load (i.e., 40, 60, and 80% of 1RM) and method of volume equating (i.e., set and repetition volume equated) can be seen in the rightward panels to visualize potential interactions with estimated RIR.

When adjusted for load and the method in which volume was equated, the marginal slope for estimated RIR was positive and the confidence interval contained a null point estimate ($\beta = 0.01007$ [95% CI: -0.00719, 0.02734; 95% PI: -0.66609, 0.68624]). The slope indicates that strength gains improve negligibly in a non-linear fashion as sets are terminated farther from failure. There was no convincing evidence of a three-way interaction between estimated RIR, load, and the method in which volume was equated ($\beta = 0.00205$ [95% CI: -0.00078, 0.00487; 95% PI: -0.67391, 0.678]) or a two-way interaction between estimated RIR and the method in which volume was equated ($\beta = -0.0216$ [95% CI: -0.0676, 0.02441; 95% PI: -0.69911, 0.65591]), as the confidence intervals of the marginal slopes contained a null point estimate.

However, there was evidence of a two-way interaction between estimated RIR and load, as the confidence interval of the marginal slope did not contain a null point estimate ($\beta = -0.00613$ [95% CI: -0.01019, -0.00208; 95% PI: -0.68209, 0.66983]). The negative slope indicates that training farther from failure results in greater strength gains as load decreases. Specifically, loads >72.5% of 1RM exhibited negative slopes but only loads >80% of 1RM did not contain a null point estimate. Lighter loads exhibited positive slopes but only loads <72.5% of 1RM did not contain a null point estimate. Therefore, training farther from failure resulted in favorable strength gains for lighter rather than heavier loads.

Model comparisons revealed that the linear-log model remained the best fit with effects as a response ratio (Figure 7: $R^2_{\text{marginal}} = 30.22\%$; $R^2_{\text{conditional}} = 57.74\%$; $I^2_{\text{study}} = 36.86\%$; $I^2_{\text{study/group}} = 0\%$; $I^2_{\text{study/group/effect}} = 56.61\%$).

Meta-regressions from Linear-log Multilevel Model for Maximal Strength (Exponentiated Response Ratio)




Figure 7: Best fit (linear-log) multi-level meta-regression for strength outcomes analyzed as a response ratio after effects with >10 estimated RIR were removed. Data are presented as exponentiated estimated marginal means (solid line) with 95% compatibility intervals (dark band = confidence, light band = prediction) after adjusting for load, method of volume equating, intervention duration, and training status of the participants. Colored circles represent the effect size of each observation included in the analysis, with the size of each circle representing its' weight determined by inverse variance weighting. The main effect for estimated RIR is presented at the mean of continuous fixed effects (i.e., load and intervention duration) and averaged across categorical fixed effects (i.e., method of volume equating and training status). Additionally, effects extracted at multiple levels of load (i.e., 40, 60, and 80% of 1RM) and method of volume equating (i.e., set and repetition volume equated) can be seen in the rightward panels to visualize potential interactions with estimated RIR.

When adjusted for load and the method in which volume was equated, the marginal slope for estimated RIR was positive and the confidence interval contained a null point estimate ($\beta_{exp} = 0.11897$ [95% CI: -0.22807, 0.46721; 95% PI: -12.98599, 15.19763]). The slope indicates that strength gains improve negligibly in a non-linear fashion as sets are terminated farther from failure. There was no convincing evidence of a three-way interaction between estimated RIR, load, and the method in which volume was equated ($\beta_{exp} = 0.04841$ [95% CI: -0.01439, 0.11124; 95% PI: -13.0437, 15.11166]) or a two-way interaction between estimated RIR and the method in which volume was equated ($\beta_{exp} = -0.27145$ [95% CI: -1.055, 0.5183; 95% PI: -

13.34079, 14.76891]), as the confidence intervals of the marginal slopes contained a null point estimate.

However, there was evidence of a two-way interaction between estimated RIR and load, as the confidence interval of the marginal slope did not contain a null point estimate ($\beta_{exp} = -0.09178$ [95% CI: -0.15001, -0.0335; 95% PI: -13.16552, 14.95034]). The negative slope indicates that training farther from failure results in greater strength gains as load decreases. Specifically, loads >72.5% of 1RM exhibited negative slopes but only loads >77.5% of 1RM did not contain a null point estimate. Lighter loads exhibited positive slopes but only loads <70% of 1RM did not contain a null point estimate. Therefore, training farther from failure resulted in favorable strength gains for lighter rather than heavier loads.

Finally, upon refitting all models with the RIR and load per set of the specific exercise evaluated as a dynamic strength outcome, a linear-log model remained the best fit, regardless of effect size. All estimates were similar to the primary models.

Muscle Hypertrophy Outcomes

Upon refitting the multi-level meta-regression models for muscle hypertrophy only with the effects with an RIR estimate of <10, they included 136 total effects from 26 studies. Model comparisons revealed that the linear-log model was the best fit with effects as a standardized mean change (Figure 8: $R^2_{marginal} = 18.62\%$; $R^2_{conditional} = 77.2\%$; $I^2_{study} = 62.66\%$; $I^2_{study/group} = 1.88\%$; $I^2_{study/group/effect} = 25.12\%$).

Meta-regressions from Linear-log Multilevel Model for Muscle Hypertrophy (Standardized Mean Change)




Figure 8: Best fit (linear-log) multi-level meta-regression for muscle hypertrophy outcomes analyzed as a standardized mean change after effects with >10 estimated RIR were removed. Data are presented as estimated marginal means (solid line) with 95% compatibility intervals (dark band = confidence, light band = prediction) after adjusting for load, method of volume equating, intervention duration, and training status of the participants. Colored circles represent the effect size of each observation included in the analysis, with the size of each circle representing its weight determined by inverse variance weighting. The main effect for estimated RIR is presented at the mean of continuous fixed effects (i.e., load and intervention duration) and averaged across categorical fixed effects (i.e., method of volume equating and training status). Additionally, effects extracted at multiple levels of load (i.e., 40, 60, and 80% of 1RM) and method of volume equating (i.e., set and repetition volume equated) can be seen in the rightward panels to visualize potential interactions with estimated RIR.

When adjusted for load and the method in which volume was equated, the marginal slope for estimated RIR was negative and the confidence interval did not contain a null point estimate ($\beta = -0.02753$ [95% CI: -0.04318, -0.01189; 95% PI: -0.61445, 0.55939]). The slope indicates that muscle hypertrophy improves in a non-linear fashion as sets are terminated closer to failure. There was no convincing evidence of a three-way interaction between estimated RIR, load, and the method in which volume was equated ($\beta = -0.00093$ [95% CI: -0.00424, 0.00238; 95% PI: -0.58765, 0.58579]) or a two-way interaction between estimated RIR and the method in which volume was equated ($\beta = 0.00004$ [95% CI: -0.04503, 0.04512; 95% PI:

-0.5884, 0.58848]), as the confidence intervals of the marginal slopes contained a null point estimate.

However, there was evidence of a two-way interaction between estimated RIR and load, as the confidence interval of the marginal slope did not contain a null point estimate ($\beta = 0.00468$ [95% CI: 0.00295, 0.00641; 95% PI: -0.58204, 0.59139]). The positive slope indicates that training closer to failure has less of an impact when training with heavy loads. Specifically, loads >77.5% of 1RM exhibited positive slopes but only loads >82.5% of 1RM did not contain a null point estimate within the confidence interval. All loads <77.5% of 1RM exhibited negative slopes and did not contain a null point estimate within the confidence interval. Therefore, training closer to failure is more beneficial for muscle hypertrophy with lighter rather than heavier loads.

Model comparisons revealed that the linear-log model remained the best fit with effects as a response ratio (Figure 9: $R_{marginal}^2 = 26.13\%$; $R_{conditional}^2 = 68.06\%$; $I_{study}^2 = 47.24\%$; $I_{study/group}^2 = 3.87\%$; $I_{study/group/effect}^2 = 38.94\%$).

Meta-regressions from Linear-log Multilevel Model for Muscle Hypertrophy (Exponentiated Response Ratio)




Figure 9: Best fit (linear-log) multi-level meta-regression for muscle hypertrophy outcomes analyzed as a response ratio after effects with >10 estimated RIR were removed. Data are presented as exponentiated estimated marginal means (solid line) with 95% compatibility intervals (dark band = confidence, light band = prediction) after adjusting for load, method of volume equating, intervention duration, and training status of the participants. Colored circles represent the effect size of each observation included in the analysis, with the size of each circle representing its weight determined by inverse variance weighting. The main effect for estimated RIR is presented at the mean of continuous fixed effects (i.e., load and intervention duration) and averaged across categorical fixed effects (i.e., method of volume equating and training status). Additionally, effects extracted at multiple levels of load (i.e., 40, 60, and 80% of 1RM) and method of volume equating (i.e., set and repetition volume equated) can be seen in the rightward panels to visualize potential interactions with estimated RIR.

When adjusted for load and the method in which volume was equated, the marginal slope for estimated RIR was negative and the confidence interval did not contain a null point estimate ($\beta_{exp} = -0.58529$ [95% CI: -0.82318, -0.34682; 95% PI: -10.77438, 10.76735]). The slope indicates that muscle hypertrophy improves in a non-linear fashion as sets are terminated closer to failure. There was no convincing evidence of a three-way interaction between estimated RIR, load, and the method in which volume was equated ($\beta_{exp} = -0.02344$ [95% CI: -0.09255, 0.04571; 95% PI: -10.26793, 11.39064]) or a two-way interaction between estimated RIR and the method in which volume was equated ($\beta_{exp} = 0.03964$ [95% CI: -0.59284, 0.67614;

95% PI: -10.22781, 11.48139]), as the confidence intervals of the marginal slopes contained a null point estimate.

However, there was evidence of a two-way interaction between estimated RIR and load, as the confidence interval of the marginal slope did not contain a null point estimate ($\beta_{exp} = 0.11787$ [95% CI: 0.07461, 0.16115; 95% PI: -10.14098, 11.54794]). The positive slope indicates that training closer to failure has less of an impact when training with heavy loads. Specifically, loads >77.5% of 1RM exhibited positive slopes but only loads >80% of 1RM did not contain a null point estimate within the confidence interval. All loads <77.5% of 1RM exhibited negative slopes and did not contain a null point estimate within the confidence interval. Therefore, training closer to failure is more beneficial for muscle hypertrophy with lighter rather than heavier loads.

Other Potentially Influential Moderators

The following sections will present the results of the additional moderator analyses. We evaluated the marginal slopes of the two-way interactions between estimated RIR and the moderators of interest. Specifically, we will identify moderators that alter the directionality (i.e., positive or negative) and magnitude (i.e., does the confidence interval contain a null point estimate?) of the relationship between estimated RIR and the outcome variable. Estimates from all the additional moderator analyses are located at <https://osf.io/7knsj/>.

Strength Outcomes

There were multiple moderators that seemed to alter the dose-response relationship between estimated RIR and strength gains with effects as standardized mean change. The following moderators exhibited positive slopes that were greater in magnitude compared the main effect and did not contain a null point estimate within the confidence interval: studies that were both set and repetition volume equated, within-participant designs, lower-body outcomes, groups that trained with multi-joint exercises, traditional set structures, isotonic strength tests, and non-1RM isotonic strength tests. These slopes suggest a shift from a negligible to a meaningful relationship in which even greater strength gains are observed training farther from failure compared to the main effect. Alternatively, isokinetic strength tests exhibited negative slopes that were lesser in magnitude compared the main effect and did not

contain a null point estimate within the confidence interval. This slope suggests a shift from a negligible to a meaningful relationship in which worse strength gains are observed training farther from failure compared to the main effect.

When strength outcomes were evaluated with effects as a response ratio, there were two remaining moderators that seemed to alter the dose-response relationship. Within-participant designs exhibited positive slopes that were greater in magnitude compared the main effect. This slopes suggests a shift from a negligible to a meaningful relationship in which even greater strength gains are observed training farther from failure compared to the main effect. In contrast, isokinetic strength tests exhibited negative slopes that were lesser in magnitude compared the main effect and did not contain a null point estimate within the confidence interval. This slope suggests a shift from a negligible to a meaningful relationship in which worse strength gains are observed training farther from failure compared to the main effect.

Muscle Hypertrophy Outcomes

There were multiple moderators that seemed to alter the dose-response relationship between estimated RIR and muscle hypertrophy with effects as standardized mean change. The following moderators exhibited slopes that were lesser in magnitude compared the main effect and contained a null point estimate within the confidence interval: studies that were set volume equated, repetition volume equated, higher biological age, trained with multi-joint exercises, trained with single-joint exercises, did not utilize progressive overload, included a failure definition, alternative set structures, higher frequency per muscle group, longer intervention duration, higher percentages of male participants, within-participant designs, and a higher number of adjusted sets per week. These slopes suggest a shift from a meaningful to a negligible relationship in which there was less of an advantage to training closer to failure compared to the main effect. There were no moderators that resulted in positive slopes that did not contain a null point estimate within the confidence interval.

When muscle hypertrophy outcomes were evaluated with effects as a response ratio, there were multiple remaining moderators that seemed to alter the dose-response relationship. The following moderators exhibited slopes that were lesser in magnitude compared the main effect and contained a null point estimate within the confidence interval: studies that were set volume equated, repetition volume equated, untrained participants, higher biological age, trained with multi-joint exercises, trained with single-joint exercises, concurrent training

interventions, included a failure definition, alternative set structures, did not utilize maximal concentric intended velocity, higher frequency per muscle group, longer intervention duration, higher percentages of male participants, within-participant designs, and a higher number of adjusted sets per week.. These slopes suggest a shift from a meaningful to a negligible relationship in which there was less of an advantage to training closer to failure compared to the main effect. There were no moderators that resulted in positive slopes that did not contain a null point estimate within the confidence interval.

Discussion

The present meta-analysis explored the dose-response relationship between proximity to failure (quantified as estimated RIR), strength gains, and muscle hypertrophy. Our results showed that strength gains are minimally influenced by proximity to failure, while muscle hypertrophy tends to increase non-linearly as sets are terminated closer to failure. Moreover, the dose-response relationships between estimated proximity to failure and both training outcomes (i.e., muscle hypertrophy and strength gains) seem to be further modified by the load used. These relationships can inform future research and potentially improve the conceptual understanding of practitioners in regards to how proximity to failure influences muscular hypertrophy and strength outcomes, respectively.

Strength

In all of the best-fit models within the present analysis, the confidence intervals of the marginal slope for estimated RIR contained a null point estimate, suggesting a negligible relationship between proximity to failure and strength gains. Additionally, there was an interaction observed between load and RIR in all of the best-fit models, indicating that, as load decreases, training farther from failure results in increasingly more favorable outcomes. There are multiple potential explanations for these findings. In the following sections, we will discuss the principle of specificity in regards to force production and opposing advantages of training far from and close to failure for strength gain to explain the observed relationship.

Force Production and The Principle of Specificity

The main estimates from the statistical models were extracted after adjusting for both load and the method in which volume was equated. Other meta-analytic data suggest that load, in

particular, exhibits a positive dose-response relationship with strength gains [3,4]. These data, in concert with the present findings, suggest that load is a better predictor of strength outcomes than proximity to failure. Importantly, as load increases, the furthest obtainable proximity to failure decreases. For example, when training with a 10RM load (~75% of 1RM) a set could be terminated after a single repetition resulting in 9 RIR. However, a set terminated after a single repetition with a 2RM load (~95% of 1RM) will be much closer to failure (i.e., 1 RIR). This example illustrates the difference between a *load-mediated* and *intra-set-fatigue-mediated* change in proximity to failure. The present analysis would suggest that *intra-set-fatigue-mediated* changes in proximity to failure have a negligible influence on strength gains, while there is more research in support of *load-mediated* changes.

The contrast between load-mediated and intra-set-fatigue mediated changes in proximity to failure and their impact on strength outcomes could be potentially explained by the principle of specificity [99] in regards to force production. To align with the principle of specificity, one would expect that training with similar forces to the strength assessment of interest would result in favorable outcomes such as training with loads >85% of 1RM to maximize performance on a 1RM test. Importantly, however, force production declines as sets are taken closer to failure [100] but increases proportionally with load (i.e., % of 1RM) [101], thereby supporting the contrast between load-mediated and intra-set-fatigue mediated changes in proximity to failure and their impact strength gains. Moreover, a recent meta-analysis by Zhang et al. [20] concluded that lower velocity loss thresholds resulted in higher strength gains per repetition than higher velocity loss thresholds. These results indicate that repetitions performed early in the set, which have the highest force production, lead to the greatest relative strength gain. Additionally, this concept may explain the interaction between RIR and load in most of the present models, in which training farther from failure seemed to be more beneficial for lower loads compared to higher loads. Indeed, lower loads tend to result in greater decrements in force production and total intra-set fatigue (i.e., higher velocity loss) than higher loads when set-equated conditions are performed to failure [102], which could potentially explain the findings of the present meta-analysis.

Advantages of Training Far From Failure for Strength

There are a number opposing advantages of training far from and close to failure for strength gains that may effectively counterbalance one another, leading to a negligible overall relationship. The first potential advantage of conditions training farther from failure is fatigue

management. Multiple acute studies have demonstrated that training closer to failure results in greater indices of neuromuscular and perceptual fatigue compared to training farther from failure [32,102,103]. Since resistance training interventions often do not employ a formal tapering period in which training stress is reduced temporarily to allow for an improvement in acute performance [104], training farther from failure may allow participants to perform strength assessments in the absence of training-related fatigue compared to conditions training closer to failure.

Additionally, there may be proximity-to-failure-specific adaptations to rate of force development (RFD), which may contribute to the attenuation of maximal strength gains. While maximal strength (i.e., absolute force production measured in a specific context) is by definition a time-independent characteristic, the rate at which force is produced could influence an individual's ability to generate sufficient absolute force i) prior to the onset of fatigue, and ii) reaching the position in the range-of-motion with the greatest force requirements. Pareja-Blanco et al. [54] reported that training the Smith-machine squat to a 40% velocity loss threshold resulted in a reduction in the RFD measured from 0-50ms while lower velocity loss thresholds (i.e., 0, 10 and 20%) improved RFD, suggesting training closer to failure may harm RFD. However, differential adaptations seem to occur in the early (<200 ms) and the late (>200 ms) phases of RFD, with the latter being suggested as more predictive of maximal strength [105].

In an additional investigation, Pareja-Blanco et al. [55] reported that only groups training the Smith-machine bench press to higher velocity loss thresholds (i.e., 25 and 50%) saw improvements in RFD measured from 0-400ms, suggesting training closer to failure may harm early but not late phase RFD. However, as previously mentioned, these studies cannot delineate if the proximity to failure or relative volume load (i.e., % of 1RM × repetitions × sets) mediates effects on RFD, as both are manipulated via velocity loss thresholds. In studies that equate relative volume load and measure RFD, conditions training farther from failure usually result in superior changes in RFD, regardless of the phase examined [23,30,31,35,42]. Finally, a recent cross-sectional examination of strength-trained individuals demonstrate significant differences in earlier, but not later phases of RFD compared to untrained controls, potentially indicating that all phases of RFD may contribute to maximal strength [106]; however, this area lacks sufficient evidence to confidently describe the relationship between proximity to failure and phase-specific alterations to RFD.

Advantages of Training Close To Failure for Strength

In contrast to the potential advantages of training farther from failure, training closer to failure may offer exposure to motor patterns and psychological experiences more specific to a maximal strength assessment. It is well established that motor patterns change with increases in load, which require that single repetition sets are terminated closer to failure, in exercises with many degrees of freedom (e.g., barbell back squat) [107]. As maximal strength assessments occur at the threshold of failure - conditions training closer to failure may be more regularly exposed to similar motor demands, thereby aligning with the principle of specificity. Similarly, there seem to be psycho-physiological inputs to maximal strength (e.g., visualization) in which greater exposure to the subjective experience of performing resistance training near or to failure could be beneficial [108]. Practically, training closer to failure also allows for greater accuracy in subjectively reported RIR [109]. Given that loads are often selected via the perception of RIR (i.e., RIR-based RPE) [110], inaccuracy in this perception of RIR could also lead individuals to train with lower loads unnecessarily and potentially harm strength outcomes.

The second advantage of training closer to failure is greater increases in muscle size, as supported by the present meta-analysis. Although the contribution of changes in muscle size to changes in strength has been a topic of debate in the scientific literature [111,112], it is plausible that some relationship exists, though the degree of which is uncertain. Given this assumption, training closer to failure would benefit from greater muscle-size mediated strength gain than conditions training farther from failure.

Taken collectively, these opposing advantages of training far from and close to failure may explain the negligible dose-response relationship between estimated proximity to failure and strength gain in the present meta-analysis.

Hypertrophy

In contrast to results from the analysis of strength outcomes, the confidence interval of the marginal slope for estimated RIR did not contain a null point estimate in all best-fit models for muscle hypertrophy. In other words, the findings suggest a meaningful non-linear relationship

between proximity to failure and changes in muscle size whereby muscle hypertrophy tends to increase as sets are performed closer to failure. Additionally, the findings demonstrate that this non-linear relationship may be mediated by the load used, as marginal slopes increased alongside load (i.e., heavier loads do not require sets to be terminated as close to failure to optimize muscle hypertrophy.) In the following sections, we will discuss the mechanistic support for this relationship, potential explanations for the non-linearity of the observed effects, and other important considerations of the present findings.

Mechanistic Support of the Observed Relationship

The non-linear relationship observed in the present meta-analysis seems to support the proposed mechanistic models of resistance-training-induced muscle hypertrophy [113]. Specifically, Henneman's size principle suggests that motor units, and the muscle fibers they innervate, are recruited sequentially based on their size as force requirements rise [114]. As higher threshold motor units tend to innervate more type II fibers, which may have a greater potential for hypertrophy [115], the goal becomes to create the necessary and sufficient conditions to recruit these motor units allowing the muscle fibers they innervate to experience mechanical tension [116]. Critically, training to failure seems to allow for a stimulus to be delivered to these muscle fibers independent of load [117], while heavier loads may allow for greater motor unit recruitment farther from failure [118]. This rationale seems to be supported by our data as greater changes in muscle size were observed close to failure for all main effects but this pattern was less pronounced as load increased to and beyond ~80% of 1RM.

Explaining the Non-linearity of the Observed Effects

It is unclear why our findings suggest a non-linear increase in changes in muscle size while terminating sets closer to failure. While training to or close to failure is associated with increased acute fatigue [102,103] and performance decrements [119,120], there is a paucity of studies that examine fatigue longitudinally. It could be that the repeated bout effect strongly diminishes the fatigue associated with training to or close to failure as one habituates to the stimulus [121]. Alternatively, ambiguity in failure definitions could be playing a role. Halperin et al. [109] demonstrated that participants often under-predict the number of repetitions they are able to perform, thus, potentially training farther from failure than intended. If underprediction does occur, studies that state participants trained to "failure" without a clear definition of criteria for set termination would result in RIR estimations that were too low (i.e.,

estimated to be too close to failure) in the present meta-analysis. However, after adjusting for load and volume, the marginal slope of RIR from the linear spline models suggests that studies which provided a failure definition resulted in greater changes in muscle size than those that did not. Because both models suggest that gains in muscle size increase non-linearly the closer to failure a set is terminated, ambiguity in failure definition does not seem to influence the directionality of the dose-response relationship. Although, the magnitude of the expected effects may differ between studies providing explicit failure definitions and those that do not. Ultimately, it is difficult to explain the non-linearity of the observed relationship and future research should aim to explore its origins.

Other Findings

Another interesting finding of the present meta-analysis was that the marginal slopes for RIR were not meaningfully influenced by the method in which volume was equated (i.e., set or repetition volume-equated). In all of the recent velocity loss meta-analyses [16–19] the authors report positive linear dose-response relationships between the velocity loss threshold in which sets are terminated and changes in muscle size. However, as previously mentioned, velocity loss thresholds influence both proximity to failure and relative volume load, which makes it difficult to utilize the positive linear dose-response relationships to inform RIR prescriptions for muscle hypertrophy. With the present meta-analysis in mind, it seems that the method in which volume was equated (i.e., set or repetition volume-equated) does not meaningfully alter the dose-response relationship between estimated proximity to failure and changes in muscle size. Thus, a particular type of volume-equated study design (i.e., either set or repetition volume-equated) does not seem to be necessary to inform the relationship between proximity to failure and muscle hypertrophy.

Finally, it is important to mention that the present analysis predominantly focuses on the proximity to failure that optimizes the *average* muscle hypertrophy of the *prime movers* of a given exercise (e.g., a training intervention that features a leg press means that the quadriceps are the prime mover). There are considerably fewer studies that examine the effect of proximity to failure on synergistic muscle groups (e.g., triceps brachii in the bench press), thus the present findings should not be generalized blindly. Moreover, very few studies measure hypertrophy of multiple regions within a given muscle (e.g., proximal, middle, distal) which could exhibit differential relationships to the proximity to failure at which training is performed.

To provide more context for these relationships, future research should aim to measure synergistic muscles and regional hypertrophy of all the sites of interest.

Limitations and Considerations


Several limitations exist with this meta-analysis. While considerable thought and collaboration were put into constructing the highest quality process of estimating RIR, the accuracy of these estimations is unknown. Previous research suggests that the number of repetitions performed at a given load is highly individual [122–124]. Thus, estimating RIR based on homogeneous RM values may be representative of the group-level RIR but likely only applies to some participants within a study. Additionally, multiple other factors that influence proximity to failure were not directly addressed. First, as participants perform multiple sets, set-to-set performance declines due to fatigue. Specifically, if the load is not adjusted from set to set and sets are performed to the same repetition target (e.g., 10 repetitions), there may be fewer RIR on later sets. Therefore, the estimated RIR values, on average, could be overestimated (i.e., the estimations are too far from failure) - although the exact extent to which this may occur is influenced by various factors (e.g., sex, exercise selection, load, rest period, and initially prescribed proximity to failure); thus, this could not be adequately accounted for.

Another factor that may influence RIR is strength gain. Specifically, if the load is not adequately adjusted as participants gain strength (i.e., progressive overload), and sets are performed to the same repetition target (e.g., 10 repetitions), there may be greater RIR in later sessions of the training program. Therefore, the estimated RIR values, on average, could be underestimated (i.e., the estimations are too close to failure) - although, again, the extent to which this may occur is influenced by various factors (e.g., exercise selection, load, weight increments available, and frequency of load progression); thus, this could not be adequately accounted for. Another limitation comes from the decision to average RIR values across all sets performed to describe each group. Therefore, this analysis does not take into account the variability in RIR throughout a training program. For example, at an average estimated RIR of 2, the group could have performed all sets at 2 RIR or an even proportion of 1 and 3 RIR sets.

From a practical perspective, the observed relationships between estimated RIR and strength or muscle hypertrophy outcomes may not hold on the individual level. For example, in the context of training volume, Damas et al. [125] demonstrated that some participants saw more

favorable outcomes in a limb training with <10 sets per week compared to a limb training with >10 sets per week. Crucially, these findings contradict a well-cited meta-analysis by Schoenfeld et al. [1], which suggests that >10 sets per week are optimal for muscle hypertrophy outcomes, on average. This example demonstrates that training recommendations derived from meta-analytic estimates may be inappropriate for some individuals; however, in the absence of robust individual-level evidence (e.g., N of 1 trials) average effects may be the best estimate [126,127]. Moreover, applying these relationships outside the context of the volumes, loads, frequencies, and study designs included in this analysis should be done with caution (see Figure 1 and supplementary file 2). For example, only volume-equated studies (i.e., either set or repetition equated) were included in the present analysis. While a dose-response relationship may exist for muscle hypertrophy in the present analysis, such that training closer to failure results in superior outcomes, these results may change if participants could modify the number of sets performed to align with their recovery capacity. As training to failure results in greater acute fatigue [102,103], training with a greater number of RIR could allow for more weekly sets and could impact longitudinal strength and muscle hypertrophy outcomes. Finally, many studies did not provide the necessary data for the analysis, so much of it had to be estimated (e.g., pre- to post-test correlation coefficients).

Our goal with this analysis was to provide reasonably precise population average RIR estimates that, in the absence of better evidence, can describe the relationship between proximity to failure and resistance training outcomes. While the limitations are notable, this analysis may be useful to explore the directionality of these relationships and identify potential RIR thresholds of interest for future research.



Note: Estimates are adjusted for i) load (% of 1RM), ii) method used to equate training volume (i.e., sets, repetitions, or both), training status, and intervention duration. Dashed lines represent effects as a standardized mean change, while solid lines represent effects as a response ratio.

Maximal Strength	Muscle Hypertrophy
<ul style="list-style-type: none"> ➔ Strength gains increase negligibly as sets are terminated farther from failure. ➔ The slope of RIR is further modified by load, as training with lighter loads results in even greater strength gains farther from failure. ➔ Load-mediated, rather than intraset-fatigue-mediated, decreases in RIR may be preferred to optimize strength gains. 	<ul style="list-style-type: none"> ➔ Changes in muscle size increase non-linearly as sets are terminated closer to failure. ➔ The slope of RIR is further modified by load, as training with heavier loads results in less of an advantage of training closer to failure. ➔ Groups training to failure (with involuntary set termination) saw favorable outcomes compared to those that did not.

Figure 10: Visual summary of the results from the best-fit linear-log meta-regressions for strength and hypertrophy outcomes in the primary analysis.

Conclusion

The dose-response relationships between estimated proximity to failure and strength gain appears to be different from that with muscle hypertrophy (Figure 10). Strength gains seem to be negligibly impacted by the proximity to failure in which sets are performed at a given load, while muscle hypertrophy improves non-linearly as sets are terminated closer to failure. These results seem to be further influenced by the load used (i.e., % of 1RM), as the absolute magnitude of both relationships (i.e., strength and muscle hypertrophy) decreased alongside heavier loads. Researchers and practitioners should be cautious interpreting the findings of the present analysis, as proximity to failure was estimated and its accuracy cannot be confirmed.

Data and Supplementary Material Accessibility

All materials, data, and code are available on the Open Science Framework project page.

Author Contributions

ZR extracted the data, performed the initial RIR estimations, performed the statistical analysis, and drafted the manuscript. JP, JR, and MZ provided the first peer-review of the RIR estimations and edited the manuscript. MR and IJ provided the second peer-review of the RIR estimations and edited the manuscript. JS assisted with the statistical analysis, and edited the manuscript.

Conflict of Interest

Zac Robinson, Joshua Pelland, Jacob Remmert, and, Michael Zourdos are all coaches and writers in the fitness industry. Martin Refalo, Ivan Jukic, and James Steele declare that they have no conflicts of interest relevant to the content of this review.

References

1. Schoenfeld BJ, Ogborn D, Krieger JW. Dose-response relationship between weekly resistance training volume and increases in muscle mass: A systematic review and meta-analysis. *Journal of Sports Sciences* [Internet]. Informa UK Limited; 2016;35:1073–82. Available from: <https://doi.org/10.1080/02640414.2016.1210197>
2. Ralston GW, Kilgore L, Wyatt FB, Baker JS. The effect of weekly set volume on strength gain: A meta-analysis. *Sports Medicine* [Internet]. Springer Science; Business Media LLC; 2017;47:2585–601. Available from: <https://doi.org/10.1007/s40279-017-0762-7>
3. Refalo MC, Hamilton DL, Paval DR, Gallagher IJ, Feros SA, Fyfe JJ. Influence of resistance training load on measures of skeletal muscle hypertrophy and improvements in maximal strength and neuromuscular task performance: A systematic review and meta-analysis. *Journal of Sports Sciences* [Internet]. Informa UK Limited; 2021;39:1723–45. Available from: <https://doi.org/10.1080/02640414.2021.1898094>
4. LOPEZ P, RADAELLI R, TAAFFE DR, NEWTON RU, GALVÃO DA, TRAJANO GS, et al. Resistance training load effects on muscle hypertrophy and strength gain: Systematic review and network meta-analysis. *Medicine & Science in Sports & Exercise* [Internet]. Ovid Technologies (Wolters Kluwer Health); 2020;53:1206–16. Available from: <https://doi.org/10.1249/mss.0000000000002585>
5. Schoenfeld BJ, Grgic J, Krieger J. How many times per week should a muscle be trained to maximize muscle hypertrophy? A systematic review and meta-analysis of studies examining the

- effects of resistance training frequency. *Journal of Sports Sciences* [Internet]. Informa UK Limited; 2018;37:1286–95. Available from: <https://doi.org/10.1080/02640414.2018.1555906>
6. Cuthbert M, Haff GG, Arent SM, Ripley N, McMahon JJ, Evans M, et al. Effects of variations in resistance training frequency on strength development in well-trained populations and implications for in-season athlete training: A systematic review and meta-analysis. *Sports Medicine* [Internet]. Springer Science; Business Media LLC; 2021;51:1967–82. Available from: <https://doi.org/10.1007/s40279-021-01460-7>
7. Pelland JC, Robinson ZP, Remmert JF, Cerminaro RM, Benitez B, John TA, et al. Methods for controlling and reporting resistance training proximity to failure: Current issues and future directions. *Sports Medicine* [Internet]. Springer Science; Business Media LLC; 2022;52:1461–72. Available from: <https://doi.org/10.1007/s40279-022-01667-2>
8. Refalo MC, Helms ER, Hamilton DL, Fyfe JJ. Towards an improved understanding of proximity-to-failure in resistance training and its influence on skeletal muscle hypertrophy, neuromuscular fatigue, muscle damage, and perceived discomfort: A scoping review. *Journal of Sports Sciences* [Internet]. Informa UK Limited; 2022;40:1369–91. Available from: <https://doi.org/10.1080/02640414.2022.2080165>
9. Grgic J, Schoenfeld BJ, Orazem J, Sabol F. Effects of resistance training performed to repetition failure or non-failure on muscular strength and hypertrophy: A systematic review and meta-analysis. *Journal of Sport and Health Science* [Internet]. Elsevier BV; 2022;11:202–11. Available from: <https://doi.org/10.1016/j.jshs.2021.01.007>
10. Vieira AF, Umpierre D, Teodoro JL, Lisboa SC, Baroni BM, Izquierdo M, et al. Effects of resistance training performed to failure or not to failure on muscle strength, hypertrophy, and power output: A systematic review with meta-analysis. *Journal of Strength and Conditioning Research* [Internet]. Ovid Technologies (Wolters Kluwer Health); 2021;35:1165–75. Available from: <https://doi.org/10.1519/jsc.0000000000003936>
11. Jukic I, Hooren BV, Ramos AG, Helms ER, McGuigan MR, Tufano JJ. The effects of set structure manipulation on chronic adaptations to resistance training: A systematic review and meta-analysis. *Sports Medicine* [Internet]. Springer Science; Business Media LLC; 2021;51:1061–86. Available from: <https://doi.org/10.1007/s40279-020-01423-4>
12. Davies TB, Tran DL, Hogan CM, Haff GG, Latella C. Chronic effects of altering resistance training set configurations using cluster sets: A systematic review and meta-analysis. *Sports Medicine* [Internet]. Springer Science; Business Media LLC; 2021;51:707–36. Available from: <https://doi.org/10.1007/s40279-020-01408-3>

13. Jukic I, Helms ER, McGuigan MR, García-Ramos A. Using cluster and rest redistribution set structures as alternatives to resistance training prescription method based on velocity loss thresholds. *PeerJ* [Internet]. *PeerJ*; 2022;10:e13195. Available from: <https://doi.org/10.7717/peerj.13195>
14. Steele J, Fisher J, Giessing J, Gentil P. Clarity in reporting terminology and definitions of set endpoints in resistance training. *Muscle & Nerve* [Internet]. Wiley; 2017;56:368–74. Available from: <https://doi.org/10.1002/mus.25557>
15. Fisher JP, Steele J, Smith D. Intensity of effort and momentary failure in resistance training: Are we asking a binary question for a continuous variable? *Journal of Sport and Health Science* [Internet]. Elsevier BV; 2022;11:644–7. Available from: <https://doi.org/10.1016/j.jshs.2022.03.002>
16. Jukic I, Castilla AP, Ramos AG, Hooren BV, McGuigan MR, Helms ER. The acute and chronic effects of implementing velocity loss thresholds during resistance training: A systematic review, meta-analysis, and critical evaluation of the literature. *Sports Medicine* [Internet]. Springer Science; Business Media LLC; 2022;53:177–214. Available from: <https://doi.org/10.1007/s40279-022-01754-4>
17. Hernández-Belmonte A, Pallarès JG. Effects of velocity loss threshold during resistance training on strength and athletic adaptations: A systematic review with meta-analysis. *Applied Sciences* [Internet]. MDPI AG; 2022;12:4425. Available from: <https://doi.org/10.3390/app12094425>
18. Gantois P, Nakamura FY, Alcazar J, Sousa Fortes L de, Pareja-Blanco F, Souza Fonseca F de. The effects of different intra-set velocity loss thresholds on lower-limb adaptations to resistance training in young adults: A systematic review and meta-analysis. *Center for Open Science*; 2021; Available from: <https://doi.org/10.31236/osf.io/v3tr9>
19. Refalo MC, Helms ER, Trexler EricT, Hamilton DL, Fyfe JJ. Influence of resistance training proximity-to-failure on skeletal muscle hypertrophy: A systematic review with meta-analysis. *Sports Medicine* [Internet]. Springer Science; Business Media LLC; 2022; Available from: <https://doi.org/10.1007/s40279-022-01784-y>
20. Zhang X, Feng S, Li H. The effect of velocity loss on strength development and related training efficiency: A doseresponse metaanalysis. *Healthcare* [Internet]. MDPI AG; 2023;11:337. Available from: <https://doi.org/10.3390/healthcare11030337>
21. Jukic I, Prnjak K, King A, McGuigan MR, Helms ER. Velocity loss is a flawed method for monitoring and prescribing resistance training volume with a free-weight back squat exercise.

European Journal of Applied Physiology [Internet]. Springer Science; Business Media LLC; 2023; Available from: <https://doi.org/10.1007/s00421-023-05155-x>

22. Pareja-Blanco F, Rodríguez-Rosell D, Sánchez-Medina L, Sanchis-Moysi J, Dorado C, Mora-Custodio R, et al. Effects of velocity loss during resistance training on athletic performance, strength gains and muscle adaptations. *Scandinavian Journal of Medicine & Science in Sports* [Internet]. Wiley; 2016;27:724–35. Available from: <https://doi.org/10.1111/sms.12678>

23. Andersen V, Paulsen G, Stien N, Baarholm M, Seynnes O, Saeterbakken AH. Resistance training with different velocity loss thresholds induce similar changes in strength and hypertrophy. *Journal of Strength and Conditioning Research* [Internet]. Ovid Technologies (Wolters Kluwer Health); 2021; Publish Ahead of Print. Available from: <https://doi.org/10.1519/jsc.0000000000004067>

24. Myrholt RB, Solberg P, Pettersen H, Seynnes O, Paulsen G. Effects of low- versus high-velocity-loss thresholds with similar training volume on maximal strength and hypertrophy in highly trained individuals. *International Journal of Sports Physiology and Performance* [Internet]. Human Kinetics; 2023;18:368–77. Available from: <https://doi.org/10.1123/ijssp.2022-0161>

25. Martorelli S, Cadore EL, Izquierdo M, Celes R, Martorelli A, Cleto VA, et al. Strength training with repetitions to failure does not provide additional strength and muscle hypertrophy gains in young women. *European Journal of Translational Myology* [Internet]. PAGEPress Publications; 2017;27. Available from: <https://doi.org/10.4081/ejtm.2017.6339>

26. Goto K, Ishii N, Kizuka T, Takamatsu K. The impact of metabolic stress on hormonal responses and muscular adaptations. *Medicine & Science in Sports & Exercise* [Internet]. Ovid Technologies (Wolters Kluwer Health); 2005;37:955–63. Available from: [https://journals.lww.com/acsm-](https://journals.lww.com/acsm-msse/Fulltext/2005/06000/The_Impact_of_Metabolic_Stress_on_Hormonal.9.aspx)

[msse/Fulltext/2005/06000/The_Impact_of_Metabolic_Stress_on_Hormonal.9.aspx](https://journals.lww.com/acsm-msse/Fulltext/2005/06000/The_Impact_of_Metabolic_Stress_on_Hormonal.9.aspx)

27. Santaniello N, Nóbrega S, Scarpelli M, Alvarez I, Otoboni G, Pintanel L, et al. Effect of resistance training to muscle failure vs non-failure on strength, hypertrophy and muscle architecture in trained individuals. *Biology of Sport* [Internet]. Termedia Sp. z.o.o.; 2020;37:333–41. Available from: <https://doi.org/10.5114/biolSport.2020.96317>

28. Rooney KJ, Herbert RD, Balnave RJ. Fatigue contributes to the strength training stimulus. *Med Sci Sports Exerc*. Ovid Technologies (Wolters Kluwer Health); 1994;26:1160–4.

29. Folland JP, Irish CS, Roberts JC, Tarr JE, Jones DA. Fatigue is not a necessary stimulus for strength gains during resistance training. *British Journal of Sports Medicine* [Internet]. BMJ; 2002;36:370–3. Available from: <https://doi.org/10.1136/bjism.36.5.370>

30. Zaras N, Stasinaki A-N, Spiliopoulou P, Mpampoulis T, Hadjicharalambous M, Terzis G. Effect of inter-repetition rest vs. Traditional strength training on lower body strength, rate of force development, and muscle architecture. *Applied Sciences* [Internet]. MDPI AG; 2020;11:45. Available from: <https://doi.org/10.3390/app11010045>
31. Iglesias-Soler E, Mayo X, Rìo-Rodrìguez D, Carballeira E, Fariñas J, Fernàndez-Del-Olmo M. Inter-repetition rest training and traditional set configuration produce similar strength gains without cortical adaptations. *Journal of Sports Sciences* [Internet]. Informa UK Limited; 2015;34:1473–84. Available from: <https://doi.org/10.1080/02640414.2015.1119299>
32. Lasevicius T, Schoenfeld BJ, Silva-Batista C, Souza Barros T de, Aihara AY, Brendon H, et al. Muscle failure promotes greater muscle hypertrophy in low-load but not in high-load resistance training. *Journal of Strength and Conditioning Research* [Internet]. Ovid Technologies (Wolters Kluwer Health); 2019;36:346–51. Available from: <https://doi.org/10.1519/jsc.0000000000003454>
33. Fariñas J, Mayo X, Giraldez-Garcìa MA, Carballeira E, Fernandez-Del-Olmo M, Rial-Vazquez J, et al. Set configuration in strength training programs modulates the cross education phenomenon. *Journal of Strength and Conditioning Research* [Internet]. Ovid Technologies (Wolters Kluwer Health); 2021;35:2414–20. Available from: <https://doi.org/10.1519/jsc.0000000000003189>
34. Nòbrega SR, Ugrinowitsch C, Pintanel L, Barcelos C, Libardi CA. Effect of resistance training to muscle failure vs. Volitional interruption at high- and low-intensities on muscle mass and strength. *Journal of Strength and Conditioning Research* [Internet]. Ovid Technologies (Wolters Kluwer Health); 2018;32:162–9. Available from: <https://doi.org/10.1519/jsc.0000000000001787>
35. Zaras N, Stasinaki A-N, Mpampoulis T, Spiliopoulou P, Hadjicharalambous M, Terzis G. Effect of inter-repetition rest vs. Traditional resistance training on the upper body strength rate of force development and triceps brachii muscle architecture. *Journal of Human Kinetics* [Internet]. Walter de Gruyter GmbH; 2022;81:189–98. Available from: <https://doi.org/10.2478/hukin-2022-0016>
36. Terada K, Kikuchi N, Burt D, Voisin S, Nakazato K. Low-load resistance training to volitional failure induces muscle hypertrophy similar to volume-matched, velocity fatigue. *Journal of Strength and Conditioning Research* [Internet]. Ovid Technologies (Wolters Kluwer Health); 2020;36:1576–81. Available from: <https://doi.org/10.1519/jsc.0000000000003690>
37. Lacerda LT, Marra-Lopes RO, Diniz RCR, Lima FV, Rodrigues SA, Martins-Costa HC, et al. Is performing repetitions to failure less important than volume for muscle hypertrophy and strength? *Journal of Strength and Conditioning Research* [Internet]. Ovid Technologies (Wolters

- Kluwer Health); 2019;34:1237–48. Available from:
<https://doi.org/10.1519/jsc.0000000000003438>
38. Rial-Vázquez J, Mayo X, Tufano JJ, Fariñas J, Rúa-Alonso M, Iglesias-Soler E. Cluster vs. Traditional training programmes: Changes in the force-velocity relationship. *Sports Biomechanics* [Internet]. Informa UK Limited; 2020;21:85–103. Available from:
<https://doi.org/10.1080/14763141.2020.1718197>
39. Bergamasco JGA, Silva DG da, Bittencourt DF, Oliveira RM de, Júnior JCB, Caruso FR, et al. Low-load resistance training performed to muscle failure or near muscle failure does not promote additional gains on muscle strength, hypertrophy, and functional performance of older adults. *Journal of Strength and Conditioning Research* [Internet]. Ovid Technologies (Wolters Kluwer Health); 2020;36:1209–15. Available from:
<https://doi.org/10.1519/jsc.0000000000003632>
40. J Fariñas, Giraldez-García Ma, E Carballeira, Rial -Vázquez, Mayo X, Fernández-Del-Olmo M, et al. CROSS EDUCATION IN a KNEE EXTENSION EXERCISE DEPENDS ON SET CONFIGURATION OF RESISTANCE TRAINING. Unpublished; 2019; Available from:
<http://rgdoi.net/10.13140/RG.2.2.19941.96485>
41. Karsten B, Fu Y (Leon), Larumbe-Zabala E, Seijo M, Naclerio F. Impact of two high-volume set configuration workouts on resistance training outcomes in recreationally trained men. *Journal of Strength and Conditioning Research* [Internet]. Ovid Technologies (Wolters Kluwer Health); 2021;35:S136–43. Available from: <https://doi.org/10.1519/jsc.0000000000003163>
42. Silva LXN da, Teodoro JL, Menger E, Lopez P, Grazioli R, Farinha J, et al. Repetitions to failure versus not to failure during concurrent training in healthy elderly men: A randomized clinical trial. *Experimental Gerontology* [Internet]. Elsevier BV; 2018;108:18–27. Available from:
<https://doi.org/10.1016/j.exger.2018.03.017>
43. IZQUIERDO-GABARREN M, EXPÓSITO RGDT, GARCÍA-PALLARÈS J, SÀNCHEZ-MEDINA L, VILLARREAL ESSD, IZQUIERDO M. Concurrent endurance and strength training not to failure optimizes performance gains. *Medicine & Science in Sports & Exercise* [Internet]. Ovid Technologies (Wolters Kluwer Health); 2010;42:1191–9. Available from:
<https://doi.org/10.1249/mss.0b013e3181c67eec>
44. Padulo J, Mignogna P, Mignardi S, Tonni F, D'Ottavio S. Effect of different pushing speeds on bench press. *International Journal of Sports Medicine* [Internet]. Georg Thieme Verlag KG; 2012;33:376–80. Available from: <https://doi.org/10.1055/s-0031-1299702>
45. Kilgallon MJ, Johnston MJ, Kilduff LP, Watsford ML. A comparison of training with a velocity loss threshold or to repetition failure on upper-body strength development in professional

- australian footballers. *International Journal of Sports Physiology and Performance* [Internet]. *Human Kinetics*; 2021;16:1758–63. Available from: <https://doi.org/10.1123/ijsp.2020-0882>
46. Held S, Hecksteden A, Meyer T, Donath L. Improved strength and recovery after velocity-based training: A randomized controlled trial. *International Journal of Sports Physiology and Performance* [Internet]. *Human Kinetics*; 2021;16:1185–93. Available from: <https://doi.org/10.1123/ijsp.2020-0451>
47. Fisher JP, Blossom D, Steele J. A comparison of volume-equated knee extensions to failure, or not to failure, upon rating of perceived exertion and strength adaptations. *Applied Physiology, Nutrition, and Metabolism* [Internet]. *Canadian Science Publishing*; 2016;41:168–74. Available from: <https://doi.org/10.1139/apnm-2015-0421>
48. Sampson JA, Groeller H. Is repetition failure critical for the development of muscle hypertrophy and strength? *Scandinavian Journal of Medicine & Science in Sports* [Internet]. *Wiley*; 2015;26:375–83. Available from: <https://doi.org/10.1111/sms.12445>
49. Izquierdo M, Ibañez J, González-Badillo JJ, Häkkinen K, Ratamess NA, Kraemer WJ, et al. Differential effects of strength training leading to failure versus not to failure on hormonal responses, strength, and muscle power gains. *Journal of Applied Physiology* [Internet]. *American Physiological Society*; 2006;100:1647–56. Available from: <https://doi.org/10.1152/jap.01400.2005>
50. Cuevas-Aburto J, Jukic I, González-Hernández JM, Janicijevic D, Barboza-González P, Chirisa-Ríos LJ, et al. Effect of resistance-training programs differing in set configuration on maximal strength and explosive-action performance. *International Journal of Sports Physiology and Performance* [Internet]. *Human Kinetics*; 2021;16:243–9. Available from: <https://doi.org/10.1123/ijsp.2019-1005>
51. LIM C, KIM HJ, MORTON RW, HARRIS R, PHILLIPS SM, JEONG TS, et al. Resistance exercise-induced changes in muscle phenotype are load dependent. *Medicine & Science in Sports & Exercise* [Internet]. *Ovid Technologies (Wolters Kluwer Health)*; 2019;51:2578–85. Available from: <https://doi.org/10.1249/mss.0000000000002088>
52. Edward Z. Pelka RC. COMPARISON OF VELOCITY-BASED & TRADITIONAL RESISTANCE EXERCISE TRAINING.
53. Sánchez-Moreno M, Cornejo-Daza PJ, González-Badillo JJ, Pareja-Blanco F. Effects of velocity loss during body mass prone-grip pull-up training on strength and endurance performance. *Journal of Strength and Conditioning Research* [Internet]. *Ovid Technologies (Wolters Kluwer Health)*; 2020;34:911–7. Available from: <https://doi.org/10.1519/jsc.0000000000003500>

54. PAREJA-BLANCO F, ALCAZAR J, SÀNCHEZ-VALDEPEÑAS J, CORNEJO-DAZA PJ, PIQUERAS-SANCHIZ F, MORA-VELA R, et al. Velocity loss as a critical variable determining the adaptations to strength training. *Medicine & Science in Sports & Exercise* [Internet]. Ovid Technologies (Wolters Kluwer Health); 2020;52:1752–62. Available from: <https://doi.org/10.1249/mss.0000000000002295>
55. Pareja-Blanco F, Alcazar J, Cornejo-Daza PJ, Sánchez-Valdepeñas J, Rodríguez-Lopez C, Mora JH, et al. Effects of velocity loss in the bench press exercise on strength gains, neuromuscular adaptations, and muscle hypertrophy. *Scandinavian Journal of Medicine & Science in Sports* [Internet]. Wiley; 2020;30:2154–66. Available from: <https://doi.org/10.1111/sms.13775>
56. Rodríguez-Rosell D, Yàñez-García JM, Mora-Custodio R, Sánchez-Medina L, Ribas-Serna J, González-Badillo JJ. Effect of velocity loss during squat training on neuromuscular performance. *Scandinavian Journal of Medicine & Science in Sports* [Internet]. Wiley; 2021;31:1621–35. Available from: <https://doi.org/10.1111/sms.13967>
57. Rodiles-Guerrero L, Pareja-Blanco F, León-Prados JA. Effect of velocity loss on strength performance in bench press using a weight stack machine. *International Journal of Sports Medicine* [Internet]. Georg Thieme Verlag KG; 2020;41:921–8. Available from: <https://doi.org/10.1055/a-1179-5849>
58. Sánchez-Moreno M, Rodríguez-Rosell D, Díaz-Cueli D, Pareja-Blanco F, González-Badillo JJ. Effects of velocity loss threshold within resistance training during concurrent training on endurance and strength performance. *International Journal of Sports Physiology and Performance* [Internet]. Human Kinetics; 2021;16:849–57. Available from: <https://doi.org/10.1123/ijsp.2020-0349>
59. Pareja-Blanco F, Sánchez-Medina L, Suárez-Arrones L, González-Badillo JJ. Effects of velocity loss during resistance training on performance in professional soccer players. *International Journal of Sports Physiology and Performance* [Internet]. Human Kinetics; 2017;12:512–9. Available from: <https://doi.org/10.1123/ijsp.2016-0170>
60. Rodríguez-Rosell D, Yàñez-García JM, Mora-Custodio R, Pareja-Blanco F, Ravelo-García AG, Ribas-Serna J, et al. Velocity-based resistance training: Impact of velocity loss in the set on neuromuscular performance and hormonal response. *Applied Physiology, Nutrition, and Metabolism* [Internet]. Canadian Science Publishing; 2020;45:817–28. Available from: <https://doi.org/10.1139/apnm-2019-0829>
61. Rissanen J, Walker S, Pareja-Blanco F, Häkkinen K. Velocity-based resistance training: Do women need greater velocity loss to maximize adaptations? *European Journal of Applied*

- Physiology [Internet]. Springer Science; Business Media LLC; 2022;122:1269–80. Available from: <https://doi.org/10.1007/s00421-022-04925-3>
62. Galiano C, Pareja-Blanco F, Mora JH de, Villarreal ES de. Low-velocity loss induces similar strength gains to moderate-velocity loss during resistance training. *Journal of Strength and Conditioning Research* [Internet]. Ovid Technologies (Wolters Kluwer Health); 2020;36:340–5. Available from: <https://doi.org/10.1519/jsc.0000000000003487>
63. Rodiles-Guerrero L, Cornejo-Daza PJ, Sánchez-Valdepeñas J, Alcazar J, Rodriguez-López C, Sánchez-Moreno M, et al. Specific adaptations to 0%, 15%, 25%, and 50% velocity-loss thresholds during bench press training. *International Journal of Sports Physiology and Performance* [Internet]. Human Kinetics; 2022;17:1231–41. Available from: <https://doi.org/10.1123/ijsp.2021-0481>
64. Davies TB, Halaki M, Orr R, Mitchell L, Helms ER, Clarke J, et al. Effect of set-structure on upper-body muscular hypertrophy and performance in recreationally-trained male and female. *Journal of Strength and Conditioning Research* [Internet]. Ovid Technologies (Wolters Kluwer Health); 2021;36:2176–85. Available from: <https://doi.org/10.1519/jsc.0000000000003971>
65. Nicholson G, Ispoglou T, Bissas A. The impact of repetition mechanics on the adaptations resulting from strength-, hypertrophy- and cluster-type resistance training. *European Journal of Applied Physiology* [Internet]. Springer Science; Business Media LLC; 2016;116:1875–88. Available from: <https://doi.org/10.1007/s00421-016-3439-2>
66. Rodríguez-Rosell D, Yàñez-García JM, Sánchez-Medina L, Mora-Custodio R, González-Badillo JJ. Relationship between velocity loss and repetitions in reserve in the bench press and back squat exercises. *Journal of Strength and Conditioning Research* [Internet]. Ovid Technologies (Wolters Kluwer Health); 2020;34:2537–47. Available from: <https://doi.org/10.1519/jsc.0000000000002881>
67. Chapman M, Larumbe-Zabala E, Goss-Sampson M, Colpus M, Triplett NT, Naclerio F. Perceptual, mechanical, and electromyographic responses to different relative loads in the parallel squat. *Journal of Strength and Conditioning Research* [Internet]. Ovid Technologies (Wolters Kluwer Health); 2019;33:8–16. Available from: <https://doi.org/10.1519/jsc.0000000000001867>
68. Chapman M, Larumbe-Zabala E, Goss-Sampson M, Triplett NT, Naclerio F. Using perceptual and neuromuscular responses to estimate mechanical changes during continuous sets in the bench press. *Journal of Strength and Conditioning Research* [Internet]. Ovid Technologies (Wolters Kluwer Health); 2019;33:2722–32. Available from: <https://doi.org/10.1519/jsc.0000000000002516>

69. Sánchez-Moreno M, Rodríguez-Rosell D, Pareja-Blanco F, Mora-Custodio R, González-Badillo JJ. Movement velocity as indicator of relative intensity and level of effort attained during the set in pull-up exercise. *International Journal of Sports Physiology and Performance* [Internet]. *Human Kinetics*; 2017;12:1378–84. Available from: <https://doi.org/10.1123/ijsp.2016-0791>
70. Vieira JG. Resistance training with repetition to failure or not on muscle strength and perceptual responses. *Journal of Professional Exercise Physiology*. 2019;
71. Zac P, Robinson MCJ, Christian T, Macarilla, Zourdos MC. The effect of resistance training proximity to failure on muscular adaptations and longitudinal fatigue in trained men. Master's Thesis. 2023;
72. Ruple BA, Plotkin DL, Smith MA, Godwin JS, Sexton CL, McIntosh MC, et al. The effects of resistance training to near failure on strength, hypertrophy, and motor unit adaptations in previously trained adults. *Physiological Reports* [Internet]. *Wiley*; 2023;11. Available from: <https://doi.org/10.14814/phy2.15679>
73. Drinkwater EJ, Lawton TW, Lindsell RP, Pyne DB, Hunt PH, McKenna MJ. Training leading to repetition failure enhances bench press strength gains in elite junior athletes. *The Journal of Strength and Conditioning Research* [Internet]. *Ovid Technologies (Wolters Kluwer Health)*; 2005;19:382. Available from: <https://doi.org/10.1519/r-15224.1>
74. Oliver JM, Jagim AR, Sanchez AC, Mardock MA, Kelly KA, Meredith HJ, et al. Greater gains in strength and power with intraset rest intervals in hypertrophic training. *Journal of Strength and Conditioning Research* [Internet]. *Ovid Technologies (Wolters Kluwer Health)*; 2013;27:3116–31. Available from: <https://doi.org/10.1519/jsc.0b013e3182891672>
75. Hansen KT, Cronin JB, Pickering SL, Newton MJ. Does cluster loading enhance lower body power development in preseason preparation of elite rugby union players? *Journal of Strength and Conditioning Research* [Internet]. *Ovid Technologies (Wolters Kluwer Health)*; 2011;25:2118–26. Available from: <https://doi.org/10.1519/jsc.0b013e318220b6a3>
76. Prestes J, Tibana RA, Araujo Sousa E de, Cunha Nascimento D da, Oliveira Rocha P de, Camarço NF, et al. Strength and muscular adaptations after 6 weeks of rest-pause vs. Traditional multiple-sets resistance training in trained subjects. *Journal of Strength and Conditioning Research* [Internet]. *Ovid Technologies (Wolters Kluwer Health)*; 2019;33:S113–21. Available from: <https://doi.org/10.1519/jsc.0000000000001923>
77. Carneiro MAS, Oliveira Júnior GN de, Sousa JFR de, Santagnello SB, Souza MVC, Orsatti FL. Effects of cluster training sets on muscle power and force-velocity relationship in

- postmenopausal women. *Sport Sciences for Health* [Internet]. Springer Science; Business Media LLC; 2019;16:257–65. Available from: <https://doi.org/10.1007/s11332-019-00599-1>
78. Samson A, Pillai P, Somashekharan. Effect of cluster training versus traditional training on muscular strength among recreationally active males—a comparative study. *Indian Journal of Physiotherapy and Occupational Therapy - An International Journal* [Internet]. Diva Enterprises Private Limited; 2018;12:122. Available from: <https://doi.org/10.5958/0973-5674.2018.00022.9>
79. Zarezadehmehrzi M, Aliasghar & Aminai. Effects of traditional and cluster resistance training on explosive power in soccer players. *Indian Journal of Health and Physical Activity*. 2013;
80. Beck M, Varner W, LeVault L, Boring J, Fahs CA. Decline in unintentional lifting velocity is both load and exercise specific. *Journal of Strength and Conditioning Research* [Internet]. Ovid Technologies (Wolters Kluwer Health); 2020;34:2709–14. Available from: <https://doi.org/10.1519/jsc.0000000000003786>
81. Richens B, Cleather D. THE RELATIONSHIP BETWEEN THE NUMBER OF REPETITIONS PERFORMED AT GIVEN INTENSITIES IS DIFFERENT IN ENDURANCE AND STRENGTH TRAINED ATHLETES. *Biology of Sport* [Internet]. Index Copernicus; 2014;31:157–61. Available from: <https://doi.org/10.5604/20831862.1099047>
82. Hernández-Belmonte A, Courel-Ibáñez J, Conesa-Ros E, Martínez-Cava A, Pallarès JG. Level of effort: A reliable and practical alternative to the velocity-based approach for monitoring resistance training. *Journal of Strength and Conditioning Research* [Internet]. Ovid Technologies (Wolters Kluwer Health); 2021;36:2992–9. Available from: <https://doi.org/10.1519/jsc.0000000000004060>
83. Hoeger DR, Werner W. K., Hopkins. Relationship between repetitions and selected percentages of one repetition maximum: a comparison between untrained and trained males and females. *Journal of Strength and Conditioning Research*. Ovid Technologies (Wolters Kluwer Health); 1990;
84. Harris DM, Oranchuk DJ, Latella C. Neuromuscular adaptations to different set configurations during a periodized power training block in elite junior judokas. *International Journal of Sports Science & Coaching* [Internet]. SAGE Publications; 2023;174795412211511. Available from: <https://doi.org/10.1177/17479541221151195>
85. Iglesias-Soler E, Carballeira E, Sánchez-Otero T, Mayo X, Fernández-del-Olmo M. Performance of maximum number of repetitions with cluster-set configuration. *International Journal of Sports Physiology and Performance* [Internet]. Human Kinetics; 2014;9:637–42. Available from: <https://doi.org/10.1123/ijspp.2013-0246>

86. Viechtbauer W. Conducting meta-analyses in R with the metafor package. *Journal of Statistical Software* [Internet]. 2010;36:1–48. Available from: <https://doi.org/10.18637/jss.v036.i03>
87. Amrhein V, Greenland S, McShane B. Scientists rise up against statistical significance. *Nature* [Internet]. Springer Science; Business Media LLC; 2019;567:305–7. Available from: <https://doi.org/10.1038/d41586-019-00857-9>
88. Gardner MJ, Altman DG. Confidence intervals rather than p values: Estimation rather than hypothesis testing. *BMJ* [Internet]. BMJ; 1986;292:746–50. Available from: <https://doi.org/10.1136/bmj.292.6522.746>
89. Noortgate WV den, Lòpez-Lòpez JA, Marìn-Martìnez F, Sànchez-Meca J. Three-level meta-analysis of dependent effect sizes. *Behavior Research Methods* [Internet]. Springer Science; Business Media LLC; 2012;45:576–94. Available from: <https://doi.org/10.3758/s13428-012-0261-6>
90. Hedges LV, Tipton E, Johnson MC. Robust variance estimation in meta-regression with dependent effect size estimates. *Research Synthesis Methods* [Internet]. Wiley; 2010;1:39–65. Available from: <https://doi.org/10.1002/jrsm.5>
91. Morris SB. Estimating effect sizes from pretest-posttest-control group designs. *Organizational Research Methods* [Internet]. SAGE Publications; 2007;11:364–86. Available from: <https://doi.org/10.1177/1094428106291059>
92. Lajeunesse MJ. On the meta-analysis of response ratios for studies with correlated and multi-group designs. *Ecology* [Internet]. Wiley; 2011;92:2049–55. Available from: <https://doi.org/10.1890/11-0423.1>
93. Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, et al., editors. *Cochrane handbook for systematic reviews of interventions* [Internet]. Wiley; 2019. Available from: <https://doi.org/10.1002/9781119536604>
94. Nakagawa S, Santos ESA. Methodological issues and advances in biological meta-analysis. *Evolutionary Ecology* [Internet]. Springer Science; Business Media LLC; 2012;26:1253–74. Available from: <https://doi.org/10.1007/s10682-012-9555-5>
95. Nakagawa S, Schielzeth H. A general and simple method for obtaining R^2 from generalized linear mixed-effects models. OHara RB, editor. *Methods in Ecology and Evolution* [Internet]. Wiley; 2012;4:133–42. Available from: <https://doi.org/10.1111/j.2041-210x.2012.00261.x>
96. Nakagawa S, Lagisz M, O’Dea RE, Rutkowska J, Yang Y, Noble DW, et al. The orchard plot: Cultivating a forest plot for use in ecology, evolution, and beyond. *Research Synthesis Methods*. 2021;12:4–12.

97. Lüdtke D, Ben-Shachar MS, Patil I, Waggoner P, Makowski D. performance: An R package for assessment, comparison and testing of statistical models. *Journal of Open Source Software*. 2021;6:3139.
98. Lenth RV. Emmeans: Estimated marginal means, aka least-squares means [Internet]. 2022. Available from: <https://CRAN.R-project.org/package=emmeans>
99. Zatsiorsky VM, Kraemer WJ. Science and practice of strength training. 2nd ed. Champaign, IL: Human Kinetics; 2006.
100. Piqueras-Sanchiz F, Cornejo-Daza PJ, Sánchez-Valdepeñas J, Bachero-Mena B, Sánchez-Moreno M, Martín-Rodríguez S, et al. Acute mechanical, neuromuscular, and metabolic responses to different set configurations in resistance training. *Journal of Strength and Conditioning Research* [Internet]. Ovid Technologies (Wolters Kluwer Health); 2021;36:2983–91. Available from: <https://doi.org/10.1519/jsc.0000000000004068>
101. McBride JM, Haines TL, Kirby TJ. Effect of loading on peak power of the bar, body, and system during power cleans, squats, and jump squats. *Journal of Sports Sciences* [Internet]. Informa UK Limited; 2011;29:1215–21. Available from: <https://doi.org/10.1080/02640414.2011.587444>
102. Pareja-Blanco F, Rodríguez-Rosell D, Aagaard P, Sánchez-Medina L, Ribas-Serna J, Mora-Custodio R, et al. Time course of recovery from resistance exercise with different set configurations. *Journal of Strength and Conditioning Research* [Internet]. Ovid Technologies (Wolters Kluwer Health); 2020;34:2867–76. Available from: <https://doi.org/10.1519/jsc.0000000000002756>
103. Morán-Navarro R, Pérez CE, Mora-Rodríguez R, Cruz-Sánchez E de la, González-Badillo JJ, Sánchez-Medina L, et al. Time course of recovery following resistance training leading or not to failure. *European Journal of Applied Physiology* [Internet]. Springer Science; Business Media LLC; 2017;117:2387–99. Available from: <https://doi.org/10.1007/s00421-017-3725-7>
104. Travis SK, Mujika I, Gentles JA, Stone MH, Bazylar CD. Tapering and peaking maximal strength for powerlifting performance: A review. *Sports* [Internet]. MDPI AG; 2020;8:125. Available from: <https://doi.org/10.3390/sports8090125>
105. Andersen LL, Andersen JL, Zebis MK, Aagaard P. Early and late rate of force development: Differential adaptive responses to resistance training? *Scandinavian Journal of Medicine & Science in Sports* [Internet]. Wiley; 2010;20:e162–9. Available from: <https://doi.org/10.1111/j.1600-0838.2009.00933.x>
106. Vecchio AD, Negro F, Falla D, Bazzucchi I, Farina D, Felici F. Higher muscle fiber conduction velocity and early rate of torque development in chronically strength-trained individuals.

- Journal of Applied Physiology [Internet]. American Physiological Society; 2018;125:1218–26. Available from: <https://doi.org/10.1152/jappphysiol.00025.2018>
107. Maddox EU, Sievert ZA, Bennett HJ. Modified vector coding analysis of trunk and lower extremity kinematics during maximum and sub-maximum back squats. *Journal of Biomechanics* [Internet]. Elsevier BV; 2020;106:109830. Available from: <https://doi.org/10.1016/j.jbiomech.2020.109830>
108. Paravlic AH, Slimani M, Tod D, Marusic U, Milanovic Z, Pisot R. Effects and dose-response relationships of motor imagery practice on strength development in healthy adult populations: A systematic review and meta-analysis. *Sports Medicine* [Internet]. Springer Science; Business Media LLC; 2018;48:1165–87. Available from: <https://doi.org/10.1007/s40279-018-0874-8>
109. Halperin I, Malleron T, Har-Nir I, Androulakis-Korakakis P, Wolf M, Fisher J, et al. Accuracy in predicting repetitions to task failure in resistance exercise: A scoping review and exploratory meta-analysis. *Sports Medicine* [Internet]. Springer Science; Business Media LLC; 2021;52:377–90. Available from: <https://doi.org/10.1007/s40279-021-01559-x>
110. Helms ER, Cronin J, Storey A, Zourdos MC. Application of the repetitions in reserve-based rating of perceived exertion scale for resistance training. *Strength & Conditioning Journal* [Internet]. Ovid Technologies (Wolters Kluwer Health); 2016;38:42–9. Available from: <https://doi.org/10.1519/ssc.0000000000000218>
111. Loenneke JP, Buckner SL, Dankel SJ, Abe T. Exercise-induced changes in muscle size do not contribute to exercise-induced changes in muscle strength. *Sports Medicine* [Internet]. Springer Science; Business Media LLC; 2019;49:987–91. Available from: <https://doi.org/10.1007/s40279-019-01106-9>
112. Taber CB, Vigotsky A, Nuckols G, Haun CT. Exercise-induced myofibrillar hypertrophy is a contributory cause of gains in muscle strength. *Sports Medicine* [Internet]. Springer Science; Business Media LLC; 2019;49:993–7. Available from: <https://doi.org/10.1007/s40279-019-01107-8>
113. Alix-Fages C, Vecchio AD, Baz-Valle E, Santos-Concejero J, Balsalobre-Fernández C. The role of the neural stimulus in regulating skeletal muscle hypertrophy. *European Journal of Applied Physiology* [Internet]. Springer Science; Business Media LLC; 2022;122:1111–28. Available from: <https://doi.org/10.1007/s00421-022-04906-6>
114. Henneman E, Somjen G, Carpenter DO. EXCITABILITY AND INHIBITIBILITY OF MOTONEURONS OF DIFFERENT SIZES. *Journal of Neurophysiology* [Internet]. American Physiological Society; 1965;28:599–620. Available from: <https://doi.org/10.1152/jn.1965.28.3.599>

115. TESCH PA. Skeletal muscle adaptations consequent to long-term heavy resistance exercise. *Medicine & Science in Sports & Exercise* [Internet]. Ovid Technologies (Wolters Kluwer Health); 1988;20:S132–4. Available from: <https://doi.org/10.1249/00005768-198810001-00008>
116. Wackerhage H, Schoenfeld BJ, Hamilton DL, Lehti M, Hulmi JJ. Stimuli and sensors that initiate skeletal muscle hypertrophy following resistance exercise. *Journal of Applied Physiology* [Internet]. American Physiological Society; 2019;126:30–43. Available from: <https://doi.org/10.1152/jappphysiol.00685.2018>
117. Morton RW, Sonne MW, Zuniga AF, Mohammad IYZ, Jones A, McGlory C, et al. Muscle fibre activation is unaffected by load and repetition duration when resistance exercise is performed to task failure. *The Journal of Physiology* [Internet]. Wiley; 2019;597:4601–13. Available from: <https://doi.org/10.1113/jp278056>
118. Murphy J, Hodson-Tole E, Vigotsky AD, Potvin JR, Fisher JP, Steele J. Motor unit recruitment patterns of the quadriceps differ between continuous high- and low-torque isometric knee extension to momentary failure. *Cold Spring Harbor Laboratory*; 2021; Available from: <https://doi.org/10.1101/2021.04.08.438966>
119. Mangine GT, Serafini PR, Stratton MT, Olmos AA, VanDusseldorp TA, Feito Y. Effect of the repetitions-in-reserve resistance training strategy on bench press performance, perceived effort, and recovery in trained men. *Journal of Strength and Conditioning Research* [Internet]. Ovid Technologies (Wolters Kluwer Health); 2021;36:1–9. Available from: <https://doi.org/10.1519/jsc.0000000000004158>
120. Refalo MC, Helms ER, Hamilton DL, Fyfe JJ. Influence of resistance training proximity-to-failure, determined by repetitions-in-reserve, on neuromuscular fatigue in resistance-trained males and females. *Sports Medicine - Open* [Internet]. Springer Science; Business Media LLC; 2023;9. Available from: <https://doi.org/10.1186/s40798-023-00554-y>
121. Hylldahl RD, Chen TC, Nosaka K. Mechanisms and mediators of the skeletal muscle repeated bout effect. *Exercise and Sport Sciences Reviews* [Internet]. Ovid Technologies (Wolters Kluwer Health); 2017;45:24–33. Available from: <https://doi.org/10.1249/jes.0000000000000095>
122. Cooke DM, Haischer MH, Carzoli JP, Bazylar CD, Johnson TK, Varieur R, et al. Body mass and femur length are inversely related to repetitions performed in the back squat in well-trained lifters. *Journal of Strength and Conditioning Research* [Internet]. Ovid Technologies (Wolters Kluwer Health); 2019;33:890–5. Available from: <https://doi.org/10.1519/jsc.0000000000003021>

123. Zourdos MC, Goldsmith JA, Helms ER, Trepeck C, Halle JL, Mendez KM, et al. Proximity to failure and total repetitions performed in a set influences accuracy of intraset repetitions in reserve-based rating of perceived exertion. *Journal of Strength and Conditioning Research* [Internet]. Ovid Technologies (Wolters Kluwer Health); 2021;35:S158–65. Available from: <https://doi.org/10.1519/jsc.0000000000002995>
124. Mitter B, Zhang L, Bauer P, Baca A, Tschan H. Modelling the relationship between load and repetitions to failure in resistance training: A bayesian analysis. *European Journal of Sport Science* [Internet]. Informa UK Limited; 2022;1–11. Available from: <https://doi.org/10.1080/17461391.2022.2089915>
125. Damas F, Barcelos C, Nóbrega SR, Ugrinowitsch C, Lixandrão ME, Santos LME d., et al. Individual muscle hypertrophy and strength responses to high vs. Low resistance training frequencies. *Journal of Strength and Conditioning Research* [Internet]. Ovid Technologies (Wolters Kluwer Health); 2019;33:897–901. Available from: <https://doi.org/10.1519/jsc.0000000000002864>
126. Steele J, Fisher J, Smith D, Schoenfeld B, Yang Y, Nakagawa S. Meta-analysis of variation in sport and exercise science. *Society for Transparency, Openness,; Replication in Kinesiology*; 2022; Available from: <https://doi.org/10.51224/srxiv.214>
127. Hecksteden A, Kraushaar J, Scharhag-Rosenberger F, Theisen D, Senn S, Meyer T. Individual response to exercise training - a statistical perspective. *Journal of Applied Physiology* [Internet]. American Physiological Society; 2015;118:1450–9. Available from: <https://doi.org/10.1152/jappphysiol.00714.2014>