

**Exercise Training in Metabolic and Bariatric Surgery Adults:
An Overview of Systematic Reviews**

-

SUPPLEMENTARY FILE

Julia Hussien, Marine Asselin, Dale Bond, Yin Wu, Valentina Ly, David Creel, Pavlos
Papasavas, Bret H. Goodpaster, & Aurélie Baillot

Correspondence concerning this manuscript and/or supplementary file should be addressed to:

Aurélie Baillot
School/Department, Université du Québec en Outaouais
283, boulevard Alexandre-Taché (CP 1250, Hull branch)
Gatineau QC, J8X 3X7
Canada
Email: aurelie.baillot@uqo.ca

Hyperlinked Table of Contents

Appendix A – Amendments to Preregistered Protocol	2
Appendix B – PRIOR Checklist.....	3
Appendix C – Search Methods and Strategy	8
Appendix D – Excluded Publications	21
Appendix E – Methodological Details of Included Reviews	23
Appendix F – AMSTAR 2 Ratings.....	35
Appendix G – Corrected Covered Area Calculations Summary	37
Appendix H – Pre-MBS Study Characteristics	38
Appendix I – Pre-MBS Primary Articles.....	43
Appendix J – Post-MBS Study Characteristics.....	46
Appendix K – Post-MBS Primary Articles	56
Appendix L – Post-MBS Subanalyses	63
Appendix M – Feasibility and Acceptability Outcomes	67
Appendix N – Authors Conclusions for Publication Bias by Outcome	69
Appendix O – Certainty of Evidence Summary (RCT-Only Meta-analyses)	75
Appendix P – Outcome Conclusions Flow Diagram.....	80
Appendix Q – Downgrade Rules.....	81
Appendix R – Summary of Outcome Conclusions	82

Appendix A – Amendments to Preregistered Protocol

Table S1: Amendments to Preregistered Protocol with Reasoning

Preregistered Details	Amendment	Reasoning
Selection: Studies screened through RAYYAN	Covidence used instead	Preference of researchers conducting screening
Selection: “To avoid overlap between primary studies, only one systematic review per outcome will be selected. Several systematic reviews could be included for the same outcomes if they provide additional analysis (e.g., by exercise training type or timing, by bariatric surgery). If several systematic reviews are available for one outcome, the highest quality systematic review (AMSTAR 2) will be considered. The most recent systematic review will be chosen if several systematic reviews for the same outcome have the same quality score.”	ALL reviews maintained for all outcomes. Overlap calculated but not avoided	Shifted study aim to synthesize the current FULL body of literature and assess concordance/ discordance between reviews
Risk of bias (quality) assessment	Item 7 was removed from critical item list	Providing a list of excluded original articles with reasons for exclusion is not required by the Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) reporting guidelines
Strategy for data synthesis: “A narrative synthesize will be performed per outcome with tables to ensure consistency of data presentation across studies.”	Added in efforts to determine concordance / discordance followed by a categorization into “what we currently know”, “what we think we know” and “what we still don’t know”	Shifted study aim to synthesize the current FULL body of literature and assess concordance/ discordance between reviews

Note. All amendments were made after protocol preregistration on PROSPERO but before any data collection began on the study. Failing to update the protocol was an oversight.

Appendix B – PRIOR Checklist

(Gates M, Gates A, Pieper D, et al. Reporting guideline for overviews of reviews of healthcare interventions: development of the PRIOR statement. *BMJ* 2022;378:e070849. doi:10.1136/bmj-2022-070849)

Page numbers will be updated based on final typeset article

Section Topic	#	Item	Location reported (Page #)
TITLE			
Title	1	Identify the report as an overview of reviews.	First page
ABSTRACT			
Abstract	2	Provide a comprehensive and accurate summary of the purpose, methods, and results of the overview of reviews.	First page
INTRODUCTION			
Rationale	3	Describe the rationale for conducting the overview of reviews in the context of existing knowledge.	Introduction Page x
Objectives	4	Provide an explicit statement of the objective(s) or question(s) addressed by the overview of reviews.	Introduction Page x
METHODS			
Eligibility criteria	5a	Specify the inclusion and exclusion criteria for the overview of reviews. If supplemental primary studies were included, this should be stated, with a rationale.	Eligibility Criteria Page x & Table 1
	5b	Specify the definition of ‘systematic review’ as used in the inclusion criteria for the overview of reviews.	Table 1
Information sources	6	Specify all databases, registers, websites, organizations, reference lists, and other sources searched or consulted to identify systematic reviews and supplemental primary studies (if included). Specify the date when each source was last searched or consulted.	Information Sources and Search Strategy Page x
Search strategy	7	Present the full search strategies for all databases, registers and websites, such that they could be reproduced. Describe any search filters and limits applied.	*Supplementary File* Appendix C
Selection process	8a	Describe the methods used to decide whether a systematic review or supplemental primary study (if included) met the inclusion criteria of the overview of reviews.	Study Selection Page x
	8b	Describe how overlap in the populations, interventions, comparators, and/or outcomes of systematic reviews was identified and managed during study selection.	Study Selection Page x
Data collection process	9a	Describe the methods used to collect data from reports.	Data Extraction Page x
	9b	If applicable, describe the methods used to identify and manage primary study overlap at the level of the comparison and outcome during data collection. For each outcome, specify the method used to illustrate and/or quantify the degree of primary study overlap across systematic reviews.	Data Synthesis Page x
	9c	If applicable, specify the methods used to manage discrepant data across systematic reviews during data collection.	Data Extraction Page x
Data items	10	List and define all variables and outcomes for which data were sought. Describe any assumptions made and/or measures taken	Data Extraction Page x & Table 2

		to identify and clarify missing or unclear information.	
Risk of bias assessment	11a	Describe the methods used to <u>assess</u> risk of bias or methodological quality of the included systematic reviews.	Risk of Bias of Included Systematic Reviews Page X
	11b	Describe the methods used to <u>collect</u> data on (from the systematic reviews) and/or <u>assess</u> the risk of bias of the primary studies included in the systematic reviews. Provide a justification for instances where flawed, incomplete, or missing assessments are identified but not re-assessed.	Data Extraction Page X & Table 2
	11c	Describe the methods used to <u>assess</u> the risk of bias of supplemental primary studies (if included).	N/A
Synthesis methods	12a	Describe the methods used to summarize or synthesize results and provide a rationale for the choice(s).	Data Synthesis Page X
	12b	Describe any methods used to explore possible causes of heterogeneity among results.	Data Synthesis Page X
	12c	Describe any sensitivity analyses conducted to assess the robustness of the synthesized results.	N/A
Reporting bias assessment	13	Describe the methods used to <u>collect</u> data on (from the systematic reviews) and/or <u>assess</u> the risk of bias due to missing results in a summary or synthesis (arising from reporting biases at the levels of the systematic reviews, primary studies, and supplemental primary studies, if included).	Table 2
Certainty assessment	14	Describe the methods used to <u>collect</u> data on (from the systematic reviews) and/or <u>assess</u> certainty (or confidence) in the body of evidence for an outcome.	Data Extraction Page X & Table 2 Certainty of Evidence of Randomized Controlled Trial-Only Meta-Analyzed Outcomes Page X
RESULTS			
Systematic review and supplemental primary study selection	15a	Describe the results of the search and selection process, including the number of records screened, assessed for eligibility, and included in the overview of reviews, ideally with a flow diagram.	Results Page X & Figure 1
	15b	Provide a list of studies that might appear to meet the inclusion criteria, but were excluded, with the main reason for exclusion.	*Supplementary File* Appendix D
Characteristics of systematic reviews and supplemental primary studies	16	Cite each included systematic review and supplemental primary study (if included) and present its characteristics.	Preoperative Exercise Training Page X Postoperative Exercise Training Page X *Supplementary File* Appendix F, H & J
Primary study overlap	17	Describe the extent of primary study overlap across the included systematic reviews.	Preoperative Exercise Training Page X Postoperative Exercise Training Page X *Supplementary File* Appendix G, I, & K

Risk of bias in systematic reviews, primary studies, and supplemental primary studies	18a	Present assessments of risk of bias or methodological quality for each included systematic review.	Preoperative Exercise Training Page x Postoperative Exercise Training Page x *Supplementary File* Appendix F
	18b	Present assessments (<i>collected</i> from systematic reviews or <i>assessed</i> anew) of the risk of bias of the primary studies included in the systematic reviews.	*Supplementary File* Appendix H & J
	18c	Present assessments of the risk of bias of supplemental primary studies (if included).	N/A
Summary or synthesis of results	19a	For all outcomes, summarize the evidence from the systematic reviews and supplemental primary studies (if included). If meta-analyses were done, present for each the summary estimate and its precision and measures of statistical heterogeneity. If comparing groups, describe the direction of the effect.	<u>Outcomes:</u> Preoperative Exercise Training Page x & Table 3 Postoperative Exercise Training Page x & Table 4 <u>Subanalyses:</u> Subanalyses Page x *Supplementary File* Appendix L <u>Feasibility and Acceptability:</u> Feasibility and Acceptability Page x & Table 5 *Supplementary File* Appendix M <u>Overall:</u> Overarching Results/Conclusions Page x
	19b	If meta-analyses were done, present results of all investigations of possible causes of heterogeneity.	N/A
	19c	If meta-analyses were done, present results of all sensitivity analyses conducted to assess the robustness of synthesized results.	N/A
Reporting biases	20	Present assessments (<i>collected</i> from systematic reviews and/or <i>assessed</i> anew) of the risk of bias due to missing primary studies, analyses, or results in a summary or synthesis (arising from reporting biases at the levels of the systematic reviews, primary studies, and supplemental primary studies, if included) for each summary or synthesis assessed.	*Supplementary File* Appendix N
Certainty of evidence	21	Present assessments (<i>collected</i> or <i>assessed</i> anew) of certainty (or confidence) in the body of evidence for each outcome.	Certainty of Evidence of RCT-Only Meta-Analyzed Outcomes Page x

			<p>Preoperative Exercise Training <i>Table 3</i></p> <p>Postoperative Exercise Training <i>Table 4</i></p>
DISCUSSION			
Discussion	22a	Summarize the main findings, including any discrepancies in findings across the included systematic reviews and supplemental primary studies (if included).	<p>Certainty of Evidence and Study Design Considerations <i>Page x</i></p> <p>Effects of Exercise Training: What We Think We Know <i>Page x</i></p> <p>Beneficial Characteristics of Exercise Training Programs <i>Page x</i></p> <p>Feasibility and Acceptability <i>Page x</i></p> <p>What We Still Don't Know Overall—Implications for Research <i>Page x</i></p>
	22b	Provide a general interpretation of the results in the context of other evidence.	<p>Certainty of Evidence and Study Design Considerations <i>Page x</i></p> <p>Effects of Exercise Training: What We Think We Know <i>Page x</i></p> <p>Beneficial Characteristics of Exercise Training Programs <i>Page x</i></p> <p>Feasibility and Acceptability <i>Page x</i></p> <p>What We Still Don't Know Overall—Implications for Research <i>Page x</i></p>
	22c	Discuss any limitations of the evidence from systematic reviews, their primary studies, and supplemental primary studies (if included) included in the overview of reviews. Discuss any limitations of the overview of reviews methods used.	<p>Strengths And Limitations <i>Page x</i></p>

	22d	Discuss implications for practice, policy, and future research (both systematic reviews and primary research). Consider the relevance of the findings to the end users of the overview of reviews, e.g., healthcare providers, policymakers, patients, among others.	Conclusion Page x
OTHER INFORMATION			
Registration and protocol	23a	Provide registration information for the overview of reviews, including register name and registration number, or state that the overview of reviews was not registered.	Materials And Methods Page x
	23b	Indicate where the overview of reviews protocol can be accessed, or state that a protocol was not prepared.	N/A
	23c	Describe and explain any amendments to information provided at registration or in the protocol. Indicate the stage of the overview of reviews at which amendments were made.	*Supplementary File* Appendix A
Support	24	Describe sources of financial or non-financial support for the overview of reviews, and the role of the funders or sponsors in the overview of reviews.	Funding Information Page x
Competing interests	25	Declare any competing interests of the overview of reviews' authors.	First page
Author information	26a	Provide contact information for the corresponding author.	First page
	26b	Describe the contributions of individual authors and identify the guarantor of the overview of reviews.	Author Contributions Page x
Availability of data and other materials	27	Report which of the following are available, where they can be found, and under which conditions they may be accessed: template data collection forms; data collected from included systematic reviews and supplemental primary studies; analytic code; any other materials used in the overview of reviews.	Reference to Supplementary File materials made where relevant throughout the manuscript

Appendix C – Search Methods and Strategy

1. SR / MA / HTA / ITC - MEDLINE, Embase, PsycInfo. In: CADTH Search Filters Database. Ottawa: CADTH; 2022: <https://searchfilters.cadth.ca/link/33>. Accessed 2022-11-21.
2. SR / MA / HTA / ITC - CINAHL. In: CADTH Search Filters Database. Ottawa: CADTH; 2022: <https://searchfilters.cadth.ca/link/98>. Accessed 2022-11-21.
3. SR / MA / HTA / ITC - Scopus. In: CADTH Search Filters Database. Ottawa: CADTH; 2022: <https://searchfilters.cadth.ca/link/105>. Accessed 2022-11-21.

Table S2: MEDLINE (Ovid) Search Strategy Ovid MEDLINE(R) ALL: 1946 to November 18, 2022		
#	Searches	Results
1	exp bariatric surgery/	32129
2	biliopancreatic diversion/	1092
3	((bariatric* or stomach* or gastric* or gastroileal* or jejuno* or ileo* or intestin* or bilio* or pancreatobiliar*) adj3 (surger* or surgical* or bypass* or diversion* or operat* or procedure*)).ti,ab,kf.	55597
4	((metabolic or weight or obesity or antiobesity or restrictive) adj2 surger*).ti,ab,kf.	6829
5	(stomach* adj2 stapl*).ti,ab,kf.	41
6	(duoden* adj2 switch*).ti,ab,kf.	936
7	((gastric or gastrect* or silicon* or lap or stomach*) adj2 (band* or sleev*)).ti,ab,kf.	11781
8	(gastrojejunostom* or gastroplast* or roux-en-y).ti,ab,kf.	17478
9	or/1-8	74528
10	exp exercise/	238285
11	exp exercise therapy/	61315
12	exp physical fitness/	35572
13	exp exercise movement techniques/	9973
14	bicycling/	12610
15	(exercis* or aerobic* or sport* or walk* or jog* or swim* or danc* or yoga or cycling or biking or bicycl* or bike* or crossfit or tai chi or tai ji or pilate* or plyometric* or fitness or calisthenic* or kinesiotherap* or kinesitherap* or gym* or movement therap*).ti,ab,kf.	832917

16	((physical* or endurance or cardio* or muscl*) adj4 (fit* or train* or activit* or conditon*)).ti,ab,kf.	243264
17	((resistance or strength* or interval* or circuit*) adj4 (train* or program*)).ti,ab,kf.	37410
18	(weight* adj4 (lift* or train*)).ti,ab,kf.	7001
19	(work* adj2 out*).ti,ab,kf.	24311
20	or/10-19	1069291
21	(systematic review or meta-analysis).pt.	291952
22	meta-analysis/ or systematic review/ or systematic reviews as topic/ or meta-analysis as topic/ or "meta analysis (topic)"/ or "systematic review (topic)"/ or exp technology assessment, biomedical/ or network meta-analysis/	328880
23	((systematic* adj3 (review* or overview*)) or (methodologic* adj3 (review* or overview*))).ti,ab,kf.	292532
24	((quantitative adj3 (review* or overview* or synthes*)) or (research adj3 (integrati* or overview*))).ti,ab,kf.	14615
25	((integrative adj3 (review* or overview*)) or (collaborative adj3 (review* or overview*)) or (pool* adj3 analy*)).ti,ab,kf.	36462
26	(data synthes* or data extraction* or data abstraction*).ti,ab,kf.	37511
27	(handsearch* or hand search*).ti,ab,kf.	10798
28	(mantel haenszel or peto or der simonian or dersimonian or fixed effect* or latin square*).ti,ab,kf.	33728
29	(met analy* or metanaly* or technology assessment* or HTA or HTAs or technology overview* or technology appraisal*).ti,ab,kf.	11635
30	(meta regression* or metaregression*).ti,ab,kf.	13406
31	(meta-analy* or metaanaly* or systematic review* or biomedical technology assessment* or bio-medical technology assessment*).mp,hw.	433902
32	(medline or cochrane or pubmed or medlars or embase or cinahl).ti,ab,hw.	315951
33	(cochrane or (health adj2 technology assessment) or evidence report).jw.	21064
34	(comparative adj3 (efficacy or effectiveness)).ti,ab,kf.	16694
35	(outcomes research or relative effectiveness).ti,ab,kf.	10880

36	((indirect or indirect treatment or mixed-treatment or bayesian) adj3 comparison*).ti,ab,kf.	4139
37	(multi* adj3 treatment adj3 comparison*).ti,ab,kf.	286
38	(mixed adj3 treatment adj3 (meta-analy* or metaanaly*)).ti,ab,kf.	176
39	umbrella review*.ti,ab,kf.	1189
40	(multi* adj2 paramet* adj2 evidence adj2 synthesis).ti,ab,kf.	13
41	(multiparamet* adj2 evidence adj2 synthesis).ti,ab,kf.	18
42	(multi-paramet* adj2 evidence adj2 synthesis).ti,ab,kf.	11
43	or/21-42	638533
44	9 and 20 and 43	213

Table S3: Embase (Ovid) Search Strategy
Embase Classic+Embase: 1947 to 2022 November 18

#	Searches	Results
1	exp bariatric surgery/	53897
2	((bariatric* or stomach* or gastric* or gastroileal* or jejuno* or ileo* or intestin* or bilio* or pancreatobiliar*) adj3 (surger* or surgical* or bypass* or diversion* or operat* or procedure*)).ti,ab,kf.	89019
3	((metabolic or weight or obesity or antiobesity or restrictive) adj2 surger*).ti,ab,kf.	11752
4	(stomach* adj2 stapl*).ti,ab,kf.	99
5	(duoden* adj2 switch*).ti,ab,kf.	1885
6	((gastric or gastrect* or silicon* or lap or stomach*) adj2 (band* or sleev*)).ti,ab,kf.	23847
7	(gastrojejunostom* or gastroplast* or roux-en-y).ti,ab,kf.	29914
8	or/1-7	115379
9	exp exercise/	433457
10	exp physical activity/	528169
11	exp kinesiotherapy/	99059
12	fitness/	44353

13	(exercis* or aerobic* or sport* or walk* or jog* or swim* or danc* or yoga or cycling or biking or bicycl* or bike* or crossfit or tai chi or tai ji or pilate* or plyometric* or fitness or calisthenic* or kinesiotherap* or kinesitherap* or gym* or movement therap*).ti,ab,kf.	1092307
14	((physical* or endurance or cardio* or muscl*) adj4 (fit* or train* or activit* or conditon*)).ti,ab,kf.	327108
15	((resistance or strength* or interval* or circuit*) adj4 (train* or program*)).ti,ab,kf.	47367
16	(weight* adj4 (lift* or train*)).ti,ab,kf.	9509
17	(work* adj2 out*).ti,ab,kf.	35972
18	or/9-17	1668824
19	(systematic review or meta-analysis).pt.	0
20	meta-analysis/ or systematic review/ or systematic reviews as topic/ or meta-analysis as topic/ or "meta analysis (topic)"/ or "systematic review (topic)"/ or exp technology assessment, biomedical/ or network meta-analysis/	570330
21	((systematic* adj3 (review* or overview*)) or (methodologic* adj3 (review* or overview*))).ti,ab,kf.	357776
22	((quantitative adj3 (review* or overview* or synthes*)) or (research adj3 (integrati* or overview*))).ti,ab,kf.	17125
23	((integrative adj3 (review* or overview*)) or (collaborative adj3 (review* or overview*)) or (pool* adj3 analy*)).ti,ab,kf.	51553
24	(data synthes* or data extraction* or data abstraction*).ti,ab,kf.	45827
25	(handsearch* or hand search*).ti,ab,kf.	13130
26	(mantel haenszel or peto or der simonian or dersimonian or fixed effect* or latin square*).ti,ab,kf.	44640
27	(met analy* or metanaly* or technology assessment* or HTA or HTAs or technology overview* or technology appraisal*).ti,ab,kf.	18720
28	(meta regression* or metaregression*).ti,ab,kf.	16465
29	(meta-analy* or metaanaly* or systematic review* or biomedical technology assessment* or bio-medical technology assessment*).mp,hw.	679977
30	(medline or cochrane or pubmed or medlars or embase or cinahl).ti,ab,hw.	411013
31	(cochrane or (health adj2 technology assessment) or evidence report).jw.	29476

32	(comparative adj3 (efficacy or effectiveness)).ti,ab,kf.	24750
33	(outcomes research or relative effectiveness).ti,ab,kf.	15951
34	((indirect or indirect treatment or mixed-treatment or bayesian) adj3 comparison*).ti,ab,kf.	7114
35	(multi* adj3 treatment adj3 comparison*).ti,ab,kf.	412
36	(mixed adj3 treatment adj3 (meta-analy* or metaanaly*)).ti,ab,kf.	256
37	umbrella review*.ti,ab,kf.	1251
38	(multi* adj2 paramet* adj2 evidence adj2 synthesis).ti,ab,kf.	27
39	(multiparamet* adj2 evidence adj2 synthesis).ti,ab,kf.	19
40	(multi-paramet* adj2 evidence adj2 synthesis).ti,ab,kf.	22
41	or/19-40	919343
42	8 and 18 and 41	653

Table S4: PsycInfo (Ovid) Search Strategy
APA PsycInfo: 1806 to November Week 2 2022

#	Searches	Results
1	bariatric surgery/	1301
2	((bariatric* or stomach* or gastric* or gastroileal* or jejuno* or ileo* or intestin* or bilio* or pancreatobiliar*) adj3 (surger* or surgical* or bypass* or diversion* or operat* or procedure*)).ti,ab.	1786
3	((metabolic or weight or obesity or antiobesity or restrictive) adj2 surger*).ti,ab.	447
4	(stomach* adj2 stapl*).ti,ab.	0
5	(duoden* adj2 switch*).ti,ab.	8
6	((gastric or gastrect* or silicon* or lap or stomach*) adj2 (band* or sleev*)).ti,ab.	225
7	(gastrojejunostom* or gastroplast* or roux-en-y).ti,ab.	270
8	or/1-7	2075
9	exercise/	26560
10	physical activity/	24885

11	active living/	275
12	physical fitness/	4765
13	walking/	6725
14	running/	2343
15	(exercis* or aerobic* or sport* or walk* or jog* or swim* or danc* or yoga or cycling or biking or bicycl* or bike* or crossfit or tai chi or tai ji or pilate* or plyometric* or fitness or calisthenic* or kinesiotherap* or kinesitherap* or gym* or movement therap*).ti,ab.	179007
16	((physical* or endurance or cardio* or muscl*) adj4 (fit* or train* or activit* or conditon*)).ti,ab.	55242
17	((resistance or strength* or interval* or circuit*) adj4 (train* or program*)).ti,ab.	7163
18	(weight* adj4 (lift* or train*)).ti,ab.	1630
19	(work* adj2 out*).ti,ab.	14933
20	or/9-19	236602
21	(systematic review or meta-analysis).pt.	0
22	meta-analysis/ or systematic review/ or systematic reviews as topic/ or meta-analysis as topic/ or "meta analysis (topic)"/ or "systematic review (topic)"/ or exp technology assessment, biomedical/ or network meta-analysis/	5913
23	((systematic* adj3 (review* or overview*)) or (methodologic* adj3 (review* or overview*))).ti,ab.	50317
24	((quantitative adj3 (review* or overview* or synthes*)) or (research adj3 (integrati* or overview*))).ti,ab.	10782
25	((integrative adj3 (review* or overview*)) or (collaborative adj3 (review* or overview*)) or (pool* adj3 analy*)).ti,ab.	6089
26	(data synthes* or data extraction* or data abstraction*).ti,ab.	3534
27	(handsearch* or hand search*).ti,ab.	1513
28	(mantel haenszel or peto or der simonian or dersimonian or fixed effect* or latin square*).ti,ab.	5890
29	(met analy* or metanaly* or technology assessment* or HTA or HTAs or technology overview* or technology appraisal*).ti,ab.	993
30	(meta regression* or metaregression*).ti,ab.	2419

31	(meta-analy* or metaanaly* or systematic review* or biomedical technology assessment* or bio-medical technology assessment*).mp,hw.	78189
32	(medline or cochrane or pubmed or medlars or embase or cinahl).ti,ab,hw.	33313
33	(cochrane or (health adj2 technology assessment) or evidence report).jx.	0
34	(comparative adj3 (efficacy or effectiveness)).ti,ab.	2301
35	(outcomes research or relative effectiveness).ti,ab.	3848
36	((indirect or indirect treatment or mixed-treatment or bayesian) adj3 comparison*).ti,ab.	494
37	(meta-analysis or systematic review).md.	59169
38	(multi* adj3 treatment adj3 comparison*).ti,ab.	51
39	(mixed adj3 treatment adj3 (meta-analy* or metaanaly*)).ti,ab.	19
40	umbrella review*.ti,ab.	208
41	(multi* adj2 paramet* adj2 evidence adj2 synthesis).ti,ab.	2
42	(multiparamet* adj2 evidence adj2 synthesis).ti,ab.	6
43	(multi-paramet* adj2 evidence adj2 synthesis).ti,ab.	2
44	or/21-43	125987
45	8 and 20 and 44	14

Table S5: Cochrane Database of Systematic Reviews (Ovid) Search Strategy
EBM Reviews - Cochrane Database of Systematic Reviews: 2005 to November 16, 2022

#	Searches	Results
1	((bariatric* or stomach* or gastric* or gastroileal* or jejuno* or ileo* or intestin* or bilio* or pancreatobiliar*) adj3 (surger* or surgical* or bypass* or diversion* or operat* or procedure*)).ti,ab,kw.	34
2	((metabolic or weight or obesity or antiobesity or restrictive) adj2 surger*).ti,ab,kw.	3
3	(stomach* adj2 stapl*).ti,ab,kw.	0
4	(duoden* adj2 switch*).ti,ab,kw.	1
5	((gastric or gastrect* or silicon* or lap or stomach*) adj2 (band* or sleev*)).ti,ab,kw.	5

6	(gastrojejunostom* or gastroplast* or roux-en-y).ti,ab,kw.	6
7	or/1-6	37
8	(exercis* or aerobic* or sport* or walk* or jog* or swim* or danc* or yoga or cycling or biking or bicycling or crossfit or tai chi or tai ji or pilate* or plyometric* or fitness or calisthenic* or kinesiotherap* or kinesitherap* or gym* or movement therap*).ti,ab,kw.	835
9	((physical* or endurance or cardio* or muscl*) adj4 (fit* or train* or activit* or conditon*)).ti,ab,kw.	259
10	((resistance or strength* or interval* or circuit*) adj4 (train* or program*)).ti,ab,kw.	78
11	(weight* adj4 (lift* or train*)).ti,ab,kw.	8
12	(work* adj2 out*).ti,ab,kw.	24
13	or/8-12	929
14	7 and 13	3

Table S6: CINAHL (EBSCOhost) Search Strategy

#	Searches	Results
S20	S8 AND S18 AND S19	63
S19	(MH "meta analysis" OR MH "systematic review" OR MH "Technology, Medical/EV" OR PT "systematic review" OR PT "meta analysis" OR (((TI systematic* OR AB systematic*) N3 ((TI review* OR AB review*) OR (TI overview* OR AB overview*))) OR ((TI methodologic* OR AB methodologic*) N3 ((TI review* OR AB review*) OR (TI overview* OR AB overview*))) OR (((TI quantitative OR AB quantitative) N3 ((TI review* OR AB review*) OR (TI overview* OR AB overview*) OR (TI synthes* OR AB synthes*))) OR ((TI research OR AB research) N3 ((TI integrati* OR AB integrati*) OR (TI overview* OR AB overview*))) OR (((TI integrative OR AB integrative) N3 ((TI review* OR AB review*) OR (TI overview* OR AB overview*))) OR ((TI collaborative OR AB collaborative) N3 ((TI review* OR AB review*) OR (TI overview* OR AB overview*))) OR ((TI pool* OR AB pool*) N3 (TI analy* OR AB analy*)) OR ((TI "data synthes*" OR AB "data synthes*") OR (TI "data extraction*" OR AB "data extraction*") OR (TI "data abstraction*" OR AB "data abstraction*")) OR ((TI handsearch* OR AB handsearch*) OR (TI "hand search*" OR AB "hand search*")) OR ((TI "mantel haenszel" OR AB "mantel haenszel") OR (TI peto OR AB peto) OR (TI "der simonian" OR AB "der simonian") OR (TI dersimonian OR AB dersimonian) OR (TI "fixed effect*" OR AB "fixed effect*") OR (TI "latin square*" OR AB "latin square*")) OR ((TI "met analy*" OR AB "met analy*"))	288,604

	OR (TI metanaly* OR AB metanaly*) OR (TI "technology assessment*" OR AB "technology assessment*") OR (TI HTA OR AB HTA) OR (TI HTAs OR AB HTAs) OR (TI "technology overview*" OR AB "technology overview*") OR (TI "technology appraisal*" OR AB "technology appraisal*")) OR ((TI "meta regression*" OR AB "meta regression*") OR (TI metaregression* OR AB metaregression*)) OR (MW meta-analy* OR MW metaanaly* OR MW "systematic review*" OR MW "biomedical technology assessment*" OR MW "bio-medical technology assessment*") OR ((TI medline OR AB medline OR MW medline) OR (TI cochrane OR AB cochrane OR MW cochrane) OR (TI pubmed OR AB pubmed OR MW pubmed) OR (TI medlars OR AB medlars OR MW medlars) OR (TI embase OR AB embase OR MW embase) OR (TI cinahl OR AB cinahl OR MW cinahl)) OR (SO Cochrane OR SO health technology assessment OR SO evidence report) OR ((TI comparative OR AB comparative) N3 ((TI efficacy OR AB efficacy) OR (TI effectiveness OR AB effectiveness))) OR ((TI "outcomes research" OR AB "outcomes research") OR (TI "relative effectiveness" OR AB "relative effectiveness")) OR (((TI indirect OR AB indirect) OR (TI "indirect treatment" OR AB "indirect treatment") OR (TI mixed-treatment OR AB mixed-treatment) OR (TI bayesian OR AB bayesian)) N3 (TI comparison* OR AB comparison*)) OR ((TI multi* OR AB multi*) N3 (TI treatment OR AB treatment) N3 (TI comparison* OR AB comparison*)) OR ((TI mixed OR AB mixed) N3 (TI treatment OR AB treatment) N3 ((TI meta-analy* OR AB meta-analy*) OR (TI metaanaly* OR AB metaanaly*))) OR (TI "umbrella review*" OR AB "umbrella review*") OR ((TI multi* OR AB multi*) N2 (TI paramet* OR AB paramet*) N2 (TI evidence OR AB evidence) N2 (TI synthesis OR AB synthesis)) OR ((TI multiparamet* OR AB multiparamet*) N2 (TI evidence OR AB evidence) N2 (TI synthesis OR AB synthesis)) OR ((TI multi-paramet* OR AB multi-paramet*) N2 (TI evidence OR AB evidence) N2 (TI synthesis OR AB synthesis))	
S18	S9 OR S10 OR S11 OR S12 OR S13 OR S14 OR S15 OR S16 OR S17	427,744
S17	TI (work* N2 out*) OR AB (work* N2 out*)	13,311
S16	TI (weight* N4 (lift* or train*)) OR AB (weight* N4 (lift* or train*))	3,305
S15	TI ((resistance or strength* or interval* or circuit*) N4 (train* or program*)) OR AB ((resistance or strength* or interval* or circuit*) N4 (train* or program*))	19,704
S14	TI ((physical* or endurance or cardio* or muscl*) N4 (fit* or train* or activit* or conditon*)) OR AB ((physical* or endurance or cardio* or muscl*) N4 (fit* or train* or activit* or conditon*))	107,433
S13	TI ((exercis* or aerobic* or sport* or walk* or jog* or swim* or danc* or yoga or cycling or biking or bicycling or crossfit or tai chi or tai ji or pilate* or plyometric* or fitness or calisthenic* or kinesiotherap* or kinesitherap* or gym* or movement therap*)) OR AB ((exercis* or aerobic* or sport* or walk* or jog* or swim* or danc* or yoga or cycling or biking or bicycling or crossfit or tai chi or tai ji or pilate* or plyometric* or fitness or calisthenic* or kinesiotherap* or kinesitherap* or gym* or movement therap*))	281,360

S12	(MH "Physical Activity")	49,322
S11	(MH "Therapeutic Exercise+")	62,339
S10	(MH "Physical Fitness+")	20,623
S9	(MH "Exercise+")	128,907
S8	S1 OR S2 OR S3 OR S4 OR S5 OR S6 OR S7	17,459
S7	TI ((gastrojejunostom* or gastroplast* or roux-en-y)) OR AB ((gastrojejunostom* or gastroplast* or roux-en-y))	3,225
S6	TI ((gastric or gastrect* or silicon* or lap or stomach*) N2 (band* or sleeve*)) OR AB ((gastric or gastrect* or silicon* or lap or stomach*) N2 (band* or sleeve*))	2,613
S5	TI (duoden* N2 switch*) OR AB (duoden* N2 switch*)	144
S4	TI (stomach* N2 stapl*) OR AB (stomach* N2 stapl*)	19
S3	TI ((metabolic or weight or obesity or antiobesity or restrictive) N2 surger*) OR AB ((metabolic or weight or obesity or antiobesity or restrictive) N2 surger*)	2,736
S2	TI ((bariatric* or stomach* or gastric* or gastroileal* or jejuno* or ileo* or intestin* or bilio* or pancreatobiliar*) N3 (surger* or surgical* or bypass* or diversion* or operat* or procedure*)) OR AB ((bariatric* or stomach* or gastric* or gastroileal* or jejuno* or ileo* or intestin* or bilio* or pancreatobiliar*) N3 (surger* or surgical* or bypass* or diversion* or operat* or procedure*))	12,124
S1	(MH "Bariatric Surgery+")	9,611

Table S7: SPORTDiscus (EBSCOhost) Search Strategy

#	Searches	Results
S9	S7 AND S8	43
S8	PT "systematic review" OR PT "meta analysis" OR (((TI systematic* OR AB systematic*) N3 ((TI review* OR AB review*) OR (TI overview* OR AB overview*))) OR ((TI methodologic* OR AB methodologic*) N3 ((TI review* OR AB review*) OR (TI overview* OR AB overview*))) OR (((TI quantitative OR AB quantitative) N3 ((TI review* OR AB review*) OR (TI overview* OR AB overview*) OR (TI syntheses* OR AB syntheses*))) OR ((TI research OR AB research) N3 ((TI integrati* OR AB integrati*) OR (TI overview* OR AB overview*))) OR (((TI integrative OR AB integrative) N3 ((TI review* OR AB review*) OR (TI overview* OR AB overview*))) OR ((TI collaborative OR AB collaborative) N3 ((TI review* OR AB review*) OR (TI overview* OR AB overview*))) OR ((TI pool* OR AB pool*) N3 (TI analy* OR AB analy*)) OR ((TI "data syntheses*" OR AB "data syntheses*") OR (TI "data extraction*" OR AB	23,657

	<p>"data extraction*" OR (TI "data abstraction*" OR AB "data abstraction*") OR ((TI handsearch* OR AB handsearch*) OR (TI "hand search*" OR AB "hand search*")) OR ((TI "mantel haenszel" OR AB "mantel haenszel") OR (TI peto OR AB peto) OR (TI "der simonian" OR AB "der simonian") OR (TI dersimonian OR AB dersimonian) OR (TI "fixed effect*" OR AB "fixed effect*") OR (TI "latin square*" OR AB "latin square*")) OR ((TI "met analy*" OR AB "met analy*") OR (TI metanaly* OR AB metanaly*) OR (TI "technology assessment*" OR AB "technology assessment*") OR (TI HTA OR AB HTA) OR (TI HTAs OR AB HTAs) OR (TI "technology overview*" OR AB "technology overview*") OR (TI "technology appraisal*" OR AB "technology appraisal*")) OR ((TI "meta regression*" OR AB "meta regression*") OR (TI metaregression* OR AB metaregression*)) OR (MW meta-analy* OR MW metaanaly* OR MW "systematic review*" OR MW "biomedical technology assessment*" OR MW "bio-medical technology assessment*") OR ((TI medline OR AB medline OR MW medline) OR (TI cochrane OR AB cochrane OR MW cochrane) OR (TI pubmed OR AB pubmed OR MW pubmed) OR (TI medlars OR AB medlars OR MW medlars) OR (TI embase OR AB embase OR MW embase) OR (TI cinahl OR AB cinahl OR MW cinahl)) OR (SO Cochrane OR SO health technology assessment OR SO evidence report) OR ((TI comparative OR AB comparative) N3 ((TI efficacy OR AB efficacy) OR (TI effectiveness OR AB effectiveness))) OR ((TI "outcomes research" OR AB "outcomes research") OR (TI "relative effectiveness" OR AB "relative effectiveness")) OR (((TI indirect OR AB indirect) OR (TI "indirect treatment" OR AB "indirect treatment") OR (TI mixed-treatment OR AB mixed-treatment) OR (TI bayesian OR AB bayesian)) N3 (TI comparison* OR AB comparison*)) OR ((TI multi* OR AB multi*) N3 (TI treatment OR AB treatment) N3 (TI comparison* OR AB comparison*)) OR ((TI mixed OR AB mixed) N3 (TI treatment OR AB treatment) N3 ((TI meta-analy* OR AB meta-analy*) OR (TI metaanaly* OR AB metaanaly*))) OR (TI "umbrella review*" OR AB "umbrella review*") OR ((TI multi* OR AB multi*) N2 (TI paramet* OR AB paramet*) N2 (TI evidence OR AB evidence) N2 (TI synthesis OR AB synthesis)) OR ((TI multiparamet* OR AB multiparamet*) N2 (TI evidence OR AB evidence) N2 (TI synthesis OR AB synthesis)) OR ((TI multi-paramet* OR AB multi-paramet*) N2 (TI evidence OR AB evidence) N2 (TI synthesis OR AB synthesis))</p>	
S7	S1 OR S2 OR S3 OR S4 OR S5 OR S6	992
S6	TI ((gastrojejunostom* or gastroplast* or roux-en-y)) OR AB ((gastrojejunostom* or gastroplast* or roux-en-y))	172
S5	TI ((gastric or gastrect* or silicon* or lap or stomach*) N2 (band* or sleeve*)) OR AB ((gastric or gastrect* or silicon* or lap or stomach*) N2 (band* or sleeve*))	135
S4	TI (duoden* N2 switch*) OR AB (duoden* N2 switch*)	7
S3	TI (stomach* N2 stapl*) OR AB (stomach* N2 stapl*)	3
S2	TI ((metabolic or weight or obesity or antiobesity or restrictive) N2 surger*) OR AB ((metabolic or weight or obesity or antiobesity or restrictive) N2 surger*)	326

S1	TI ((bariatric* or stomach* or gastric* or gastroileal* or jejuno* or ileo* or intestin* or bilio* or pancreatobiliar*) N3 (surger* or surgical* or bypass* or diversion* or operat* or procedure*)) OR AB ((bariatric* or stomach* or gastric* or gastroileal* or jejuno* or ileo* or intestin* or bilio* or pancreatobiliar*) N3 (surger* or surgical* or bypass* or diversion* or operat* or procedure*))	778
----	--	-----

Table S8: Scopus Search Strategy

#	Searches	Results
1	((TITLE-ABS-KEY ((bariatric* OR stomach* OR gastric* OR gastroileal* OR jejuno* OR ileo* OR intestin* OR bilio* OR pancreatobiliar*) W/3 (surger* OR surgical* OR bypass* OR diversion* OR operat* OR procedure*)) OR (TITLE-ABS-KEY ((metabolic OR weight OR obesity OR antiobesity OR restrictive) W/2 surger*)) OR (TITLE-ABS-KEY (stomach* W/2 stapl*)) OR (TITLE-ABS-KEY (duoden* W/2 switch*)) OR (TITLE-ABS-KEY ((gastric OR gastrect* OR silicon* OR lap OR stomach*) W/2 (band* OR sleeve*))) OR (TITLE-ABS-KEY (gastrojejunistom* OR gastroplast* OR roux-en-y))) AND ((TITLE-ABS-KEY (exercis* OR aerobic* OR sport* OR walk* OR jog* OR swim* OR danc* OR yoga OR cycling OR biking OR bicycl* OR bike* OR crossfit OR "tai chi" OR "tai ji" OR pilate* OR plyometric* OR fitness OR calisthenic* OR kinesiotherap* OR kinesitherap* OR gym* OR "movement therap*")) OR (TITLE-ABS-KEY ((physical* OR endurance OR cardio* OR muscl*) W/4 (fit* OR train* OR activit* OR conditon*))) OR (TITLE-ABS-KEY ((resistance OR strength* OR interval* OR circuit*) W/4 (train* OR program*))) OR (TITLE-ABS-KEY (weight* W/4 (lift* OR train*))) OR (TITLE-ABS-KEY (work* W/2 out*))) AND (TITLE-ABS-KEY ((systematic* W/3 (review* OR overview*)) OR (methodologic* W/3 (review* OR overview*))) OR TITLE-ABS-KEY ((quantitative W/3 (review* OR overview* OR synthes*)) OR (research W/3 (integrati* OR overview*))) OR TITLE-ABS-KEY ((integrative W/3 (review* OR overview*)) OR (collaborative W/3 (review* OR overview*)) OR (pool* W/3 analy*)) OR TITLE-ABS-KEY ("data synthes*" OR "data extraction*" OR "data abstraction*")) OR TITLE-ABS-KEY (handsearch* OR "hand search*") OR TITLE-ABS-KEY ("mantel haenszel" OR peto OR "der simonian" OR dersimonian OR "fixed effect*" OR "latin square*")) OR TITLE-ABS-KEY ("met analy*" OR metanaly* OR "technology assessment*" OR hta OR htas OR "technology overview*" OR "technology appraisal*") OR TITLE-ABS-KEY ("meta regression*" OR metaregression*) OR TITLE-ABS-KEY (meta-analy* OR metaanaly* OR "systematic review*" OR "biomedical technology assessment*" OR "bio-medical technology assessment*")) OR TITLE-ABS-KEY (medline OR cochrane OR pubmed OR medlars OR embase OR cinahl) OR SRCTITLE (cochrane OR (health W/2 "technology assessment") OR "evidence report")) OR TITLE-ABS-KEY (comparative W/3 (efficacy OR	635

	effectiveness)) OR TITLE-ABS-KEY ("outcomes research" OR "relative effectiveness") OR TITLE-ABS-KEY ((indirect OR "indirect treatment" OR mixed-treatment OR bayesian) W/3 comparison*) OR TITLE-ABS-KEY (multi* W/3 treatment W/3 comparison*) OR TITLE-ABS-KEY (mixed W/3 treatment W/3 (meta-analy* OR metaanaly*)) OR TITLE-ABS-KEY ("umbrella review*") OR TITLE-ABS-KEY (multi* W/2 paramet* W/2 evidence W/2 synthesis) OR TITLE-ABS-KEY (multiparamet* W/2 evidence W/2 synthesis) OR TITLE-ABS-KEY (multi-paramet* W/2 evidence W/2 synthesis))	
--	---	--

Appendix D – Excluded Publications

Table S9: Publications Excluded from the Overview of Reviews

Author(s), Year	Title	Reason for Exclusion
de Aquino Chamis et al., 2022	Abordagem fisioterapêutica no pós-operatório de cirurgia bariátrica: revisão sistemática	Wrong language (i.e., not available in English or French)
Cobos-Fernández et al., 2021	Effects of exercise in bariatric surgery	
Chaves-Alves et al., 2020	Efeito do treinamento de força sobre a aptidão física em pós - bariátricos: revisão sistemática	
DosSantos & Palmeira, 2022	Long-term association between physical activity, weight gain, metabolic risk factors and quality of life in patients undergoing bariatric surgery - systematic literature review	
Zhao et al., 2023	Dietary and physical activity behavior promotion strategies after bariatric surgery from a cognitive perspective: an integrative review	
Amarodos Santos et al., 2022	Associação no longo-termo entre a prática de atividade física, o reganho de peso, fatores de risco metabólico e qualidade de vida, em pacientes submetidos a cirurgia bariátrica - revisão sistemática da literatura	
Barrientos-Sanchez et al., 2022	Physical exercise and loss of weight and body mass index in bariatric surgery: a systematic review	Abstract poster
Cobos-Fernández et al., 2021	Efectos del ejercicio en cirugía bariátrica	
Pouwels et al., 2016	Aspects of exercise before or after bariatric surgery: a systematic review	
Barreto et al., 2015	Physical activity and bariatric surgery - a review	
Pouwels et al., 2017	Comparative analysis of respiratory muscle strength before and after bariatric surgery using five predictive equations exercise and bariatric surgery	
Marshall et al., 2019	Do intensive preoperative and postoperative behavioural interventions impact on health-related bariatric surgery outcomes? A systematic review	
Pattyn et al., 2018	The effects of exercise training on body composition and exercise capacity following bariatric surgery: a systematic review and meta-analysis	Abstract oral presentation
Pouwels et al., 2015	Aspects of exercise before or after bariatric surgery: a systematic review	
Marshall et al., 2019	Do intensive preoperative and postoperative multidisciplinary interventions impact health-related bariatric surgery outcomes? a systematic review	
Egberts et al., 2012	Does exercise improve weight loss after bariatric surgery? A systematic review	No intervention
Livhits et al., 2010	Exercise following bariatric surgery: systematic review	
Smith et al., 2022	Preoperative assessment and prehabilitation in patients with obesity undergoing non-bariatric surgery: a systematic review	

Stewart & Avenell, 2016	Behavioural interventions for severe obesity before and/or after bariatric surgery: a systematic review and meta-analysis	Wrong intervention (i.e., not exercise training)
James et al., 2022	A systematic review of interventions to increase physical activity and reduce sedentary behaviour following bariatric surgery	
Swierz et al., 2020	Systematic review and meta-analysis of perioperative behavioral lifestyle and nutritional interventions in bariatric surgery: a call for better research and reporting	
Jiménez-Loaisa et al., 2015	Psychosocial effects of surgery and physical activity in bariatric patients: a systematic review	
Robinson et al., 2021	Digital technology to support lifestyle and health behaviour changes in surgical patients: systematic review	
Roman et al., 2019	Meta-analysis of the influence of lifestyle changes for preoperative weight loss on surgical outcomes	
Tabesh et al., 2023	Nutrition, physical activity, and prescription of supplements in pre- and post-bariatric surgery patients: an updated comprehensive practical guideline	Wrong study design (i.e., not a systematic review or meta-analysis)
Cornejo-Pareja et al., 2021	Factors related to weight loss maintenance in the medium-long term after bariatric surgery: a review	
Petering & Webb, 2009	Exercise, fluid, and nutrition recommendations for the postgastric bypass exerciser	
Mao et al., 2023	Prehabilitation in metabolic and bariatric surgery: a narrative review	

Appendix E – Methodological Details of Included Reviews

Table S10: Methodological Details of Pre- and Post-Metabolic and Bariatric Surgery (MBS) Systematic Reviews/Meta-Analyses (Alphabetical Order)

Author (Year) Design i.e., SR and/or MA	Aim/Objective	Selection Criteria				Search Information 1. Date (+ Date Restrictions If Applicable) 2. Databases Searched + Other Search Methods
		Patient and/or Problem	Intervention and Comparator	Outcome(s)	Study Design	
Baillot (2014) SR	To appraise current knowledge on the impact of physical activity and physical fitness on the health of adults living with class II and III obesity and those awaiting or who have undergone MBS	Adult population of subjects living with class II and III obesity or subjects awaiting or having undergone MBS	Intervention or recommendations on physical activity or with an evaluation of physical fitness or physical activity level	At least one of these outcomes: anthropometric parameters, body composition, cardiometabolic risk factors, physical fitness, quality of life or psychological parameters.	Peer reviewed original studies	November 16 th , 2012
		When more than one publication used the same cohort, we included only the results from the publication with the largest sample size, unless specific findings (e.g. on selected outcomes) were present only in the other papers.	Comparator requirements not reported	Studies evaluating only weight loss after MBS were excluded since this was previously reviewed.		Medline Scopus CINAHL Sportdiscus + reference lists of included articles + consultation with physical activity and obesity experts
Baillot (2022) MA	To assess the evidence, and identifying factors, of the feasibility and acceptability	Adults awaiting or having undergone MBS.	Articles were excluded if they only presented a study focused	Number of participants who discontinued the intervention and	Primary research involving a controlled trial,	April 2021

	of exercise intervention in adults awaiting or having undergone MBS		on behavioral interventions to promote exercise engagement Comparing an exercise group to a control group without exercise.	reasons; participants' satisfaction ratings/scores; reported attendance rate related to session frequency; reported compliance rate related to session duration and intensity; number and type of adverse events related to exercise intervention only and other quantitative or qualitative data showing feasibility and acceptability of the intervention.	either randomized or non randomized published in a peer-reviewed journal in English	Searched articles in a systematic review and meta-analysis published in June 2021 on exercise training and MBS. +Search for articles published from January 1, 2019, onwards on PubMed, Web of Science, and EMBASE. +hand-search reference lists from eligible articles and relevant reviews.
Bellicha (2021) MA	To examine the impact of physical activity interventions performed before or after MBS in subjects living with obesity	Adults undergoing MBS with at least one obesity comorbidity	Exercise training based on aerobic and/or resistance and/or high-intensity interval training Comparison group of patients undergoing MBS receiving usual care without following an exercise training	Preintervention to postintervention changes reported for at least one of the following outcome category: anthropometry or body composition, objectively measured physical activity or physical	Originals, RCT, NRCT	October 2019 PubMed, Web of Science, and EMBASE + Reference lists from the resulting reviews and

			program	fitness, health-related quality of life, and other relevant health outcomes.		articles were also screened to identify additional articles
Bond (2023) MA	To determine the effect of exercise interventions on weight change 12 months following MBS	Adults who had undergone metabolic and MBS	Exercise intervention Control group that included at least usual post-surgery care	Measured weight change beyond 12 months post surgery	RCT	February 2022 PubMed, Cochrane, Scopus, SPORTDiscus, and CINAHL + Cross referenced lists of included studies, relevant reviews, and meta-analyses manually
Boppre (2021) MA	To determine if exercise favors weight loss and promotes additional benefits on body composition compared to those elicited solely by MBS To determine the characteristics of exercise interventions (mode, duration, and onset after MBS) that were more likely to favor weight loss and body composition benefits	Adults living with severe obesity that underwent MBS	Supervised and semi-supervised training protocols, with a minimum of 1-month duration in addition to the usual medical follow-up, were included and no restrictions were applied on exercise mode, intensity, and the timing onset after surgery	Body weight, anthropometric measures, and body composition (BMI, waist circumference, fat mass and lean body mass) Only studies in which body composition was assessed by dual-energy X-ray absorptiometry were included in fat mass and lean body mass analysis	RCTs published in English	Between 2000 and November 2020. PubMed/MEDLINE®, EBSCO®, Web of Science® and Scopus® + Manual inspection of select articles references

			Control group that received usual follow-up care			
Boppre (2022) MA	To determine the effects of different exercise types, duration, and onset after MBS on cardiometabolic risk factors	Adults living with severe obesity who underwent MBS	Exercise intervention program with a minimum of 1-month duration. No restrictions were applied regarding exercise type, intensity, and onset timing after surgery Post-MBS patients receiving usual medical care only as control group	Cardiometabolic risk factors: primary (lipid profile and glucose metabolism) and secondary outcomes (resting heart rate, systolic and diastolic blood pressure)	RCTs published in English	July 2021 PubMed, Web of Science, Scopus, and EBSCO + A reference inspection from selected articles
Carretero-Ruiz (2019) MA	To analyze the effects of exercise training, and type of training, after MBS in relation to weight loss	Adults having undergone MBS	Exercise interventions performed after MBS Excluded articles combining physical activity with other types of intervention, such as medications, nutrition, other surgeries and lifestyle interventions Comparator requirements not reported	Weight loss	RCT, NRCT	May 23, 2019 MEDLINE, EMBASE, Scopus, Cochrane, and Web of Science
Carretero-Ruiz (2021) MA	To review the evidence on the effectiveness of exercise training to improve cardiometabolic risk in	Adults having undergone MBS	Intervention based on physical activity	At least one outcome related to metabolic risk (VO ₂ max or peak, heart rate, blood	RCT, NRCT	December 6, 2020 MEDLINE,

	patients living with obesity who are undergoing MBS.		Excluded articles combining physical activity with other types of intervention, such as medications, nutrition, other surgeries and lifestyle interventions	pressure, lipid profile, glucose, insulin or inflammation markers)		EMBASE, Scopus, Cochrane, and Web of Science
			Comparator requirements not reported			
Da Silva (2019) MA	To perform a meta-analysis in order to determine the effect size of exercise training on VO ₂ max in adults following MBS weight loss	Adults who have undergone MBS	Intervention including aerobic exercise, resistance exercise, or both Comparator requirements not reported	Association between cardiometabolic risk factors and measured cardiorespiratory variables following MBS by cardiopulmonary exercise testing (VO ₂ max)	Original studies including all eligible prospective cohort models that (1) is written in English language, (2) investigates the association between cardiometabolic risk factors and measured cardio-respiratory variables following MBS and (3) included a description of the exercise training protocol	August 21, 2018 MEDLINE (through PubMed)
Diniz-Sousa (2022) MA	To compare the effect of exercise and usual medical care in the prevention of bone	Adults living with severe obesity submitted to MBS	Supervised or unsupervised structured exercise	Areal bone mineral density from relevant clinical skeletal sites	RCTs and NRCTs	January 2021

	<p>mineral density loss following MBS</p> <p>To identify which skeletal sites might be more responsive to exercise following MBS</p>		<p>training program with a minimum of 3-month duration with any training characteristics (except swimming); including those combined with other interventions (e.g., nutrition)</p> <p>Patients receiving usual medical post-MBS care as control</p>	<p>(total hip, femoral neck, lumbar spine, and one-third distal radius) assessed by dual-energy X-ray absorptiometry</p>		<p>PubMed/MEDLINE, Web of Science, Scopus, and EBSCO</p>
<p>Durey (2022)</p> <p>MA</p>	<p>To summarise the evidence comparing the effects of preoperative whole-body exercise-based interventions on fitness and clinical outcomes for MBS patients</p>	<p>Adult MBS patients</p>	<p>Preoperative exercise intervention</p> <p>Excluded any studies without a control group (didn't define control group)</p>	<p>The primary outcomes assessed were (1) all-cause mortality in the short-term (30 days) and/or longer-term (maximal follow-up), (2) post-operative short-term morbidity, (3) overall quality of life and (4) serious adverse events (short term and longer term).</p> <p>The secondary outcomes were treatment-associated costs, length of hospital stay, number of days of lost work (maximal follow-up), changes in fitness (preoperative and maximal follow-up),</p>	<p>RCT</p>	<p>First search on May 1 2020 and updated on March 2 2021</p> <p>MEDLINE, Embase, Cochrane Central Register of Controlled Clinical Trials, SPORTDiscus, Web of Science and Scopus</p>

				re-operation/re-intervention and its classification of severity, change in weight, diabetes status, technical complications of the specific operation and micronutrient status		
Fonseca-Junior (2013) SR	To investigate the effects of exercise training programs in clinical and surgical treatment of patients living with morbid obesity	Adults living with morbid obesity awaiting or having undergone MBS	Intervention program of physical exercise in the treatment for weight loss or any other aspect of health Could use control group or not	Weight loss or any other aspect of health in general	Pre-and post-intervention test design published in English, Spanish or Portuguese	From January 2000 until July 2012 Virtual Health Library in the database of the “Health Sciences in General” (Medline, Lilacs and IBECS) and PubMed
Gasmi (2022) MA	To evaluate the influence of physical activity on different biological markers of patients’ post-MBS	Adult patients who had undergone MBS	Physical activity intervention Control was MBS	Weight loss and specified biomarkers (BMI, fat mass, fat-free mass, hip to waist ratio, and waist circumference).	RCT published in English	From January 2000 to December 2020 PubMed, Embase, OVID, CINAHL, and Cochrane Library + search of reference list of included articles

Herrera-Santelices (2022) MA	To determine the effect of rehabilitation on the body composition, functional capacity, quality of life and surgical outcomes in patients who are candidates for MBS	Adults awaiting their first MBS who were included in a rehabilitation program	Supervised physical exercise programs described as aerobic exercise training, resistance exercise training or included both, with a duration of at least one week and performed before MBS. The control received only the standard care	One or more of the following outcomes: (1) Body composition, (2) Functional capacity (e.g., six-minute walk test, VO2 max, sit to stand etc.), (3) Quality of life, (4) Surgical outcomes (e.g., number of hospital stay days)	RCT	Between July 1-31 2021 PubMed, Web of Science, SciELO, Scopus, MEDLINE and CINAHL
Jabbour (2022) SR	To review the available evidence for the beneficial health impact of adding exercise to standard care preoperatively and to address metabolic health and surgical outcomes compared to standard care alone in MBS patients	MBS candidates	Preoperative physical activity and/or exercise intervention Standard care control	Fitness level, body weight and composition, physical activity level, physical functioning and muscular performance, aerobic fitness, metabolic parameters, hospital stay	Randomized control trial, intervention trial, and prospective studies	Search date not specified (published before article publication in July 2022) PubMed, Institute for Scientific Information Web of Knowledge, Web of Science, and SPORTDiscus + manual search of references
Karaaslan (2020) SR	To determine the optimal exercise program to prevent weight gain and maintain	Adult MBS patients	Exercise therapy program included after MBS with a study	Changes in body composition, muscular strengths, aerobic capacity, functional	RCT	Articles published from January 2008 through

	weight loss in patients after undergoing MBS.		length of at least 4 weeks	capacity, walking distance, or relevant health outcomes		September 2018.
			Comparator requirements not reported			PubMed (NLM), Pedro, and the Web of Science
Lodewijks (2022) SR	To systematically review the current literature on the overall effects of a preoperative programme concerning exercise, behaviour and/or diet in patients eligible for MBS with a primary focus on weight loss	Preoperative adults who were eligible for MBS and who participated in a preconditioning programme	Preconditioning programme concerning exercise, behaviour and/or diet Comparator requirements not reported	The primary outcome of interest was weight loss. Secondary outcomes were all other available outcomes of the included studies	RCT, NRCT, and pilot studies published in English	Studies published between January 2010 and September 2021 Embase, Cinahl, PubMed and Cochrane Library
Marshall (2020) SR and MA	To evaluate the effect of preoperative and/or post-operative support for adults who elect MBS delivered by an multidisciplinary team on postoperative body composition, mental health, co-morbidities, quality of life, and side effects	Adults who elect MBS delivered by a multidisciplinary team	Intervention had to be implemented by a multidisciplinary team (≥ 3 health disciplines including the surgeon and nurse). Intervention duration needed to be ≥ 2 weeks if delivered preoperatively and ≥ 3 months if delivered post-operatively. Post-operative interventions that commenced >12 months post-operatively were excluded.	Postoperative body composition, mental health, co-morbidities, quality of life and side effects	RCTs, pseudo-RCTs, or non-RCTs.	July 19 2018 Medline (PubMed), CENTRAL, EMBASE, CINAHL, PsycINFO, and Web of Science + snowball search of Google Scholar and key papers

			Prospectively compared a preoperative and/or post-operative intervention delivered by an multidisciplinary team against a comparator group that had less engagement with the multidisciplinary team or had no follow-up			
Morales-Marroquin (2020) SR	To evaluate the effect of resistance training on body composition and strength post-metabolic and MBS	Adults who've undergone MBS	Exercise interventions conducted post-metabolic and MBS that include resistance exercise as part of the exercise intervention (either alone or in combination with cardiovascular exercise)	Body composition (changes in fat-free mass, fat mass, visceral adipose tissue, bone mineral density) and strength	RCT, NRCT written in English	December 2019 PUBMED, Web of Science, and Science Direct
			Comparator requirements not reported			
Pouwels (2015) SR	To determine the kind/type/mode of exercise a patient living with obesity should be advised to undertake What is the most beneficial timing of exercise delivery -	Patients who have undergone or are awaiting MBS	Intervention that includes strength training and/or endurance training or a combination of both. Also, multimodal programs with	Improvement of anthropometric and physical fitness variables (VO ₂ max and/or heart rate reserve / heart rate kinetics), complications, effect	Randomized controlled trial or prospective trial.	July 2014 Pubmed, Embase, and CINAHL

	pre- or postoperatively or a combination of both?		exercise components were included	on weight, and quality of life.		+ cross-references were screened
			Intervention of interest was exercise training compared to regular care			
Ren (2018) MA	To determine whether engaging in exercise after surgery can provide additional weight loss and improvement in physical function.	Adults living with obesity who had undergone MBS	Intervention includes aerobic exercise, resistance training or a combination of both. Control group received only standard care	Primary outcomes: weight loss and physical function (6 min walk test). Secondary outcomes: body mass index, total body fat percentage, fat and fat-free mass, waist and hip circumference, systolic and diastolic blood pressure, and heart rate	RCT	May 2018 PubMed, Embase, the Cochrane Library, OVID and the CINAHL + searched the reference lists of the retrieved articles and relevant review articles
Roth (2022) MA	To investigate the effect of exercise training, protein, calcium, and vitamin D supplementation on the preservation of fat free mass during non-surgical and surgical weight loss and of the combination of all interventions together in adults living with obesity	Overweight or obese adults undergoing diet- or surgery-induced weight loss	Exercise training, or being physically active, alone or combined with dietary supplementation Compared to placebo intervention, controlled comparison intervention or standard care	Fat free mass, bone mineral density and muscle mass	RCT	August 27 2020 Ovid Medline, Ovid Embase, Cochrane Central Register of Controlled Trials, ISI Web of Science
Schurmans (2022) SR	To summarize the effects of different physical exercise programs on various health	Adult	Physical exercise intervention	At least one of the following parameters as an outcome: BMI,	RCTs and quasi-experimental	Search date not specified but

	variables in patients living with obesity in peri-operative MBS	subjects with a body mass index >30 and scheduled to undergo or already have undergone MBS	Compared to usual care	weight loss, muscle strength, lean body or fat free mass, cardiorespiratory endurance, quality of life, or functional capacity	Studies published in English, Dutch, French or German	article sent on February 2021 PubMed and the Cochrane Library
Vieira (2022) MA	To assess the effect of exercise on muscle strength in individuals following MBS To conduct separate meta-analyses for studies that used different muscle strength tests	Adults who underwent MBS	Physical exercise intervention Utilized a control group	Muscle strength (using any method)	RCT and NRCT	October 27, 2021 Embase, Medline, Scopus, SPORTDiscus, and Web of Science + searched 2 grey literature databases (ProQuest and Google Scholar) + reference lists of the included studies were also searched

Note. MA=meta-analysis, SR=systematic review, RCT=randomized controlled trial, NRCT=non randomized controlled trial, MBS=metabolic and bariatric surgery, BMI=body mass index.

Appendix F – AMSTAR 2 Ratings

Figure S1: Breakdown of AMSTAR2 Ratings for the Included Reviews

Reviews	AMSTAR 2 Items																Overall
	1 PICO search	2 a priori review methods	3 Justified design inclusion	4 Comprehensive search strategy	5 Duplicate study selection	6 Duplicate data extraction	7 Justified exclusions	8 Described included studies	9 Risk of bias assessed	10 Funding sources reported	11 Meta-analysis statistics	12 Meta-analysis risk of bias	13 Risk of bias impact	14 Result heterogeneity	15 Publication bias assessed	16 Conflicts of interest declared	
Preoperative Intervention Reviews																	
Durey et al 2022	⊕	⊕	⊖	⊖	⊕	⊕	⊖	⊖	⊕	⊖	⊖	⊖	⊖	⊖	⊖	⊕	C-LOW
Herrera-Santelices et al 2022	⊕	⊕	⊖	⊖	⊕	⊕	⊖	?	⊕	⊖	⊖	⊖	⊕	⊕	⊖	⊕	C-LOW
Jabbour et al 2022	⊕	⊖	⊖	?	⊖	⊖	⊖	?	⊖	⊖	N/A	N/A	⊖	⊕	N/A	⊕	C-LOW
Lodewijks et al 2022	⊖	⊖	⊖	?	⊖	⊖	⊖	?	⊕	⊖	N/A	N/A	⊕	⊕	N/A	⊕	LOW
Postoperative Intervention Reviews																	
Bond et al 2023	⊖	⊕	⊖	?	⊕	⊕	⊖	⊕	⊕	⊖	⊖	⊖	⊕	⊕	⊕	⊕	LOW
Roth et al 2022	⊕	⊕	⊖	?	⊖	⊖	⊖	?	⊕	⊖	⊕	⊖	⊕	⊕	⊕	⊕	MODERATE
Vieira et al 2022	⊖	?	⊖	?	⊕	⊕	⊖	?	⊕	⊖	⊖	⊖	⊕	⊕	⊕	⊕	LOW
Gasmi et al 2022	⊕	⊖	⊖	?	⊖	⊕	⊖	?	⊕	⊖	⊖	⊖	⊖	⊖	⊕	⊕	C-LOW
Diniz-Sousa et al 2022	⊕	⊕	⊖	?	⊕	⊖	⊖	?	⊕	⊖	⊕	⊖	⊕	⊕	⊕	⊕	MODERATE
Boppre et al 2022	⊕	?	⊖	⊖	⊖	⊖	⊖	⊖	⊕	⊖	⊖	⊖	⊖	⊕	⊕	⊕	C-LOW
Boppre et al 2021	⊕	?	⊖	⊖	⊖	⊖	⊖	⊖	⊕	⊖	⊖	⊖	⊕	⊕	⊕	⊕	C-LOW
Carretero-Ruiz et al 2021	⊖	⊖	⊖	?	⊕	⊕	⊖	?	⊕	⊖	⊕	⊖	⊖	⊕	⊕	⊕	C-LOW

Carretero-Ruiz et al 2019	⊖	⊕	⊖	?	⊕	⊕	⊖	?	⊕	⊖	⊕	⊖	⊖	⊕	⊕	⊕	LOW
da Silva et al 2019	⊖	⊖	⊖	⊖	⊕	⊖	⊖	?	⊕	⊖	⊖	⊖	⊖	⊕	⊕	⊕	C-LOW
Ren et al 2018	⊕	⊖	⊖	?	⊕	⊕	⊖	⊖	⊕	⊖	⊕	⊖	⊕	⊕	⊕	⊕	LOW
Morales-Marroquin et al 2020	⊖	⊖	⊖	?	⊖	⊖	⊖	⊕	⊕	⊖	N/A	N/A	⊕	⊕	N/A	⊕	LOW
Civi Karaaslan et al 2020	⊖	⊖	⊖	?	⊖	⊖	⊖	⊖	⊕	⊖	N/A	N/A	⊖	⊖	N/A	⊕	C-LOW
Pre- and Postoperative Intervention Reviews																	
Baillot et al 2022	⊕	⊕	⊖	?	⊕	⊖	⊖	⊕	⊕	⊖	⊖	⊕	⊕	⊕	⊕	⊕	LOW
Marshall et al 2020	⊕	⊕	⊖	?	⊕	⊖	⊖	?	⊕	⊖	⊕	⊕	⊕	⊕	⊖	⊕	LOW
Schurmans et al 2022	⊕	?	⊖	?	⊕	⊖	⊖	⊕	⊕	⊖	N/A	N/A	⊖	⊕	N/A	⊕	LOW
Bellicha et al 2021	⊕	?	⊖	?	⊖	⊖	⊖	⊕	⊕	⊖	⊖	⊕	⊖	⊕	⊕	⊕	C-LOW
Pouwels et al 2015	⊕	⊖	⊖	?	⊕	⊖	⊖	⊖	⊕	⊖	N/A	N/A	⊖	⊖	N/A	⊕	C-LOW
Baillot et al 2014	⊖	⊖	⊖	?	⊕	⊖	⊖	⊖	⊕	⊖	N/A	N/A	⊕	⊕	N/A	⊕	LOW
Fonseca-Junior et al 2013	⊖	⊖	⊖	?	⊖	⊖	⊖	⊖	⊖	⊖	N/A	N/A	⊖	⊖	N/A	⊕	C-LOW

Note. Critical categories highlighted in grey. Green positive symbol=full completion of criteria, yellow question mark=partial completion of criteria, and red negative symbol=lack of completion of criteria. C-LOW= critically low. 1=PICO criteria used for search, 2=explicit a priori established review methods, 3=justification of study design inclusion, 4=comprehensive literature search strategy, 5=study selection completed in duplicate, 6=data extraction completed in duplicate, 7=list and justification for excluded individual studies, 8=adequate description of included studies, 9=satisfactory technique for risk of bias (RoB) assessment, 10=sources of funding of included studies listed, 11=appropriate statistics to combine results for meta-analysis, 12=impact of RoB on meta-analysis evidence synthesis, 13=RoB considered for result interpretation/discussion, 14=explanation and discussion of result heterogeneity, 15=publication bias assessed and interpreted for quantitative synthesis and 16=declared author conflicts of interest.

Appendix G – Corrected Covered Area Calculations Summary

Table S11: Summary of the Corrected Covered Area (CCA) Calculations

	Number of Times Studies Appeared in Reviews	Number of Studies	Number of Reviews	CCA
Pre-MBS				
Overall	65	23	11	0.18
Body weight, Body mass index, Weight Loss	30	15	6	0.20
Fat Mass	8	4	4	0.33
Fat Free and Lean Body Mass	6	3	4	0.33
VO ₂ max	10	8	3	0.13
6-Min Walk Test Distance	10	6	4	0.22
Muscle strength	6	4	2	0.50
Resting Heart Rate	4	3	2	0.33
Blood Pressure	11	6	4	0.28
Quality of Life	11	7	4	0.19
Glucose/Lipid Metabolism	3	3	2	0.00
Physical Activity	6	5	2	0.20
Adverse Events	1	1	1	NA
Hospital Stay Length	2	2	2	0.00
Post-MBS				
Overall	196	42	20	0.19
Body weight, Body mass index, Weight Loss	78	29	8	0.24
Waist Circumference*	6	5	2	0.20
Fat Mass*	19	11	4	0.24
Fat Free and Lean Body Mass*	35	16	6	0.24
Bone Mineral Density	11	4	4	0.58
VO ₂ max	32	12	5	0.42
6-Min Walk Test Distance	10	6	3	0.33
Muscle Strength	29	16	4	0.27
Resting Heart Rate*	17	9	4	0.30
Blood Pressure*	30	13	5	0.33
Quality of Life	4	3	2	0.33
Glucose Metabolism	19	10	4	0.30
Lipid Metabolism	25	12	5	0.27

Note. *This outcome was included in Ren et al., 2018; however, Ren et al., 2018 was not included in the CCA calculation since the studies been reviewed were not disclosed. MBS=metabolic and MBS.

Appendix H – Pre-MBS Study Characteristics

Table S12: Summary of Meta-Analyses and Systematic Reviews for Pre-Metabolic and Bariatric Surgery Exercise Interventions

Author Year (Country ^a)	Study Designs	Sample Description (N, Age Range, BMI Range, % Women) ^b	Intervention Description	Control Group	Outcomes Assessed ^c	Main Results ^d Level of Evidence	Quality Scores of the Studies Included (<i>Quality Assessment Tool</i>)	AMSTAR Rating
Meta-Analyses								
Baillot 2022 (Canada+1) <i>k=7</i>	RCT (<i>k=4</i>) NRCT (<i>k=3</i>)	NO SYNTHESIS ONLY FOR PRE-MBS	Exercise training (<i>k=7</i>) <u>Duration:</u> 4-26 weeks <u>Type:</u> E, E/R, HIIT, aquatic <u>Frequency:</u> 2-5x/wk. <u>Intensity:</u> Moderate to vigorous <u>Session duration:</u> 25-70 min <u>Supervision:</u> supervised (<i>k=6</i>), partial (<i>k=1</i>)	Usual care (<i>k=7</i>)	1. Adverse events during exercise 2. Attendance 3. Dropout 4. Compliance 5. Satisfaction	1. None (<i>k=2</i>), occasional pain, fatigue, or dyspnea (<i>k=1</i>), hypoglycemia or hypotension (<i>k=1</i>) 2. Pooled percentage: 79.4% [67.7; 89.4], I ² =0%; <i>n</i> =NR, <i>k=4</i> 3. Pooled percentage: 3.6% [0.0; 14.6], I ² =44%; <i>n</i> =NR, <i>k=7</i> 4. No-metanalysis due to small number of studies (<i>k=3</i>) and various operationalization of compliance 5. all participants very satisfied/satisfied, except 1 participant = moderately satisfied by location and schedule (<i>k=1</i>) <u>Sub-analysis:</u> NS differences in attendance, dropout rates based on studies quality, exercise intervention timing, and exercise intervention duration	Good (<i>k=1</i>) Fair (<i>k=3</i>) Poor (<i>k=3</i>) (<i>National Heart, Lung and Blood Institute quality assessment tool</i>)	Low
Durey 2022 (UK+1) <i>k=5</i>	RCT (<i>k=5</i>) included 2 conference abstracts	<i>N</i> = 199 38-47 years 45 to 47 kg.m ² 68-86% women	Exercise training (<i>k=3</i>) PA counseling (<i>k=2</i>) <u>Duration:</u> 2-12 wk. <u>Type:</u> E, R, E/R, HIIT, BC <u>Frequency:</u> 3-7x/week <u>Intensity:</u> Moderate to vigorous	Usual care (<i>k=5</i>)	1. Adverse events 2. Pre-MBS VO ₂ max change 3. VO ₂ max change at	1. RR: 6.00 adverse events [0.27; 131.34], I ² =N/A; <i>n</i> =22; <i>k=1</i> 2. MD: 0.73 mL/kg/min [0.61; 0.86], I ² =62%; <i>n</i> =79, <i>k=3</i> 3. MD: 0.98 mL/kg/min [0.05; 1.90], I ² =0%; <i>n</i> =131, <i>k=3</i> 4. MD: 0.94% [-1.61; 3.48], I ² =70%; <i>n</i> =142; <i>k=3</i>	Some concerns (<i>k=2</i>) High risk (<i>k=3</i>) (<i>Cochrane risk of bias tool 2</i>)	Critically low

			<u>Session duration:</u> 120-240 min/week <u>Supervision:</u> unsupervised (<i>k</i> =2), partial (<i>k</i> =2), full-supervised (<i>k</i> =1)		maximal follow-up 4. %WL 5. Length of hospital stay	5. NS ≠ bw intervention and control; <i>n</i> =22; <i>k</i> =1		
Herrera-Santelices 2022 (Chili+1) <i>k</i> =5 including 2 publications with the same intervention and overlapping data in the results	RCT (<i>k</i> =5)	<i>N</i> = 139* 28-54 years BMI = NR 76-95% women * <i>N</i> =114 (<i>k</i> =4), however Baillot 2016/2018 included in same analyses	Exercise training (<i>k</i> =5) <u>Duration:</u> 8-16 wk. <u>Type:</u> E, R, E/R, Water aerobics <u>Frequency:</u> 2-3x/wk. <u>Intensity:</u> Moderate to vigorous <u>Session duration:</u> 25-80 min <u>Supervision:</u> supervised (<i>k</i> =5)	Usual care (<i>k</i> =4) No intervention (<i>k</i> =1)	1. BMI 2. %FM 3. FFM 4. 6MWTD 5. QoL	1. SMD: -0.71 [-1.55; 0.12], <i>I</i> ² =76%; <i>n</i> =115; <i>k</i> =4, very low 2. SMD: 0.38 [-0.08; 0.84], <i>I</i> ² =0%; <i>n</i> =75; <i>k</i> =3, moderate 3. SMD: - 0.41[-1.00; 0.18], <i>I</i> ² =0%; <i>n</i> =46; <i>k</i> =2, moderate 4. SMD: 2.59 [1.89; 3.30], <i>I</i> ² =0%; <i>n</i> =61; <i>k</i> =2, high 5. SMD: 0.88 [-0.23; 1.99], <i>I</i> ² =67%; <i>n</i> =53; <i>k</i> =3, moderate	High risk (<i>k</i> =1) Uncertain risk (<i>k</i> =4) (Cochrane risk of bias tool)	Critically low
Marshall (2020) (Australia) <i>k</i> = 3 (data resulting in 8 publications)	RCT (<i>k</i> =3)	<i>N</i> =208 43-47 years 45.8-47.4 kg.m ² 80-90% women	Exercise training (<i>k</i> =1) PA counselling (<i>k</i> =2) <u>Duration:</u> 6 weeks <u>Type:</u> E/R, aqua fitness <u>Frequency:</u> 3 x/wk. <u>Intensity:</u> NR <u>Session duration:</u> 30-80 min <u>Supervision:</u> Supervised (<i>k</i> =1), NR (<i>k</i> = 2)	Usual care (<i>k</i> =3)	Pre- and post-MBS 1. RHR 2. DBP 3. SBP	Pre- and post-MBS intervention results merged 1. MD: -3.06 bpm [-5.65; -0.47], <i>I</i> ² =0%; <i>n</i> =111, <i>k</i> =4 (8 arms; very low level of evidence) 2. MD: -1.31 mmHg [-2.33; -0.29], <i>I</i> ² =23%; <i>n</i> =251, <i>k</i> =6 (13 arms; very low level of evidence) 3. MD: -1.59 mmHg [-3.74; 0.56], <i>I</i> ² =27%; <i>n</i> =239, <i>k</i> =6 (5 with 1 outlier removed; 13 arms) (very low level of evidence)	Overall risk of bias not reported (Cochrane risk of bias tool)	Low
Systematic Literature Reviews								
Schurmans 2022 (Belgium) <i>k</i> =4 including 2 publications with the same intervention	RCT (<i>k</i> =4)	<i>N</i> = 104 (<i>k</i> =2) 41.1-48.1 years 44.4 to 47.8 kg.m ² 75-100% women	Exercise training (<i>k</i> =2) PA counseling (<i>k</i> =2) <u>Duration:</u> 6-12wk. <u>Type:</u> E/R, E <u>Frequency:</u> 3-7/wk. <u>Intensity:</u> Moderate to vigorous <u>Session duration:</u> 30-80 min	Usual care (<i>k</i> =4)	1. WL 2. BMI 3. FFM 4. BP 5. 6MWTD 6. QoL 7. RHR	1. NS (<i>k</i> =1) 4. NS (<i>k</i> =2) 5. + (<i>k</i> =1) 6. + (<i>k</i> =1 except role-emotional domain) 7. NS (<i>k</i> =2) 1yr after MBS 2. NS (<i>k</i> =1) 3. + (<i>k</i> =1) 5. NS (<i>k</i> =1) 6. NS (<i>k</i> =1)	3/10 (<i>k</i> =1) Good (6/10 <i>k</i> =2; 7/10 <i>k</i> =1) (PEDro scale)	Low

		<u>Supervision:</u> supervised (<i>k</i> =2), unsupervised (<i>k</i> =2)						
Jabbour 2022 (Qatar+2)	RCT (<i>k</i> =10) Intervention trial (<i>k</i> =3) <i>k</i> =13 including 2 and 3 publications with the same intervention	<i>N</i> = 261 (<i>k</i> =10) 37.3-50.1 years 43.9 to 51.4 kg.m ² 67-100% women	Exercise training (<i>k</i> =11) PA counseling (<i>k</i> =3) <u>Duration:</u> 30d.-24 wk. <u>Type:</u> E, R, E/R, Water exerc, respiratory. <u>Frequency:</u> 1-7x/wk. <u>Intensity:</u> Low to vigorous <u>Session duration:</u> 25- 219 min <u>Supervision:</u> supervised (<i>k</i> =11), unsupervised (<i>k</i> =2)	Usual care (<i>k</i> =8) A 3-day outpatient control Period (<i>k</i> =1) NR (<i>k</i> =5)	1. BW 2. BMI 3. FM% 4. 6MWTD 5. Sit-to-stand 6. Arm curl 7. Leg strength/ muscle quality 8. Maximum aerobic capacity 9. SI 10. Adipokines 11. SBP 12. DBP 13. Length of hospital stay	Difficult interpretation of text and table results 1. NS (<i>k</i> =3), + (<i>k</i> =2) * 2. NS (<i>k</i> =1), + (<i>k</i> =3) 3. NS (<i>k</i> =1) 4. NS (<i>k</i> =1), + (<i>k</i> =2) 5. NS (<i>k</i> =1), + (<i>k</i> =1) 6. + (<i>k</i> =2) 7. + (<i>k</i> =3)* 8. NS (<i>k</i> =1, METS), + (<i>k</i> =1, VO ₂ peak) 9. NS (<i>k</i> =1) 10. NS (<i>k</i> =1) 11. NS (<i>k</i> =2), + (<i>k</i> =1) 12. NS (<i>k</i> =2), + (<i>k</i> =1) 13. + (<i>k</i> =1) *Daniels 2018 removed from results as it is a postMBS intervention	No evaluation	Critically low
Lodewijks 2022 (Netherlands)	RCT (<i>k</i> =8) NRCT (<i>k</i> =3) <i>k</i> =11 including 2 publications each from 2 interventions	<i>N</i> = 305 (<i>k</i> =9) NO SYNTHESIS ONLY FOR EXERCISE	Exercise training (<i>k</i> =7) PA counseling (<i>k</i> =4) <u>Duration:</u> 2 wk.-52wk. NO OTHER INFORMATION COLLECTED AND REPORTED	Usual care (<i>k</i> =7) No intervention (<i>k</i> =4)	1. Pre-MBS WL 2. Post-MBS WL 3. BMI 4. FM 5. Physical activity 6. QoL 7. FFM	1. NS (<i>k</i> =9), + (<i>k</i> =1)* 2. NS (<i>k</i> =1)* 3. + (<i>k</i> =1) 4. + (<i>k</i> =1) 5. + (<i>k</i> =4) 6. + (<i>k</i> =2, 1 for physical functioning, general health perceptions, mental health and social functioning, and 1 for all except role- emotional) 1-year postMBS 2. + (<i>k</i> =1)* 3. + (<i>k</i> =1) 5. + (<i>k</i> =1) 7. + (<i>k</i> =1) *Unclear if comparison is pre- vs post-MBS in intervention group or ex vs c	3/ NR (<i>k</i> =4) 2/5 (<i>k</i> =4) (Jadad) Serious bias (<i>k</i> =1) Moderate bias (<i>k</i> =2) (ROBINS-1 tool)	Low
Bellicha 2021 (France+6)	RCT (<i>k</i> =3) NRCT (<i>k</i> =1) <i>k</i> = 4 including 2 publications	<i>N</i> = 104 (<i>k</i> =3) 37.5 to 50.1 years 41.5 to 50.8 kg.m ²	Exercise training (<i>k</i> =4) <u>Duration:</u> 12-16 wk. <u>Type:</u> E, E/R, HIIT/R <u>Frequency:</u> 2-4x/wk. <u>Intensity:</u> Moderate to vigorous	Usual care (<i>k</i> =4)	1. BW/BMI 2. FM 3. LBM 4. Muscle Strength 5. Walking distance	1. NS (<i>k</i> =1), + (<i>k</i> =2) 2. NS (<i>k</i> =1), + (<i>k</i> =1), 3. NS (<i>k</i> =1) 4. NS (<i>k</i> =1), + (<i>k</i> =1) 5. + (<i>k</i> =2) 6. NS (<i>k</i> =2), + (<i>k</i> =1) 7. NS (<i>k</i> =1), + (<i>k</i> =1)	Good (<i>k</i> =1) Fair (<i>k</i> =1) Poor (<i>k</i> =1) (National Heart, Lung and Blood	Critically low

with the same intervention		76-90% women	<u>Session duration:</u> 25-50 min <u>Supervision:</u> Full-supervised ($k=4$)		6. VO ₂ max 7. QoL 8. BP 9. Glucose metabolism 10. Lipid profile 11. Habitual physical activity	8. NS ($k=2$), + ($k=1$) 9. NS ($k=1$), + ($k=1$) 10. NS ($k=1$), + ($k=1$) 1 yr post-MBS effects in one publication 1. + ($k=1$) 2. NS ($k=1$) 3. - ($k=1$) 4. NS ($k=1$) 5. + ($k=1$) 6. NS ($k=1$) 7. NS ($k=1$) 8. NS ($k=1$) 11. + ($k=1$)	<i>Institute quality assessment tool</i>	
Pouwels 2015 (Netherlands) $k=4$	RCT ($k=1$) NRCT ($k=3$)	NO SYNTHESIS ONLY FOR PRE-MBS	Exercise training ($k=4$) <u>Duration:</u> 1-24 wk. <u>Type:</u> E, E/R, Water exercise. <u>Frequency:</u> 1-7/wk. <u>Intensity:</u> Low to vigorous <u>Session duration:</u> 60-219 min <u>Supervision:</u> Full-supervised ($k=3$), partial ($k=1$)	NR	1. BW/BMI 2. FM 3. BP 4. Glucose metabolism 5. Lipid profile 6. Walking distance 7. VO ₂ max 8. QoL	Unclear: only one RCT with no systematic clear comparison between ex vs. c in the original article due to small sample size lack of control groups	Good 6/10 ($k=1$) Fair (5/10 $k=2$; 4/10 $k=1$) (PEDro scale)	Critically low
Baillot 2014 (Canada) $k=3$	RCT ($k=1$) UCT ($k=2$)	$N = 46$ 29.6 to 49.3 years 48.5 to 48.7 kg.m ² 0-86% women	Exercise training ($k=3$) <u>Duration:</u> 1-24 wk. <u>Type:</u> E, Water exercise. <u>Frequency:</u> 1-7/wk. <u>Intensity:</u> Low to vigorous <u>Session duration:</u> 30-60 min <u>Supervision:</u> Full-supervised ($k=3$)	NR	1. BW/BMI 2. Walking distance 3. Insulin action 4. BP 5. QoL/Depression scores 6. Bodily pain 7. Cardiovascular risk	Unclear as only one RCT with no systematic clear comparison between ex vs. c in the original article due to small sample size and other articles lack of control group	Moderate ($k=2$) Weak ($k=1$) (<i>Effective Public Health Practice Project</i>)	Low
Fonseca-Junior 2013 (Brazil) $k=1$	NR	$N = 61$ NR	Exercise training ($k=1$) <u>Duration:</u> 24 wk. <u>Type:</u> E <u>Frequency:</u> 1/wk. <u>Intensity:</u> Low <u>Session duration:</u> 30 min	NR	1. BMI 2. BP 3. Glucose metabolism 4. Lipid profile 5. Functional capacity	Explicit comparison between ex vs control groups not made clear in text and no table to reinforce data	No evaluation	Critically low

Supervision: Full-
supervised

Note. All data were reported as they were presented in the manuscript. Details presented in blue were obtained directly or calculated from tables/figures in the manuscripts rather than from the text. N =Total sample size, n =subsample size, k =number of studies, RCT=randomized control trial, NCRT=non randomized control trial, E=endurance, R=resistance, E/R=combination endurance and resistance, HIIT=high intensity interval training, BC=behavioral component, MBS=metabolic and bariatric surgery, BW=body weight, WL=weight loss, BMI=body mass index, WL=weight loss, FM=fat mass, FFM=fat-free mass, LBM=lean body mass, 6MWTD=6 minute walking test difference, QoL=quality of life, BP=blood pressure, SBP=systolic blood pressure, DBP=diastolic blood pressure, RHR=resting heart rate, RR=risk ratio, MD=mean difference, SMD=standardized mean difference, NR=not reported, N/A=not applicable, NS=non-significant.

^a First country listed in first author's affiliations plus number of additional countries in all authors' reported affiliations; ^b Values reported from details provided in tables include minimum and maximum or mean/median age and BMI, and % women; ^c only outcomes with a synthesis within the meta-analysis or in the systematic review text are reported; ^d For meta-analyses, summary includes effect size estimate, 95% confidence interval, I^2 = values of heterogeneity, sample size (n) and number of studies (k) included in the analysis. For systematic reviews, NS indicates no significant difference, while + and - indicate a significant improvement and deterioration respectively, following intervention compared to control.

Appendix I – Pre-MBS Primary Articles

Table S13: Pre-MBS Primary Articles

All Articles	Baillot et al 2022	Durey et al 2022	Herrera-Santelices et al 2022	Marshall et al 2020	Schurmans et al 2022	Jabbour et al 2022	Lodewijks et al 2022	Bellicha et al 2021	Pouwels et al 2015	Baillot et al 2014	Fonseca-Junior et al 2013	Total Count
Arman et al 2021	X		X				X					3
Baillot et al 2014				X		X			X			3
Baillot et al 2016	X		X		X	X	X	X				6
Baillot et al 2017						X	X					2
Baillot et al 2018		X	X	X	X	X	X	X				7
Bond et al 2015a				X		X	X					3
Bond et al 2015b				X	X	X	X					4
Bond et al 2016				X								1
Bond et al 2017a		X		X	X	X						4
Bond et al 2017b				X								1
Creel et al 2016		X		X			X					3
Daniels et al 2018						X						1
Funderburk et al 2010	X		X			X	X		X	X		6
Garcia Delegado et al 2021						X						1
Gilbertson et al 2020	X					X	X					3
Hickey et al 1999									X	X		2
Kwok et al 2016		X										1
Li et al 2013		X										1
Marc-Hernandez et al 2019	X						X	X				3
Marcon et al 2011						X			X	X	X	4

Marcon et al 2017	X		X			X		X			4
Parikh et al 2012							X				1
Pico-Sirvent et al 2019	X										1

Note. Black = primary studies published after systematic review/meta-analysis and therefore not possible for inclusion.

Grey = primary articles published the same year as the systematic review/meta-analysis and therefore not likely for inclusion.

1. Arman N, Tokgoz G, Seyit H, Karabulut M. The effects of core stabilization exercise program in obese people awaiting bariatric surgery: A randomized controlled study. *Complement Ther Clin Pract*. 2021;43:101342. doi:10.1016/j.ctcp.2021.101342
2. Baillot A, Mampuya WM, Comeau E, Méziat-Burdin A, Langlois MF. Feasibility and impacts of supervised exercise training in subjects with obesity awaiting bariatric surgery: A pilot study. *Obesity surgery*. 2014;23:882-91.
3. Baillot A, Mampuya WM, Dionne IJ, Comeau E, Méziat-Burdin A, Langlois MF. Impacts of supervised exercise training in addition to interdisciplinary lifestyle management in subjects awaiting bariatric surgery: A randomized controlled study. *Obes Surg*. 2016;26(11):2602-2610. doi:10.1007/s11695-016-2153-9
4. Baillot A, Boissy P, Tousignant M, Langlois MF. Feasibility and effect of in-home physical exercise training delivered via telehealth before bariatric surgery. *J Telemed Telecare*. 2017;23(5):529-535. doi:10.1177/1357633X16653511
5. Baillot A, Vallée CA, Mampuya WM, et al. Effects of a pre-surgery supervised exercise training 1 year after bariatric surgery: A randomized controlled study. *Obes Surg*. 2018;28(4):955-962. doi:10.1007/s11695-017-2943-8
6. Bond DS, Thomas JG, King WC, et al. Exercise improves quality of life in bariatric surgery candidates: Results from the Bari-Active trial. *Obesity*. 2015;23(3):536-542. doi:10.1002/oby.20988
7. Bond DS, Thomas JG, Vithiananthan S, et al. Intervention-related increases in preoperative physical activity are maintained 6-months after Bariatric surgery: results from the bari-active trial. *Int J Obes*. 2017;41(3):467-470. doi:10.1038/ijo.2016.237
8. Bond DS, Vithiananthan S, Thomas JG, et al. Bari-Active: A randomized controlled trial of a preoperative intervention to increase physical activity in bariatric surgery patients. *Surg Obes Relat Dis*. 2015;11(1):169-177. doi:10.1016/j.soard.2014.07.010
9. Bond DS, Raynor HA, Thomas JG, et al. Greater adherence to recommended morning physical activity is associated with greater total intervention-related physical activity changes in bariatric surgery patients. *J Phys Act Health*. 2017;14(6):492-498. doi:10.1123/jpah.2016-0529
10. Bond DS, Graham Thomas J, Vithiananthan S, et al. Changes in enjoyment, self-efficacy, and motivation during a randomized trial to promote habitual physical activity adoption in bariatric surgery patients. *Surg Obes Relat Dis*. 2016;12(5):1072-1079. doi:10.1016/j.soard.2016.02.009
11. Creel DB, Schuh LM, Reed CA, et al. A randomized trial comparing two interventions to increase physical activity among patients undergoing bariatric surgery. *Obesity*. 2016;24(8):1660-1668. doi:10.1002/oby.21548

12. Daniels P, Burns RD, Brusseau TA, et al. Effect of a randomised 12-week resistance training programme on muscular strength, cross-sectional area and muscle quality in women having undergone Roux-en-Y gastric bypass. *J Sports Sci.* 2018;36(5):529-535. doi:10.1080/02640414.2017.1322217
13. Funderburk JA, Callis S. Aquatic intervention effect on quality of life prior to obesity surgery: a pilot study. [Conference Abstract]. *Annu Ther Recreat.* 2010;18:66-78.
14. García-Delgado Y, López-Madrado-Hernández MJ, Alvarado-Martel D, et al. prehabilitation for bariatric surgery: A randomized, controlled trial protocol and pilot study. *Nutrients.* 2021;13(9):2903. doi:10.3390/nu13092903
15. Gilbertson NM, Gaitán JM, Osinski V, et al. Pre-operative aerobic exercise on metabolic health and surgical outcomes in patients receiving bariatric surgery: A pilot trial. *PLOS ONE.* 2020;15(10):e0239130. doi:10.1371/journal.pone.0239130
16. Hickey M, Gavigan KE, McCammon MR, et al. Effects of 7 days of exercise training on insulin action in morbidly obese men. *Clin Exerc Physiol.* 1:24-28.
17. Kwok K, Hardy K, Bouchard D, Vergis A. The impact of a pre-operative exercise program on patients awaiting bariatric surgery. [Conference Abstract]. *Surg Obes Relat Dis.* 2016;12(7):S201-S202. doi:10.1016/j.soard.2016.08.350
18. Li C, Zavorsky GS, Kim DJ, Christou NV, Feldman LS, Carli F. Effects of a bariatric preoperative exercise program: A pilot randomized study. [Conference Abstract]. *Surg Endosc Interv Tech.* 2013;27:S431.
19. Marc-Hernández A, Ruiz-Tovar J, Aracil A, Guillén S, Moya-Ramón M. Impact of exercise on body composition and cardiometabolic risk factors in patients awaiting bariatric surgery. *Obes Surg.* 2019;29(12):3891-3900. doi:10.1007/s11695-019-04088-9
20. Marcon ER, Baglioni S, Bittencourt L, Lopes CLN, Neumann CR, Trindade MRM. What is the best treatment before bariatric surgery? Exercise, exercise and group therapy, or conventional waiting: A randomized controlled trial. *Obes Surg.* 2017;27(3):763-773. doi:10.1007/s11695-016-2365-z
21. Marcon ER, Gus I, Neumann CR. Impact of a minimum program of supervised exercises in the cardiometabolic risk in patients with morbid obesity. *Arq Bras Endocrinol Metab.* 2011;55(5):331-338. doi:10.1590/S0004-27302011000500006
22. Parikh M, Dasari M, McMacken M, Ren C, Fielding G, Ogedegbe G. Does a preoperative medically supervised weight loss program improve bariatric surgery outcomes? A pilot randomized study. *Surg Endosc.* 2012;26(3):853-861. doi:10.1007/s00464-011-1966-9
23. Picó-Sirvent I, Aracil-Marco A, Pastor D, Moya-Ramón M. Effects of a combined high-intensity interval training and resistance training program in patients awaiting bariatric surgery: A pilot study. *Sports.* 2019;7(3):72. doi:10.3390/sports7030072

Appendix J – Post-MBS Study Characteristics

Table S14: Summary of Meta-Analyses and Systematic Reviews for Post-Metabolic and Bariatric Surgery Exercise Interventions

Author Year (Country ^a)	Study Designs	Sample Description (N, Age Range, BMI Range, % Women) ^b	Intervention Description	Control Group	Outcomes Assessed ^c	Main Results ^d Level of Evidence	Quality Scores of the Studies Included (Quality Assessment Tool) ^e	AMSTAR Rating
Meta-Analyses								
Bond 2023 (USA+1) k=5	RCT (k=5)	N = 189 47.8 ± 4.2 years 36.1 ± 6.3.8 kg/m ² 83.2 ± 9.5% women	Exercise training (k=5) <u>Duration:</u> 19.2 ± 7.0 (12 - 26) weeks <u>Type:</u> E/R, E, HIIT <u>Frequency:</u> 2-5x/wk. <u>Intensity:</u> Moderate to vigorous <u>Session duration:</u> 40-80 min <u>Supervision:</u> supervised k=4, semi supervised k=1 <u>Start:</u> for post-MBS intervention = 18.9 ± 11.5 (12 - 37) months	Usual care (k=4) Standard diet and PA recommendations (k=1)	1. WL (≥12 months postMBS)	1. SMD: - 2.26 [-2.07; 1.55], I ² =0%; n=189, k=5	High risk (k=3) Some concerns (k=1) Low risk (k=1) (Cochrane risk of bias tool 2)	Low
Baillot 2022 (Canada+1) k=21	RCT (k=15) NRCT (k=6)	NO SYNTHESIS ONLY FOR POST-MBS	Exercise training (k=21) <u>Duration:</u> 4-104 weeks <u>Type:</u> R, E, E/R, HIIT balance <u>Frequency:</u> 2-5x/wk. <u>Intensity:</u> Moderate to vigorous <u>Session duration:</u> 25-90 min <u>Supervision:</u> supervised k=14, semi supervised k=4, unsupervised k=1, not reported k=2 <u>Start:</u> for post-MBS intervention = 0-7 years	Usual care (k=13) Diet education (k=1) Instruction to continue normal activities (k=1) Protein supplementation (k=1)* Placebo (k=1)* Not reported (k=5) *1 study with two control groups	1. Adverse events during exercise 2. Attendance 3. Dropout 4. Compliance	1. None (k=7), occasional pain, fatigue, or dyspnea (k=3), hypoglycemia or hypotension (k=1), back bruise after a fall (k= 1) 2. Pooled percentage: 87.4% [76.7; 95.6], I ² =0%; n=NR, k=6 3. Pooled percentage: 5.6% [0.6; 13.8], I ² =69%; n=NR, k=12 4. No metaanalysis due to small number of studies (k=3) and various operationalization of term “compliance” <u>Sub-analysis:</u> NS differences in attendance, dropout rates based on studies quality, exercise intervention timing, and exercise intervention duration	Poor (k=7) Fair (k=7) Good (k=7) (National Heart, Lung and Blood Institute quality assessment tool)	Low

Roth 2022 (Switzerland) k=6	RCT (k=6)	N = 433 35.4-42.5 years 41.7to 49.8kg.m ² 55-100% women	Exercise training (k=6) <u>Duration:</u> 12-104 wk. <u>Type:</u> R, E, E/R <u>Frequency:</u> 2-3x/wk. <u>Intensity:</u> Moderate to vigorous <u>Session duration:</u> 20-85 min <u>Supervision:</u> NR <u>Start:</u> Combined synthesis NR individual results	Control (k=4) High protein (k=2)	1. FFM 2. BMD	1. Ex. vs. C SMD: 0.39 [-0.01; 0.78], I ² =0%; n=132, k=3 (Very Low level of evidence) 1. Ex+Protein vs. Protein, SMD: 0.25 [-1.15; 1.65], I ² =0%; n=91, k=2 (Low level of evidence) 1. Ex+Protein+vit. D+Ca ²⁺ vs. Control, SMD: 5.16 [4.60; 5.71], I ² =N/A; n=220, k=1 (Moderate level of evidence) 2. Ex vs. C, SMD:0.51 [0.01; 1.01], I ² =N/A; n=63, k=1 (Moderate level of evidence) 2. Ex+Protein+vit. D+Ca ²⁺ vs. Control, SMD: 3.88 [3.43; 4.34], I ² =N/A; n=220, k=1 (Moderate level of evidence)	High risk (k=2) Some concerns (k=1) Low risk (k=3) (Cochrane risk of bias tool 2)	Moderate
Vieira 2022 (Brazil) k=15	RCT (k=10) NRCT (k=5)	N = 638 18-65 years majority>30 kg.m ² up to 100% women	Exercise training (k=15) <u>Duration:</u> 12-54 wk. <u>Type:</u> R, E/R, respiratory <u>Frequency:</u> 2-5x/wk. <u>Intensity:</u> Moderate to vigorous <u>Session duration:</u> 5-110min <u>Supervision:</u> Full-supervised (k=11), partial (k=2), unsupervised (k=1), NR (k=1) <u>Start:</u> 1-84 months after MBS	Usual care (k=11) Usual care + PA encouragement (k=4)	1. Upper muscle strength (1RM) 2. Lower muscle strength (1RM) 3. Muscle strength (sit-to stand) 4. Muscle strength (dynamometer) 5. Muscle strength (handgrip test)	1. ES: 0.71 [0.41; 1.01], I ² =0%; n=NR, k=4 (Very low level of evidence) 2. ES: 1.37 [0.84; 1.91], I ² =46%; n=NR, k=5 (Very low level of evidence) 3. ES: 0.60 [0.20; 1.01], I ² =69%; n=NR, k=8 (Very low level of evidence) 4. ES: 0.46 [0.06; 0.87], I ² =31%; n=NR, k=4 (Very low level of evidence) 5. ES: 0.11 [-0.42; 0.63], I ² =73%; n=NR, k=6 (Very low level of evidence)	None of the studies had a low risk of bias (Joanna Briggs Institute critical appraisal tools)	Low
Gasmi 2022 (France+7) k=11 including 3 publications with the same intervention	RCT (k=11)	N = 495 (k= 9) 31.0-50.6 years NR kg.m ² NR women	Exercise training (k=10) Physical activity (k=1) WBS + dynamic exercise (k=1) <u>Duration:</u> NR <u>Type:</u> E, R, E/R <u>Frequency:</u> NR <u>Intensity:</u> NR clearly <u>Session duration:</u> NR <u>Supervision:</u> NR <u>Start:</u> early-3 years after MBS	Usual care (k=1) Control group (k=10)	1. BMI 2. FFM 3. FM 4. Hip to waist 5. WC	1. SMD: -0.93 [-1.65; -0.20], I ² =85%; n=341, k=5 2. SMD: 0.23 [-0.31; 0.77], I ² =0%; n=54, k=2 3. SMD: -0.08 [-0.54; 0.38], I ² =0%; n=74, k=3 4. SMD: -0.25 [-0.76; 0.26], I ² =0%; n=60, k=2 5. SMD: -0.18 [-0.79; 0.43], I ² =0%; n=42, k=2 2,3,4,5 based on final data intervention vs. control groups	No results reported (Cochrane risk of bias tool)	Critically low
Diniz-Sousa 2022 (Portugal+1) k=4	RCT (k=3) NRCT (k=1)	N = 340 37-47 years Pre-MBS 41.8-49.8 kg.m ² 72% women	Exercise training (k=4) <u>Duration:</u> 6-11.5 months <u>Type:</u> R, E/R, high impact/balance <u>Frequency:</u> 2-5x/wk. <u>Intensity:</u> Moderate to vigorous <u>Session duration:</u> 30-90 min <u>Supervision:</u> Full-supervised (k=3), semi-supervised (k=1)	Usual care (k=4)	1. BMD total hip 2. BMD femoral neck 3. BMD lumbar spine 4. BMD 1/3 radius	1. SMD: 0.37 [0.02; 0.71], I ² =50%; n=340, k=4 (very low certainty evidence) 2. SMD: 0.63 [0.19; 1.06], I ² =19%; n=112, k=2 (low certainty evidence) 3. SMD: 0.41 [0.19; 0.62], I ² =0%; n=341, k=4 (low certainty evidence) 4. SMD: 0.58 [0.19; 0.97], I ² =0%; n=112, k=2 (low certainty evidence)	Some concerns (k=3) (Cochrane risk of bias tool 2) Moderate risk of bias (k=1)	Moderate

<u>Start:</u> 2 wk.-3 months after MBS					(ROBINS-I)			
Boppre 2022 (Portugual)	k=11 including 3, 2 and 2 publications with the same intervention	RCT (k=11)	N = 386 (k = 7) 36.0-53.9 years NR kg.m² 84% women	Exercise training (k=11) <u>Duration:</u> 12-26 wk. <u>Type:</u> E, E/R <u>Frequency:</u> 2-5x/wk. <u>Intensity:</u> Moderate <u>Session duration:</u> 40-60 min <u>Supervision:</u> NR <u>Start:</u> 1-24 months after MBS	Usual care (k=11)	1. MD: 0.26 L/min [-0.11; 0.63], I²=0%; n=, k=3 2. MD: - 5.33 mmHg [-8.99; -1.66], I²=0%; n=314, k=6 (moderate certainty evidence) 3. MD: -2.66 mmHg [-6.72; 1.40], I²=59%; n=NR, k=6 4. MD: -2.05 bpm [-6.64; 2.54], I²=0%; n= NR, k=3 5. MD: -1.58 µIU/mL [-5.14; 1.98], I²=71%; n= NR, k=4 6. MD: 0.94 mg/dL [-3.31; 5.19], I²=0%; n= NR, k=4 7. MD: 1.39 [-1.30; 4.08], I²=89%; n= NR, k=2 8. MD: -0.65 mmol/mol [-2.22; 0.93], I²=0%; n= NR, k=2 9. MD: -3.08 mg/dL [-12.04; 5.87], I²=0%; n= NR, k=5 10. MD: 0.61 mg/dL [-3.05; 4.28], I²=26%; n= NR, k=5 11. MD: -8.17 mg/dL [-20.35; 4.00], I²=57%; n= NR, k=5 12. MD: -8.38 mg/dL [-19.81; 3.04], I²=0%; n= NR, k=5	High risk (k=3) Some concerns (k=1) Low risk (k=7)	Critically low
					Sub-analysis: Endurance exercise vs MBS: NS change for VO₂max (k=2), SBP (k=3), DBP (k=3), insulin (k=2), glucose (k=2), TC (k=2), HDL-C (k=2), LDL-C (k=2), and TG (k=2) Combined exercise vs. MBS: 2. MD: -7.18 mmHg [-12.42; -1.94], I²=0%; n=137, k=3 (moderate certainty evidence) 12. MD: -17.56 mg/dL [-34.15; -0.96], I²=0%; n=171, k=3 (low certainty evidence) NS change for DBP (k=3), insulin (k=2), glucose (k=2), HbA1C (k=2), TC (k=3), HDL-C (k=3), and LDL-C (k=3)			
					Studies starting < 6 months after MBS = NS change for SBP (k=4), DBP (k=4), insulin (k=3), glucose (k=3), TC (k=4), HDL-C (k=4), LDL-C (k=4), and TG (k=4) Studies starting > 6 months after MBS:			

					<p>2. SBP, MD: -7.71 mmHg [-13.12; -2.31], I²=0%; n=84, k=2 (high certainty evidence) NS change for DBP (k=2) and RHR (k=4)</p> <p>Intervention duration < 12 week = NS change for SBP (k=3), DBP (k=3), TC (k=2), HDL-C (k=2), LDL-C (k=2), and TG (k=2)</p> <p>Intervention duration > 12 week: 2. SBP, MD: -5.78 mmHg [-9.91; -1.66], I²=0%; n=212, k=3 (high certainty evidence) NS change for VO₂max (k=2), DBP (k=3), RHR (k=2), insulin (k=3), glucose (k=3), HOMA-IR (k=2), HbA1C (k=2), TC (k=3), HDL-C (k=3), LDL-C (k=3), and TG (k=3)</p>		
					<p>1. MD: -2.51 kg [-4.74; -0.27], I²=0%; n=496, k=10 (11 arms) 2. MD: -0.84 kg/m² [-1.60; -0.08], I²=0%; n=401, k=7 (8 arms) 3. MD: 0.49 kg [-1.71; 2.69], I²=0%; n=173, k=2 4. MD: 0.87 kg [-0.65; 2.40], I²=0%; n=201, k=3 5. MD: -4.14 cm [-8.16; -0.12], I²=9%; n=201, k=4</p> <p><u>Sub-analysis:</u> Endurance exercise vs. MBS: NS change for BW (k=4), BMI (k=3), LBM (k=2), and WC (k=3) Resistance exercise vs. MBS: NS change for BW (k=2) Combined exercise vs. MBS: 1. BW, MD: -5.02 kg [-8.13; -1.90], I²=0%; n=221, k=5 2. BMI, MD: -1.62 kg/m² [-2.72; -0.59], I²=0%; n=170, k=4</p> <p>Studies starting < 6 months after MBS: NS change for BW (k=7), BMI (k=5), FM (k=2), LBM (k=3), and WC (k=3) Studies starting > 6 months after MBS: 1. BW, MD: -5.25 kg [-8.52; -1.97], I²=0%; n=135, k=3 2. BMI, MD: -1.84 kg/m² [-3.04; -0.64], I²=0%; n=84, k=2</p> <p>Intervention duration ≤ 12 week NS change for BW (k=5), FM (k=3), and WC (k=1)</p>		
Boppre 2021 (Portugal) k=10	RCT (k=10)	<p>N = 487</p> <p>35.4-53.9 years</p> <p>NR kg.m²</p> <p>85% women</p>	<p>Exercise training (k=10) <u>Duration:</u> 12-26 wk. <u>Type:</u> E, E/R, R <u>Frequency:</u> 3-5x/wk. <u>Intensity:</u> Moderate-Vigorous <u>Session duration:</u> 40-80 min <u>Supervision:</u> Full-supervised (k=6), partial-supervised (k=1), semi-supervised (k=3) <u>Start:</u> 1-24 months after MBS</p>	Usual care (k=10)	<p>1. BW 2. BMI 3. FM 4. LBM 5. WC</p>	<p>Fair 5/10 (k=6) Good 6/10 (k=4) (PEDro scale)</p> <p>Critically low</p>	

					Intervention duration >12 week: NS change for BW (k=4), BMI (k=4), FM (k=3), LBM (k=2), and WC (k=2)				
Bellicha 2021 (France+6)		N = 587							
k=14 included in meta-analysis ^f	RCT (k=9) NRCT (k=5)	33.3-53.9 years 29.6-47.9 kg.m ² 57-100% women	Exercise training (k=14) <u>Duration:</u> 3-10 months <u>Type:</u> E, R, E/R <u>Frequency:</u> 2-5x/wk. <u>Intensity:</u> Moderate to vigorous <u>Session duration:</u> 20-60 min <u>Supervision:</u> Totally supervised (k=10), partially supervised (k=2), not supervised (k=1), NR (k=1) <u>Start:</u> 1-42 months after MBS	Usual care (k=14)	1. BW 2. FM 3. LBM 4. Muscle Strength 5. Walking distance 6. VO ₂ max 7. QoL physical 8. QoL mental 9. HOMA-IR 10.HDL-C 11.LDL-C 12. TG 13. BMD 14. SBP 15. DBP 16. MVPA	1. MD: -1.8 kg [-3.2; -0.4], I ² =35%; n=NR, k=13 (14 arms) 2. MD: -2.1 kg [-3.7; -0.5], I ² =50%; n=NR, k=8 (9 arms) 3. MD: 0.7 kg [-0.2; 1.6], I ² =45%; n=NR, k=10 (11 arms) 4. SMD: 0.82 [0.48; 1.16], I ² =42%; n=NR, k=8 (9 arms) 5. SMD: 1.46 [0.27; 2.66], I ² =89%; n=NR, k=5 (6 arms) 6. SMD: 0.70 [0.35; 1.06], I ² =42%; n=NR, k=8 7. MD: -2.5 [-5.1; 0.2], I ² =0%; n=NR, k=2 8. MD: 3.9 [-0.5; 8.3], I ² =0%; n=NR, k=2 9. SMD: 0.14 [-0.10; 0.38], I ² =0%; n=NR, k=2 10. SMD: 0.10 [-0.16; 0.37], I ² =0%; n=NR, k=4 11. SMD: -0.18 [-0.46; 0.09], I ² =0%; n=NR, k=3 12. SMD: 0.01 [-0.26; 0.27], I ² =0%; n=NR, k=4 13. SMD: 0.44 [0.21; 0.67], I ² =0%; n=NR, k=3 14. MD: -4.2 mmHg [-9.3; 1.0], I ² =47%; n=NR, k=4 15. MD: -2.3 mmHg [-8.5; 3.9], I ² =77%; n=NR, k=4 16. MD: -0.20 min/d [-13.19; 12.79], I ² =77%; n=NR, k=3 Sub analyses to have MD 6. VO ₂ max relative to BW, MD: 2.73 mL/kg/min [0.81; 4.64], I ² =79%; n=NR, k=6	Poor (k=5) Fair (k=3) Good (k=6) (National Heart, Lung and Blood Institute quality assessment tool)	Critically low	
Carretero-Ruiz 2021 (Spain+2)		N = 469 (k = 11)							
k=14 in meta-analysis ^g including 6 publications from 3	RCT (k=10) NRCT (k=4)	36.0-53.9 years NR kg.m ² 66.6-100% women	Exercise training (k=14) <u>Duration:</u> 12-40 weeks <u>Type:</u> E, R, E/R <u>Frequency:</u> 2-5x/wk. <u>Intensity:</u> Moderate to vigorous <u>Session duration:</u> 25-75 min <u>Supervision:</u> Supervised (k=10), supervised/programmed (k=4)	Usual care (k=14)	1. VO ₂ max/peak relative to BW 2. VO ₂ max/peak 3. RHR 4. HDL-C 5.SBP 6. DBP	1. ES: 0.67 [0.29; 1.06] (MD: 1.25 ml/kg/min [0.48; 2.02]), I ² =23%; n=NR, k=6 2. ES: 0.32 [0.07; 0.57], I ² =0%; n=NR, k=5 3. ES: -0.44 [-0.75; -0.02] (MD: -3.93 bpm [-6.54; 1.31]), I ² =0%; n=NR, k=5 4. ES: 0.22 [0.01; 0.43], I ² =0%; n=NR, k=6 5. ES: -0.16 [-0.40; 0.08] (MD = -2.65 mmHg [-7.32; -1.11]), I ² =0%; n=NR, k=5 6. ES: -0.12 [-0.45, 0.21] (MD: -1.41 mmHg [-5.56, 2.75]), I ² =34%; n=NR, k=5	Poor (3/10 k=2) Fair (4/10 k=1; 5/10 k=3) Good (7/10 k=4; 8/10 k=3; 9/10 k=1) (PEDro scale)	Critically low	

interventions (2 each)		<u>Start:</u> 1-102 months after MBS			
Marshall (2020) (Australia)	RCT (k=4) Pseudo RCT (k=1) NRCT (k=2)	N = 282 36-54 years 32.7-45.6 kg.m ² 70-100% women	Exercise training (k=7) <u>Duration:</u> 12-26 weeks <u>Type:</u> E, R, E/R <u>Frequency:</u> 2-5x/wk. <u>Intensity:</u> NR <u>Session duration:</u> 40-80 min <u>Supervision:</u> Supervised (k=6), Semi-supervised (k=1) <u>Start:</u> 1-6 months after MBS	Usual care (k=4) Health education (k=3)	Pre- and post-MBS 1. RHR 2. DBP 3. SBP PostMBS 4. TG 5. HDL-C 6. LDL-C 7.TC 8.Fasting insulin 9.Fasting glucose
k=7 (resulting in 13 publications)					Pre- and post-MBS 1. MD: -3.06 bpm [-5.65; -0.47], I ² =0%; n=111, k=4* (very low level of evidence) 2. MD: -1.31 mmHg [-2.33; -0.29], I ² =23%; n=251, k=6 (very low level of evidence) 3. MD: -1.59 mmHg [-3.74; 0.56], I ² =27%; n=239, k=6 (5 with 1 outlier removed) (very low level of evidence) Post-MBS 4. MD: 0.01 mmol/L [-0.15; 0.16], I ² =0%; n=180, k=2*** (low level of evidence) 5. MD: -0.00 mmol/L [-0.01; 0.01], I ² =0%; n=180, k=2*** (low level of evidence) 6. MD: -0.06 mmol/L [-0.21; 0.09], I ² =0%; n=180, k=2*** (low level of evidence) 7. MD: -0.08 mmol/L [-0.26; 0.11], I ² =0%; n=180, k=2*** (low level of evidence) 8. MD: 4.88 pmol/L [-2.09; 11.84], I ² =0%; n=180, k=2*** (low level of evidence) 9. MD: 0.05 mmol/L [-0.14; 0.24], I ² =0%; n=180, k=2*** (low level of evidence) * k=1 with 2 pre-MBS intervention publications and k=1 with 4 post-MBS intervention publications ** k=1 with 2 pre-MBS intervention publications and k=2 with 4 post-MBS intervention publications each *** k=2 with 4 publications each
Carretero-Ruiz 2019 (Spain+2)	RCT (k=10) NRCT (k=6)	N = 604 33.3 -53.9 years 29.6 - 47.8 kg.m ² 66.6 -100% women	Exercise training (k=14) Physiotherapy (k=1) Respiratory training (k=1) <u>Duration:</u> 1-40 wk. <u>Type:</u> E, R, E/R, Respiratory training, balance training <u>Frequency:</u> 2-5x/wk. <u>Intensity:</u> Low to vigorous	Usual care (k=16)	1. WL
k=16 included in meta-analysis ^h					1. SMD: 0.15 [-0.02; 0.32], I ² =0%; n=NR, k=16 (17 arms) <u>Sub-analysis:</u> Endurance exercise vs MBS: NS change (k=5) Resistance exercise vs MBS: NS change (k=2) Combined E/R exercise vs MBS: NS change (k=7) Alternative training: NS change (k=3)
					Overall risk of bias not reported (Cochrane risk of bias tool)
					Low
					High risk (k=7) Moderate risk (k=6) Low risk (k=3) (Cochrane risk of bias tool)
					Low

			<u>Session duration:</u> 30 - 80 min <u>Supervision:</u> Supervised/programmed ($k=4$), supervised ($k=7$), programmed ($k=4$), NR ($k=1$) <u>Start:</u> 1 day – 3 years after MBS			Programmed PA: NS change ($k=3$) Supervised PA: NS change ($k=9$) Programmed/Supervised PA: NS change ($k=4$) ≤ 3 months postMBS: NS change ($k=11$) >3 months posMBS: NS change ($k=5$) ≤ 16 weeks: NS change ($k=12$) >16 weeks: NS change ($k=5$) ≤ 150 min/week exercise: NS change ($k=4$) >150 min/week exercise: NS change ($k=13$) <u>Meta-regression analyses</u> Mean age: SMD: -0.00 [-0.04; 0.04], $I^2=0\%$; $n=NR$, $k=NR$ Length of intervention: SMD: 0.01 [-0.01; 0.00], $I^2=0\%$; $n=NR$, $k=NR$ Time per session: SMD: -0.00 [-0.01; 0.01], $I^2=0\%$; $n=NR$, $k=NR$.		
da Silva 2019 (Brazil/USA)	RCT ($k=7$) *error in reporting as $k=4$ are NRCTs*	$N = 193$ ($k = 6$) 38.5-53.9 years 98.2-130.8 kg (baseline weight) 87% women	Exercise training ($k=7$) <u>Duration:</u> 3-10 months <u>Type:</u> E, R, E/R <u>Frequency:</u> 2-5x/wk. <u>Intensity:</u> Moderate-vigorous <u>Session duration:</u> 44-75 min <u>Supervision:</u> NR <u>Start:</u> 1-102 months after MBS	NR	1. VO ₂ max	1. SMD: 0.43 [0.16; 0.70], $I^2=0\%$; $n=215$, $k=7$ 4/10 ($k=3$) 6/10 ($k=1$) 7/10 ($k=3$) (PEDro scale)	Critically low	
Ren 2018 (China)	RCT ($k=8$) $k=8$	$N = 347$ 36.0-53.9 years NR kg.m ² NR women	Exercise training ($k=8$) <u>Duration:</u> 12-26 wk. <u>Type:</u> E, R, E/R <u>Frequency:</u> 2-5/wk. <u>Intensity:</u> Moderate to vigorous <u>Session duration:</u> 1.3-4h/wk. <u>Supervision:</u> NR <u>Start:</u> immediately-19.3 months after MBS	Usual care/No exercise training ($k=8$)	1. BW 2. 6MWT 3. BMI 4. BF% 5. FM 6. FFM 7. WC 8. SBP 9. DBP 10. RHR 11. HC	1. WMD: -1.94 kg [-3.18; -0.69], $I^2=51\%$; $n=347$, $k=8$ (Moderate level of evidence) <u>Sub-analysis:</u> Endurance exercise vs MBS: NS change ($k=3$) Resistance exercise vs MBS: NS change ($k=1$) Combined E/R exercise: WMD: -3.12 kg [-4.56; - 1.68], $I^2=32\%$; $n=NR$, $k=4$ Studies starting < 6 months after MBS: WMD: -1.71 kg [-3.06; -0.36], $I^2=32\%$; $n=NR$, $k=5$ Studies starting ≥ 12 months after MBS: WMD: -3.63 kg [-5.35; -1.91], $I^2=0\%$; $n=NR$, $k=2$ Varying starting times: NS change ($k=1$)	No overall risk reported (Cochrane's collaboration tool)	Low

2. WMD: 29.67 m [25.97; 33.37], $I^2=0\%$; $n=65$, $k=2$ (Low level of evidence)
3. WMD: -0.40 kg/m² [-0.81; 0.00], $I^2=44\%$; $n=259$, $k=5$ (Moderate level of evidence)
4. WMD: -1.93% [-4.06; 0.20], $I^2=93\%$; $n=107$, $k=4$ (Low level of evidence)
5. WMD: -3.35 kg [-7.99; 1.29], $I^2=95\%$; $n=186$, $k=3$ (Low level of evidence)
6. WMD: 0.53 kg [-1.88; 2.94], $I^2=71\%$; $n=58$, $k=2$ (Very low level of evidence)
7. WMD: -5.25 cm [-10.48; -0.03], $I^2=94\%$; $n=198$, $k=4$ (Low level of evidence)
8. WMD: -4.12 mmHg [-6.68; -1.55], $I^2=6\%$; $n=229$, $k=4$ (Low level of evidence)
9. WMD: -3.56 mmHg [-8.61; 1.48], $I^2=83\%$; $n=229$, $k=4$ (Very low level of evidence)
10. WMD: -4.39 bpm [-8.11; -0.68], $I^2=0\%$; $n=94$, $k=3$ (Low level of evidence)
11. WMD: -3.91 cm [-11.09; 3.26], $I^2=88\%$; $n=70$, $k=3$ (Very low level of evidence)

Systematic Literature Reviews

Schurmans 2022 (Belgium) $k=16$ including 5, 3 and 2 publications with the same intervention	RCT ($k=15$) NRCT ($k=1$)	$N=403$ ($k=9$) 31.0-53.9 years 32.7-48.1 kg.m ² 66-100% women	Exercise training ($k=16$) <u>Duration</u> : 8 weeks-12 months <u>Type</u> : E, R, E/R <u>Frequency</u> : 2-5x/wk. <u>Intensity</u> : Moderate-vigorous <u>Session duration</u> : 30-80 min <u>Supervision</u> : Supervised ($k=4$), semi supervised ($k=10$), NR ($k=2$) <u>Start</u> : 8 weeks-12 months after MBS	Usual care ($k=16$)	1. BMI	1. NS ($k=7$), + ($k=3$; 1 only at 24 months)	3/10 ($k=1$) Fair (4/10 $k=2$; 5/10 $k=8$) Good (6/10 $k=3$; 7/10 $k=1$) NR ($k=1$) (PEDro scale)	Low
					2. WL	2. NS ($k=10$), + ($k=4$; 1 only at 24 months)		
					3. LBM	3. NS ($k=5$)		
					4. FFM	4. NS ($k=3$), + ($k=2$, 1 only for combined E/R vs control)		
					5. Muscle strength	5. + ($k=1$)		
					6. 6MWT	6. NS ($k=3$)		
					7. VO ₂ max	7. NS ($k=2$), + ($k=4$)		
					8. QoL	8. NS ($k=2$) except for general health domain		
					9. SBP	9. NS ($k=4$), + ($k=1$)		
					10. DBP	10. NS ($k=2$), + ($k=2$, 1 only at 24 months post-op)		
					11. Mean/RHR	11. NS ($k=3$)		
					12. Dyspnea perception	12. NS ($k=1$), + ($k=1$)		
					13. Leg exertion symptoms	13. NS ($k=1$)		
					14. Blood lipids	14. NS ($k=1$), + ($k=1$ for HDL-C)		
					15. Insulin sensitivity	15. NS ($k=1$), + ($k=3$)		
					16. AIRg/Di	16. NS ($k=1$), + ($k=1$)		
					17. SPISE	17. NS ($k=1$)		
					18. HOMA-IR	18. NS ($k=2$)		
						19. + ($k=1$)		
						20. NS ($k=1$)		

				19. Glucose effectiveness 20. Forced vital capacity				
Morales-Marroquin 2020 (USA) k=9	RCT (k=6) NRCT (k=3)	N = 344 33.3-53.6 years NR kg.m ² 100% W (k=2)	Exercise training (k=9) <u>Duration:</u> 12-36 wk. <u>Type:</u> R, E/R <u>Frequency:</u> 2-5x/wk. <u>Intensity:</u> Moderate to vigorous <u>Session duration:</u> 40-90 min <u>Supervision:</u> Supervised (k=7), not supervised (k=1), NR (k=1) <u>Start:</u> 1-12 months after MBS	NR	1. BW 2. FM 3. FFM 4. Muscle Strength 5. BMD	1. NS (k=4), + (k=2) 2. NS (k=4), + (k=2) 3. NS (k=5), + (k=1 only for combined E/R v control) 4. + (k=5, 1 only for combined E/R v control and 1 for exercise+protein supplementation) 5. + (k=2)	Fair (k=6) Good (k=3) (National Institutes of Health rating system)	Low
Civi Karaaslan 2020 (Turkey) k=7 including 2 publications with the same intervention	RCT (k=7)	N = 234 (k = 6) 33.9±13.1 years NR kg.m ² 100% W (k=3) remaining studies ≥80% female (k=4)	Exercise training (k=6) Physiotherapy (k=1) <u>Duration:</u> 4-26 wk. <u>Type:</u> E, R, E/R <u>Frequency:</u> 2-5x/wk. <u>Intensity:</u> Moderate to vigorous <u>Session duration:</u> 30-80 min <u>Supervision:</u> NR <u>Start:</u> 1month-3.5 y. after MBS	NR	NO NARRATIVE SYNTHESIS not the aim	NO NARRATIVE SYNTHESIS	Fair (4/10 k=2; 5/10 k=2) Good (6/10 k=3) (PEDro scale)	Critically low
Pouwels 2015 (Netherlands) k=3 (Berggren et al. not included because no control group)	RCT (k=2) Prospective trial (k=1)	No synthesis only for post-MBS and no details in tables	Exercise training (k=3) <u>Duration:</u> 12-16 weeks <u>Type:</u> E, E/R <u>Frequency:</u> 3-5x/week <u>Intensity:</u> Moderate-Vigorous <u>Session duration:</u> 30-60 min <u>Supervision:</u> Full-supervised (k=2), partial (k=1) <u>Start:</u> NR (k=4)	NR	1. BW 2. BMI 3. WC 4. HC 5. Cardiovascular risk 6. Aerobic capacity 7. Muscle strength 8. QoL	1. NS (k=2) 2. NS (k=1) 3. NS (k=1) 4. NS (k=1) 5. NS (k=1) 6. NS (k=1), 7. + (k=1) 8. + (k=1, emotional well-being, energy levels and mental QoL)	4/10 (k=1) 5/10 (k=1) 7/10 (k=2) (PEDro scale)	Critically low
Baillot 2014 (Canada) k=3	RCT (k=2) NRCT (k=1)	N = 64 36.0 to 53.9 years 40.4 to 45.6 kg.m ²	Exercise training (k=3) <u>Duration:</u> 12wk. <u>Type:</u> E, E/R <u>Frequency:</u> 3-5/wk. <u>Intensity:</u> Moderate - Vigorous	NR	1. FM 2. FFM 3. 6MWT 4. Muscle strength 5. QoL	1. NS (k=3, 1 is %FM) 2. NS (k=3) 3. NS (k=1), + (k=1) 4. + (k=1) 5. + (k=1)	Moderate (k=3) (Quality assessment tool Effective Public	Low

		57-100% women	<u>Session duration:</u> 20-60 min <u>Supervision:</u> Full-supervised (<i>k</i> =2), partial (<i>k</i> =1) <u>Start:</u> NR			Health Practice Project)	
Fonseca- Junior 2013 (Brazil) <i>k</i> =3	NR	<i>N</i> = 64 NR	Exercise training (<i>k</i> =3) <u>Duration:</u> 12weeks to 3 months <u>Type:</u> E, E/R <u>Frequency:</u> 3-5/wk. <u>Intensity:</u> Moderate - Vigorous <u>Session duration:</u> ~60 min <u>Supervision:</u> NR <u>Start:</u> NR	NR	1. DBP 2. Aerobic capacity 3. Functional capacity 4. Autonomous cardiac functional capacity 5. Post-pandrial glucose 6. Muscle strength 7. Weight loss	Explicit comparison between ex vs control groups not made clear in text and no table to reinforce data No evaluation	Critically low

Note. All data were reported as they were presented in the manuscript. Details presented in blue were obtained directly or calculated from tables/figures in the manuscripts rather than from the text. *N*=Total sample size, *n*=subsample size, *k*=number of studies, RCT=randomized control trial, NCRT=non randomized control trial, E=endurance, R=resistance, E/R=combination endurance and resistance, HIIT=high intensity interval training, BC=behavioral component, MBS=metabolic and bariatric surgery, BMI=body mass index, WL=weight loss, FM=fat mass, FFM=fat-free mass, LBM=lean body mass, BF=body fat, BW=body weight, 1RM=1 rep maximum, Ex=exercise group. C=control group, 6MWTD=6 minute walking test difference, MVPA=moderate to vigorous physical activity, WC=waist circumference, HC=hip circumference, QoL=quality of life, SBP=systolic blood pressure, DBP=diastolic blood pressure, RHR=resting heart rate, HDL-C=high density lipoprotein cholesterol, LDL=low density lipoprotein cholesterol, TC=total cholesterol, TG=triglycerides, HOMA-IR=homeostatic model assessment for insulin resistance, SPISE=single-point insulin sensitivity estimator, AIRg=acute insulin response, Di=disposition index, HbA1C=hemoglobin A1C, ES=Hedge's *g* effect size, MD=mean difference, WMD=weighted mean difference, SMD=standardized mean difference, NR=not reported, N/A=not applicable.

^a First country listed in first author's affiliations plus number of additional countries in all authors' reported affiliations; ^b Values reported from details provided in tables include minimum and maximum or mean/median age and BMI, and % women; ^c only outcomes with a synthesis within the meta-analysis or in the systematic review text are reported; ^d For meta-analyses, summary includes effect size estimate, 95% confidence interval, *I*² = values of heterogeneity, sample size (*n*) and number of studies (*k*) included in the analysis. For systematic reviews, NS indicates no significant difference and + indicate a significant improvement following intervention compared to control; ^e RoB = Cochrane risk of bias tool, RoB2 = Cochrane risk of bias tool 2, PEDro = Physiotherapy evidence database, ROBINS-I = Risk of bias in non randomized studies of interventions; ^f *k*=17 in systematic review; ^g *k*=20 in systematic review including 4, 3, 2, 2 and 2 publications with the same interventions; ^h *k*=26 in systematic review.

Appendix K – Post-MBS Primary Articles

Read second table as if continuously scrolling to the right of the first table

Table S15: Post-MBS Primary Articles

All Articles	Bond et al 2023	Baillet et al 2022	Roth et al 2022	Vieira et al 2022	Gasmi et al 2022	Diniz- Sousa et al 2022	Boppre et al 2022	Boppre et al 2021	Bellicha et al 2021	Carretero- Ruiz et al 2021	Marshall et al 2020
Auclair et al 2021							X			X	
Campanha-Versiani et al 2017		X	X	X		X			X		
Carnero et al 2017											
Casali et al 2011											
Castello et al 2011		X	X				X	X	X	X	X
Castello et al 2013					X						
Coen et al 2015a							X			X	X
Coen et al 2015b		X					X	X	X	X	X
Coleman et al 2017	X	X		X				X	X		
Daniels et al 2018		X		X				X	X		X
Dantas et al 2020							X			X	
de Oliveira et al 2021				X							
Diniz-sousa et al 2021		X		X		X					
Freitas et al 2017					X						
Galle et al 2020				X							
Gil et al 2021				X							
Hassannejad et al 2017		X	X	X	X			X	X		
Herring et al 2017	X	X		X	X		X	X	X	X	
Huck 2015		X		X					X	X	X
Kelley 2019				X							
Lamarca et al 2021		X		X							
Marc-Hernandez et al 2020	X	X			X						
Marchesi et al 2015		X							X	X	

Mundberg et al 2018a		X			X		X	X	X	X	X
Mundberg et al 2018b	X			X	X		X			X	X
Murai et al 2019		X	X			X					
Muschitz et al 2016		X	X			X					
Noack-Segovia et al 2019				X							
Nunez Lopez et al 2017							X				X
Oliveira et al 2016											
Onofre et al 2017		X							X	X	X
Oppert et al 2018		X	X		X			X	X		
Ricci et al 2020					X						
Rojhani-Shirazi et al 2016		X									
Sellberg et al 2019					X						
Shah et al 2011	X	X					X	X	X	X	
Stegen et al 2011		X		X					X	X	X
Stolberg et al 2018a											X
Stolberg et al 2018b											
Stolberg et al 2018c					X						X
Tardif et al 2020		X					X	X		X	
Woodlief et al 2015											X

Note. Black=primary studies published after systematic review/meta-analysis and therefore not possible for inclusion.

Grey=primary articles published the same year as the systematic review/meta-analysis and therefore not likely for inclusion.

ALL ARTICLES	Carretero-Ruiz et al 2019	da Silva et al 2019	Ren et al 2018	Schurmans et al 2022	Morales-Marroquin et al 2020	Civi Karaaslan et al 2020	Pouwels et al 2015	Baillot et al 2014	Fonseca-Junior et al 2013	Total Count
Auclair et al 2021										2
Campanha-Versiani et al 2017	X				X					7
Carnero et al 2017				X						1
Casali et al 2011	X									1

Castello et al 2011	X		X	X		X		X	X	13
Castello et al 2013				X		X	X			4
Coen et al 2015a	X	X		X						6
Coen et al 2015b			X	X						8
Coleman et al 2017	X		X	X		X				9
Daniels et al 2018	X		X	X	X	X				10
Dantas et al 2020										2
de Oliveira et al 2021										1
Diniz-sousa et al 2021										3
Freitas et al 2017										1
Galle et al 2020										1
Gil et al 2021										1
Hassannejad et al 2017	X		X	X	X					10
Herring et al 2017	X		X	X	X					12
Huck 2015	X	X		X	X					9
Kelley 2019										1
Lamarca et al 2021										2
Marc-Hernandez et al 2020										3
Marchesi et al 2015	X	X								5
Mundberg et al 2018a	X		X	X	X					11
Mundberg et al 2018b						X				7
Murai et al 2019					X					4
Muschitz et al 2016										3
Noack-Segovia et al 2019										1
Nunez Lopez et al 2017		X		X						4
Oliveira et al 2016	X					X				2
Onofre et al 2017	X	X								6
Oppert et al 2018					X					6
Ricci et al 2020										1
Rojhani-Shirazi et al 2016	X									2

Sellberg et al 2019										1
Shah et al 2011	X	X	X	X		X	X	X	X	14
Stegen et al 2011	X	X			X		X	X	X	11
Stolberg et al 2018a				X						2
Stolberg et al 2018b				X						1
Stolberg et al 2018c										2
Tardif et al 2020										4
Woodlief et al 2015				X						2

Note. Black=primary studies published after systematic review/meta-analysis and therefore not possible for inclusion.

Grey=primary articles published the same year as the systematic review/meta-analysis and therefore not likely for inclusion.

1. Auclair A, Harvey J, Leclerc J, et al. Determinants of cardiorespiratory fitness after bariatric surgery: Insights from a randomised controlled trial of a supervised training program. *Can J Cardiol.* 2021;37(2):251-259. doi:10.1016/j.cjca.2020.03.032
2. Campanha-Versiani L, Pereira DAG, Ribeiro-Samora GA, et al. The effect of a muscle weight-bearing and aerobic exercise program on the body composition, muscular strength, biochemical markers, and bone mass of obese patients who have undergone gastric bypass surgery. *Obes Surg.* 2017;27(8):2129-2137. doi:10.1007/s11695-017-2618-5
3. Carnero EA, Dubis GS, Hames KC, et al. Randomized trial reveals that physical activity and energy expenditure are associated with weight and body composition after RYGB. *Obesity.* 2017;25(7):1206-1216. doi:10.1002/oby.21864
4. Casali CCC, Pereira APM, Martinez JAB, de Souza HCD, Gastaldi AC. Effects of inspiratory muscle training on muscular and pulmonary function after bariatric surgery in obese patients. *Obes Surg.* 2011;21(9):1389-1394. doi:10.1007/s11695-010-0349-y
5. Castello V, Simões RP, Bassi D, Catai AM, Arena R, Borghi-Silva A. Impact of aerobic exercise training on heart rate variability and functional capacity in obese women after gastric bypass surgery. *Obes Surg.* 2011;21(11):1739-1749. doi:10.1007/s11695-010-0319-4
6. Castello-Simões V, Polaquini Simões R, Beltrame T, et al. Effects of aerobic exercise training on variability and heart rate kinetic during submaximal exercise after gastric bypass surgery – A randomized controlled trial. *Disabil Rehabil.* 2013;35(4):334-342. doi:10.3109/09638288.2012.694575
7. Coen(a) PM, Menshikova EV, Distefano G, et al. Exercise and weight loss improve muscle mitochondrial respiration, lipid partitioning, and insulin sensitivity after gastric bypass surgery. *Diabetes.* 2015;64(11):3737-3750. doi:10.2337/db15-0809
8. Coen(b) PM, Tanner CJ, Helbling NL, et al. Clinical trial demonstrates exercise following bariatric surgery improves insulin sensitivity. *J Clin Invest.* 2015;125(1):248-257. doi:10.1172/JCI78016
9. Coleman KJ, Caparosa SL, Nichols JF, et al. Understanding the capacity for exercise in post-bariatric patients. *Obes Surg.* 2017;27(1):51-58. doi:10.1007/s11695-016-2240-y

10. Daniels P, Burns RD, Brusseau TA, et al. Effect of a randomised 12-week resistance training programme on muscular strength, cross-sectional area and muscle quality in women having undergone Roux-en-Y gastric bypass. *J Sports Sci.* 2018;36(5):529-535. doi:10.1080/02640414.2017.1322217
11. Dantas WS, Roschel H, Murai IH, et al. Exercise-induced increases in insulin sensitivity after bariatric surgery are mediated by muscle extracellular matrix remodeling. *Diabetes.* 2020;69(8):1675-1691. doi:10.2337/db19-1180
12. de Oliveira Júnior GN, Goessler KF, Santos JVP, et al. Home-based exercise training during COVID-19 pandemic in post-bariatric patients: A randomized controlled trial. *Obes Surg.* 2021;31(11):5071-5078. doi:10.1007/s11695-021-05621-5
13. Diniz-Sousa F, Veras L, Bopp G, et al. The effect of an exercise intervention program on bone health after bariatric surgery: A randomized controlled trial. *J Bone Miner Res.* 2021;36(3):489-499. doi:10.1002/jbmr.4213
14. Freitas PD, Ferreira PG, Silva AG, et al. The role of exercise in a weight-loss program on clinical control in obese adults with asthma. A randomized controlled trial. *Am J Respir Crit Care Med.* 2017;195(1):32-42. doi:10.1164/rccm.201603-0446OC
15. Gallé F, Marte G, Cirella A, et al. An exercise-based educational and motivational intervention after surgery can improve behaviors, physical fitness and quality of life in bariatric patients. *PLOS ONE.* 2020;15(10):e0241336. doi:10.1371/journal.pone.0241336
16. Gil S, Peçanha T, Dantas WS, et al. Exercise enhances the effect of bariatric surgery in markers of cardiac autonomic function. *Obes Surg.* 2021;31(3):1381-1386. doi:10.1007/s11695-020-05053-7
17. Hassannejad A, Khalaj A, Mansournia MA, Rajabian Tabesh M, Alizadeh Z. The effect of aerobic or aerobic-strength exercise on body composition and functional capacity in patients with BMI ≥ 35 after bariatric surgery: A randomized control trial. *Obes Surg.* 2017;27(11):2792-2801. doi:10.1007/s11695-017-2717-3
18. Herring LY, Stevinson C, Carter P, et al. The effects of supervised exercise training 12–24 months after bariatric surgery on physical function and body composition: A randomised controlled trial. *Int J Obes.* 2017;41(6):909-916. doi:10.1038/ijo.2017.60
19. Huck CJ. Effects of supervised resistance training on fitness and functional strength in patients succeeding bariatric surgery. *J Strength Cond Res.* 2015;29(3):589. doi:10.1519/JSC.0000000000000667
20. Kelley J. Maintaining skeletal muscle through eccentric exercise after bariatric surgery: A randomized controlled trial. Theses Diss. Published online December 1, 2019. <https://scholarsarchive.byu.edu/etd/7742>
21. Lamarca F, Vieira FT, Lima RM, et al. Effects of resistance training with or without protein supplementation on body composition and resting energy expenditure in patients 2–7 years post-Roux-en-Y gastric bypass: a Controlled Clinical Trial. *Obes Surg.* 2021;31(4):1635-1646. doi:10.1007/s11695-020-05172-1
22. Marc-Hernández A, Ruiz-Tovar J, Aracil A, Guillén S, Moya-Ramón M. Effects of a high-intensity exercise program on weight regain and cardio-metabolic profile after 3 years of bariatric surgery: A randomized trial. *Sci Rep.* 2020;10(1):3123. doi:10.1038/s41598-020-60044-z
23. Marchesi F, De Sario G, Reggiani V, et al. Road running after gastric bypass for morbid obesity: Rationale and results of a new protocol. *Obes Surg.* 2015;25(7):1162-1170. doi:10.1007/s11695-014-1517-2
24. Mundbjerg(a) LH, Stolberg CR, Cecere S, et al. Supervised physical training improves weight loss after Roux-en-Y gastric bypass surgery: A randomized controlled trial. *Obesity.* 2018;26(5):828-837. doi:10.1002/oby.22143

25. Mundbjerg(b) LH, Stolberg CR, Bladbjerg EM, Funch-Jensen P, Juhl CB, Gram B. Effects of 6 months supervised physical training on muscle strength and aerobic capacity in patients undergoing Roux-en-Y gastric bypass surgery: A randomized controlled trial. *Clin Obes.* 2018;8(4):227-235. doi:10.1111/cob.12256
26. Murai IH, Roschel H, Dantas WS, et al. Exercise mitigates bone loss in women with severe obesity after Roux-en-Y gastric bypass: A randomized controlled trial. *J Clin Endocrinol Metab.* 2019;104(10):4639-4650. doi:10.1210/je.2019-00074
27. Muschitz C, Kocijan R, Haschka J, et al. The impact of vitamin d, calcium, protein supplementation, and physical exercise on bone metabolism after bariatric surgery: The BABS study. *J Bone Miner Res.* 2016;31(3):672-682. doi:10.1002/jbmr.2707
28. Noack-Segovia JP, Sánchez-López AM, García-García I, Rodríguez-Blanco R, León-Ríos XA, Aguilar-Cordero MJ. Physical exercise and grip strength in patients intervened through bariatric surgery. *AQUICHAN.* 2019;19(3). doi:10.5294/aqui.2019.19.3.6
29. Nunez Lopez YO, Coen PM, Goodpaster BH, Seyhan AA. Gastric bypass surgery with exercise alters plasma microRNAs that predict improvements in cardiometabolic risk. *Int J Obes.* 2017;41(7):1121-1130. doi:10.1038/ijo.2017.84
30. Oliveira JJJ de, Freitas ACT de, Almeida AA de. Postoperative effect of physical therapy related to functional capacity and respiratory muscle strength in patients submitted to bariatric surgery. *Arq Bras Cir Dig.* 2016;29Suppl 1(Suppl 1):43-47. doi:10.1590/0102-6720201600s10012
31. Onofre T, Carlos R, Oliver N, et al. Effects of a physical activity program on cardiorespiratory fitness and pulmonary function in obese women after bariatric surgery: A pilot study. *Obes Surg.* 2017;27(8):2026-2033. doi:10.1007/s11695-017-2584-y
32. Oppert JM, Bellicha A, Roda C, et al. Resistance training and protein supplementation increase strength after bariatric surgery: A randomized controlled trial. *Obesity.* 2018;26(11):1709-1720. doi:10.1002/oby.22317
33. Ricci PA, Di Thommazo-Luporini L, Jürgensen SP, et al. Effects of Whole-Body Electromyostimulation Associated with Dynamic Exercise on Functional Capacity and Heart Rate Variability After Bariatric Surgery: A randomized, double-blind, and sham-controlled trial. *Obes Surg.* 2020;30(10):3862-3871. doi:10.1007/s11695-020-04724-9
34. Rojhani-Shirazi Z, Azadeh Mansoriyan S, Hosseini SV. The effect of balance training on clinical balance performance in obese patients aged 20–50 years old undergoing sleeve gastrectomy. *Eur Surg.* 2016;48(2):105-109. doi:10.1007/s10353-015-0379-8
35. Sellberg F, Possmark S, Willmer M, Tynelius P, Persson M, Berglund D. Meeting physical activity recommendations is associated with health-related quality of life in women before and after Roux-en-Y gastric bypass surgery. *Qual Life Res.* 2019;28(6):1497-1507. doi:10.1007/s11136-019-02120-0
36. Shah M, Snell PG, Rao S, et al. High-volume exercise program in obese bariatric surgery patients: A randomized, controlled trial. *Obesity.* 2011;19(9):1826-1834. doi:10.1038/oby.2011.172
37. Stegen S, Derave W, Calders P, Van Laethem C, Pattyn P. Physical fitness in morbidly obese patients: Effect of gastric bypass surgery and exercise training. *Obes Surg.* 2011;21(1):61-70. doi:10.1007/s11695-009-0045-y
38. Stolberg(a) CR, Mundbjerg LH, Funch-Jensen P, Gram B, Juhl CB, Bladbjerg EM. Effects of gastric bypass followed by a randomized study of physical training on markers of coagulation activation, fibrin clot properties, and fibrinolysis. *Surg Obes Relat Dis.* 2018;14(7):918-926. doi:10.1016/j.soard.2018.03.022

39. Stolberg(b) CR, Mundbjerg LH, Bladbjerg EM, Funch-Jensen P, Gram B, Juhl CB. Physical training following gastric bypass: Effects on physical activity and quality of life - A randomized controlled trial. *Qual Life Res.* 2018;27(12):3113-3122. doi:10.1007/s11136-018-1938-9
40. Stolberg(c) CR, Mundbjerg LH, Funch-Jensen P, Gram B, Bladbjerg EM, Juhl CB. Effects of gastric bypass surgery followed by supervised physical training on inflammation and endothelial function: A randomized controlled trial. *Atherosclerosis.* 2018;273:37-44. doi:10.1016/j.atherosclerosis.2018.04.002
41. Tardif I, Auclair A, Piché ME, et al. Impact of a 12-week randomized exercise training program on lipid profile in severely obese patients following bariatric surgery. *Obes Surg.* 2020;30(8):3030-3036. doi:10.1007/s11695-020-04647-5
42. Woodlief TL, Carnero EA, Standley RA, et al. Dose response of exercise training following Roux-en-Y gastric bypass surgery: A randomized trial. *Obesity.* 2015;23(12):2454-2461. doi:10.1002/oby.21332

Appendix L – Post-MBS Subanalyses

Table S16: Post-MBS Exercise Interventions: Systematic Reviews/Meta-Analyses Sub Analysis Results, Considerations and Conclusions

Author (Year)	Number of Primary Studies Included	Outcomes Assessed ^a	Main Results ^b <i>Level of Evidence</i>	Special Considerations	AMSTAR2	Conclusions						
Intervention Type												
Boppre 2022 k=11	1. VO ₂ max 2. SBP 3. DBP 4. Insulin 5. Glucose 6. HbA1C 7. TC 8. HDL-C 9. LDL-C 10. TG	<u>Endurance</u> 1. MD: 0.75 L/min [-1.06; 2.56], I ² =49%; n=NR, k=2 2. MD: -3.55 mmHg [-8.68; 1.58], I ² =0%; n=NR, k=3 3. MD: -0.89 mmHg [-5.61; 3.82], I ² =0%; n=NR, k=3 4. MD: 0.69 μIU/mL [-0.62; 2.01], I ² =0%; n=NR, k=2 5. MD: 1.13 mg/dL [-3.66; 5.93], I ² =0%; n=NR, k=2 7. MD: -2.14 mg/dL [-15.37; 11.09], I ² =0%; n=NR, k=2 8. MD: 0.22 mg/dL [-4.48; 4.92], I ² =0%; n=NR, k=2 9. MD: -1.79 mg/dL [-12.49; 8.90], I ² =0%; n=NR, k=2 10. MD: -0.13 mg/dL [-15.87; 15.61], I ² =0%; n=NR, k=2			Only RCT	Critically Low	Combined E/R = + SBP and TG					
		<u>Combined Endurance/Resistance</u> 2. MD: -7.18 mmHg [-12.42; -1.94], I ² =0%; n=137, k=3 (moderate certainty evidence) 3. MD: -4.01 mmHg [-9.49; 1.46], I ² =53%; n=NR, k=3 4. MD: -3.36 μIU/mL [-7.03; 0.31], I ² =45%; n=NR, k=2 5. MD: 0.23 mg/dL [-8.95; 9.42], I ² =0%; n=NR, k=2 6. MD: -0.65 mmol/mol [-2.22; 0.93], I ² =0%; n=NR, k=2 7. MD: -3.88 mg/dL [-16.04; 8.27], I ² =0%; n=NR, k=3 8. MD: -11.77 mg/dL [-30.17; 6.64], I ² =67%; n=NR, k=3 9. MD: 1.36 mg/dL [-5.18; 7.91], I ² =59%; n=NR, k=3 10. MD: -17.56 mg/dL [-34.15; -0.96], I ² =0%; n=171, k=3 (low certainty evidence)										
		<u>Endurance</u> 1. MD: -0.80 kg [-7.19; 5.58], I ² =0%; n=214, k=4 2. MD: -0.21 kg/m ² [-2.42; 2.00], I ² =0%; n=186, k=3 4. MD: -0.10 kg [-3.61; 3.41], I ² =0%; n=156, k=2 5. MD: -4.30 cm [-11.30; 2.70], I ² =39%; n=177, k=3										
		<u>Resistance</u> 1. MD: 0.46 kg [-3.24; 4.16], I ² =0%; n=61, k=2										
		<u>Combined E/R</u> 1. MD: -5.02 kg [-8.13; -1.90], I ² =0%; n=221, k=5 2. MD: -1.62 kg/m ² [-2.72; -0.59], I ² =0%; n=170, k=4										
		Boppre 2021 k=10	1. BW 2. BMI 3. FM 4. LBM 5. WC					Only RCT	Critically Low	Combined E/R = + BW and BMI		

		<u>Endurance</u> Pooled SMD: 0.21 [-0.07; 0.49], I ² =0%; n=NR, k=5			
		<u>Resistance</u> Pooled SMD: 0.15 [-0.56; 0.85], I ² =0%; n=NR, k=2			
		<u>Combined E/R</u> Pooled SMD: 0.19 [-0.09; 0.46], I ² =0%; n=NR, k=7			
Carretero-Ruiz 2019 k=16	WL	<u>Alternative</u> Pooled SMD: -0.08 [-0.48; 0.33], I ² =0%; n=NR, k=3	RCT, NRCT Respiratory (k=1), and physiotherapy (k=1) interventions	Low	NS differences
		<u>Programmed</u> Pooled SMD: 0.15 [-0.23; 0.54], I ² =0%; n=NR, k=3			
		<u>Supervised</u> Pooled SMD: 0.10 [-0.16; 0.36], I ² =0%; n=NR, k=9			
		<u>Combined Programmed/Supervised</u> Pooled SMD: 0.20 [-0.09; 0.50], I ² =0%; n=NR, k=4			
Ren 2018 k=8	BW	<u>Endurance</u> WMD: -0.24 kg [-1.56; 1.09], I ² =0%; n=NR, k=3 <u>Resistance</u> WMD: -2.20 kg [-5.35; 0.95], I ² =N/A, n=NR, k=1 <u>Combined E/R</u> WMD: -3.12 kg [-4.56; -1.68], I ² =32%; n=NR, k=4	Only RCT	Low	Combined E/R = + BW
Intervention Start Time					
		<u><6 months</u> 1. MD: -3.30 mmHg [-8.29; 1.69], I ² =0%; n=NR, k=4 2. MD: -0.08 [-3.25; 3.09], I ² =0%; n=NR, k=4 4. MD: -1.84 µIU/mL [-6.89; 3.20], I ² =80%; n=NR, k=3 5. MD: 0.77 mg/dL [-3.87; 5.42], I ² =0%; n=NR, k=3 6. MD: -1.40 mg/dL [-11.85; 9.05], I ² =0%; n=NR, k=4 7. MD: 2.10 mg/dL [-1.58; 5.78], I ² =0%; n=NR, k=4 8. MD: -1.84 mg/dL [-10.48; 6.80], I ² =0%; n=NR, k=4 9. MD: -4.51 mg/dL [-17.09; 8.07], I ² =%; n=NR, k=4			
Boppre 2022 k=11	1. SBP 2. DBP 3. RHR 4. Insulin 5. Glucose 6. TC 7. HDL-C 8. LDL-C 9. TG	<u>>6 months</u> 1. MD: -7.71 mmHg [-13.12; -2.31], I ² =0%; n=84, k=2 (high certainty evidence) 2. MD: -5.20 mmHg [-11.72; 1.33], I ² =%; n=NR, k=2 3. MD: -0.08 bpm [-3.25; 3.09], I ² =%; n=NR, k=4	Only RCT	Critically Low	<u>>6 months</u> = + SBP

Boppre 2021 k=10	1. BW 2. BMI 3. FM 4. LBM 5. WC	<u><6 months</u> 1. MD: -0.12 kg [-3.17; 2.93], I ² =0%; n=361, k=7 (8 arms) 2. MD: -0.16 kg/m ² [-1.15; 0.82], I ² =0%; n=317, k=5 (6 arms) 3. MD: 0.49 kg [-1.71; 2.69], I ² =0%; n=173, k=2 4. MD: 0.87 kg [-0.65; 2.40], I ² =0%; n=201, k=3 5. MD: -4.30 cm [-11.30; 2.70], I ² =39%; n=177, k=3	Only RCT	Critically Low	<u>>6 months</u> = + BW and BMI
		<u>>6 months</u> 1. MD: -5.25 kg [-8.52; -1.97], I ² =0%; n=135, k=3 2. MD: -1.84 kg/m ² [-3.04; -0.64], I ² =0%; n=84, k=2			
Carretero-Ruiz 2019 k=16	WL	<u><3 months postMBS</u> Pooled SMD: 0.12 [-0.08; 0.33], I ² =0%; n=NR, k=11 <u>>3 months posMBS</u> Pooled SMD: 0.20 [-0.13; 0.53], I ² =0%; n=NR, k=5	RCT, NRCT Respiratory (k=1), and physiotherapy (k=1) interventions	Low	NS differences
Ren 2018 k=8	BW	<u><6 months</u> WMD: -1.71 kg [-3.06; -0.36], I ² =32%; n=NR, k=5 <u>>12 months</u> WMD: -3.63 kg [-5.35; -1.91], I ² =0%; n=NR, k=2 <u>Varying starting times</u> WMD: 0.50 kg [-1.79; 2.79], I ² =N/A; n=NR, k=1	Only RCT	Low	<u><6 months and >12 months</u> = + BW *significantly greater effects for >12 months compared to <6 months
Intervention Duration					
Boppre 2022 k=11	1. VO ₂ max 2. SBP 3. DBP 4. RHR 5. Insulin 6. Glucose 7. HOMA-IR 8. HbA1C 9. TC 10. HDL-C 11. LDL-C 12. TG	<u><12 weeks</u> 2. MD: -3.61 mmHg [-11.61; 4.39], I ² =0%; n=NR, k=3 3. MD: -2.97 mmHg [-8.01; 2.08], I ² =0%; n=NR, k=3 9. MD: 1.96 mg/dL [-19.93; 23.85], I ² =0%; n=NR, k=2 10. MD: 4.39 mg/dL [-2.58; 11.36], I ² =0%; n=NR, k=2 11. MD: -1.10 mg/dL [-20.00; 17.80], I ² =0%; n=NR, k=2 12. MD: -7.30 mg/dL [-34.65; 20.04], I ² =0%; n=NR, k=2	Only RCT	Critically Low	<u>>12 weeks</u> = + SBP
		<u>>12 weeks</u> 1. MD: 0.23 L/min [-0.15; 0.60], I ² =0%; n=NR, k=2 2. MD: -5.78 mmHg [-9.91; -1.66], I ² =0%; n=212, k=3 (high certainty evidence) 3. MD: -2.51 mmHg [-9.22; 4.19], I ² =82%; n=NR, k=3 4. MD: -2.16 bpm [-6.98; 2.66], I ² =0%; n=NR, k=2 5. MD: -1.61 µIU/mL [-5.41; 2.19], I ² =81%; n=NR, k=3 6. MD: 0.91 mg/dL [-3.35; 5.18], I ² =0%; n=NR, k=3 7. MD: 1.39 [-1.30; 4.08], I ² =89%; n=NR, k=2 8. MD: -0.65 mmol/mol [-2.22; 0.93], I ² =0%; n=NR, k=2 9. MD: -4.10 mg/dL [-13.91; 5.71], I ² =0%; n=NR, k=3 10. MD: -0.41 mg/dL [-5.08; 4.26], I ² =45%; n=NR, k=3 11. MD: -10.66 mg/dL [-27.69; 6.36], I ² =76%; n=NR, k=3 12. MD: -10.23 mg/dL [-26.02; 5.56], I ² =31%; n=NR, k=3			

Boppre 2021 k=10	1. BW 2. BMI 3. FM 4. LBM 5. WC	<u>≤12 weeks</u> 1. MD: - 1.68 kg [-7.68; 4.32], I ² =0%; n=188, k=5 (6 arms) 2. MD: -0.40 kg/m ² [-2.54; 1.74], I ² =0%; n=144, k=3 (4 arms) 5. MD: -6.51 cm [-14.30; 1.29], I ² =N/A; n=74, k=1 (2 arms)	Only RCT	Critically Low	NS differences
		<u>>12 weeks</u> 1. MD: - 2.28 kg [-6.31; 1.75], I ² =48%; n=257, k=4 2. MD: - 0.85 kg/m ² [-2.05; 0.35], I ² =40%; n=257, k=4 3. MD: -1.40 kg [-4.84; 2.03], I ² =68%; n=197, k=3 4. MD: 0.92 kg [-0.65; 2.50], I ² =0%; n=173, k=2 5. MD: -3.38cm [-7.66; 0.90], I ² =0%; n=152, k=2			
Carretero-Ruiz 2019 k=16	WL	<u>≤16 weeks</u> Pooled SMD: 0.06 [-0.17; 0.29], I ² =0%; n=NR, k=12 <u>>16 weeks</u> Pooled SMD: 0.25 [-0.01; 0.50], I ² =0%; n=NR, k=5	RCT, NRCT Respiratory (k=1), and physiotherapy (k=1) interventions	Low	NS differences
Intervention Exercise Time/Week					
Carretero-Ruiz 2019 k=16	WL	<u>≤150 min/week</u> Pooled SMD: 0.17 [-0.11; 0.44], I ² =0%; n=NR, k=4 <u>>150 min/week</u> Pooled SMD: 0.13 [-0.08; 0.35], I ² =0%; n=NR, k=13	RCT, NRCT Respiratory (k=1), and physiotherapy (k=1) interventions	Low	NS differences

Note. All data were reported as they were presented in the manuscript. Red text represents significant findings. MBS=metabolic and bariatric surgery, RCT=randomized control trial, NCRT=non randomized control trial, E=endurance, R=resistance, E/R=combination endurance and resistance, BMI=body mass index, WL=weight loss, FM=fat mass, LBM=lean body mass, BW=body weight, WC=waist circumference, SBP=systolic blood pressure, DBP=diastolic blood pressure, HDL-C=high density lipoprotein cholesterol, LDL=low density lipoprotein cholesterol, TC=total cholesterol, TG=triglycerides, HOMA-IR=homeostatic model assessment for insulin resistance, HbA1C=hemoglobin A1C, MD=mean difference, WMD=weighted mean difference, SMD=standardized mean difference, NR=not reported.

^a only outcomes with a synthesis within the meta-analysis or in the systematic review text are reported; ^b For meta-analyses, summary includes effect size estimate, 95% confidence interval, I² = values of heterogeneity, sample size (n) and number of studies (k) included in the analysis.

Appendix M – Feasibility and Acceptability Outcomes

Table S17: Summary of Baillot et al., 2022 Feasibility and Acceptability Outcomes

Effects	<i>n</i>	<i>k</i>	arms	I ²	Studies included	Study Design
Total Attendance Rate (%; Exercise Training)						
Pre MBS pooled percentage: 79.4% [67.7; 89.4]	NR	3	4	0%	Baillot 2016, Marcon 2017, Picó-Servant 2019	RCT only
Post MBS pooled percentage: 87.4% [76.7; 95.6]	NR	5	6	0%	Castello 2011, Herring 2017, Huck 2015, Lamarca 2021, Murai 2019	RCT/NRCT
Total Refusal Rate (%; Exercise Training and Control Group)						
Pre MBS pooled percentage: 30.7% [0.0%; 81.0%]	NR	4	4	92%	Arman 2021, Baillot 2016, Gilbertson 2020, Marcon 2017	RCT/NRCT
Post MBS pooled percentage: 20.3% [6.5%; 38.7%]	NR	12	12	95%	Campanha-Versiani 2017, Castello 2011, Diniz Souza 2020, Hassanejad 2017, Herring 2017, Lamarca 2021, Marc-Hernandez 2020, Mundberg 2018a, Murai 2019, Onofre 2017, Oppert 2018, Tardif 2020	RCT/NRCT
Total Recruitment Rate (n/month; Exercise Training and Control Group)						
Pre MBS pooled percentage: 8.1% [0.7; 20.2]	NR	3	3	0%	Arman 2021, Baillot 2016, Gilbertson 2020	RCT/NRCT
Post MBS pooled percentage: 7.0% [2.9; 12.4]	NR	9	9	70%	Castello 2011, Coen 2015b, Diniz Souza 2020, Hassanejad 2017, Herring 2017, Lamarca 2021, Mundberg 2018a, Oppert 2018, Tardif 2020	RCT/NRCT
Total Enrollment Speed (n/month; Exercise Training and Control Group)						
Pre MBS pooled percentage: 1.1% [0.0; 6.2]	NR	3	3	0%	Arman 2021, Baillot 2016, Gilbertson 2020	RCT/NRCT
Post MBS pooled percentage: 2.9% [1.6; 4.4]	NR	10	10	0%	Castello 2011, Coen 2015b, Diniz Souza 2020, Hassanejad 2017, Herring 2017, Lamarca 2021, Marc-Hernandez 2020, Mundberg 2018a, Oppert 2018, Tardif 2020	RCT/NRCT
Total Enrollment Rate (%; Exercise Training and Control Group)						
Pre MBS pooled percentage: 17.4% [0.4; 46.5]	NR	4	4	75%	Arman 2021, Baillot 2016, Gilbertson 2020, Marcon 2017	RCT/NRCT
Post MBS pooled percentage: 50.8% [36.4; 65.2]	NR	14	14	93%	Campanha-Versiani 2017, Castello 2011, Coen 2015b, Coleman 2017, Diniz Souza 2020, Hassanejad 2017, Herring 2017, Lamarca 2021, Marc-Hernandez 2020, Mundberg 2018a, Murai 2019, Onofre 2017, Oppert 2018, Tardif 2020	RCT/NRCT

Total Dropout Rate (%; Pre MBS)							
Exercise pooled percentage: 3.6% [0.0; 14.6]	NR	6	7	44%	Arman 2021, Baillot 2016, Gilbertson 2020, Marc-Hernandez 2019, Marcon 2017, Picó-Servant 2019	RCT/NRCT	
Control pooled percentage: 0.0% [0.0; 0.3]	NR	4	4	0%	Baillot 2016, Gilbertson 2020, Marcon 2017, Picó-Servant 2019	RCT/NRCT	
Total Dropout Rate (%; Post MBS)							
Exercise pooled percentage: 5.6% [0.6; 13.8]	NR	12	12	69%	Castello 2011, Coen 2015b, Coleman 2017, Daniels 2017, Herring 2017, Marc-Hernandez 2020, Marchesi 2015, Murai 2019, Onofre 2017, Oppert 2018, Shah 2011, Tardif 2020	RCT/NRCT	
Control pooled percentage: 2.8% [0.0; 8.2]	NR	13	14	62%	Castello 2011, Coen 2015b, Coleman 2017, Daniels 2017, Herring 2017, Marc-Hernandez 2020, Marchesi 2015, Murai 2019, Muschitz 2016, Onofre 2017, Oppert 2018, Shah 2011, Tardif 2020	RCT/NRCT	
Total Retention Rate (%; Pre MBS)							
Exercise pooled percentage: 96.4% [84.6; 100]	NR	6	8	50%	Arman 2021, Baillot 2016, Gilbertson 2020, Marc-Hernandez 2019, Marcon 2017, Picó-Servant 2019	RCT/NRCT	
Control pooled percentage: 89.5% [77.8; 97.8]	NR	6	6	0%	Arman 2021, Baillot 2016, Gilbertson 2020, Marc-Hernandez 2019, Marcon 2017, Picó-Servant 2019	RCT/NRCT	
Total Retention Rate (%; Post MBS)							
Exercise pooled percentage: 83.6% [74.1; 91.5]	NR	17	18	83%	Campanha-Versiani 2017, Castello 2011, Coen 2015b, Coleman 2017, Daniels 2017, Herring 2017, Lamarca 2021, Marc-Hernandez 2020, Marchesi 2015, Mundberg 2018a, Murai 2019, Muschitz 2016, Onofre 2017, Oppert 2018, Shah 2011, Stegen 2011, Tardif 2020	RCT/NRCT	
Control pooled percentage: 85.6% [78.1; 92.1]	NR	18	20	73%	Campanha-Versiani 2017, Castello 2011, Coen 2015b, Coleman 2017, Daniels 2017, Diniz Souza 2020, Herring 2017, Lamarca 2021, Marc-Hernandez 2020, Marchesi 2015, Mundberg 2018a, Murai 2019, Muschitz 2016, Onofre 2017, Oppert 2018, Shah 2011, Stegen 2011, Tardif 2020	RCT/NRCT	

Note. All data were reported as they were presented in the manuscript. MBS=metabolic and bariatric surgery, n =sample size, k =number of studies, I^2 =measure of heterogeneity, NR=not reported, RCT=randomized control trial, NRCT=non randomized control trial.

Appendix N – Authors Conclusions for Publication Bias by Outcome

Table S18: Pre-MBS Outcomes

Systematic Review/ Meta-Analysis	Author Comments on Publication Bias
Body Weight (BW), Body Mass Index (BMI) and Weight Loss (WL)	
Jabbour 2022	None
Lodewijks 2022	None
Durey 2022	None
Herrera-Santelices 2022	Very low level of evidence with serious risk of bias but no reported concerns about publication bias
Schurmans 2022	None
Bellicha 2021	None
Fat Mass (FM)	
Jabbour 2022	None
Lodewijks 2022	None
Herrera-Santelices 2022	Moderate level of evidence with serious risk of bias but no reported concerns about publication bias
Bellicha 2021	None
Fat-Free Mass (FFM) and Lean Body Mass (LBM)	
Lodewijks 2022	None
Herrera-Santelices 2022	Moderate level of evidence with serious risk of bias but no reported concerns about publication bias
Schurmans 2022	None
Bellicha 2021	None
VO₂max/Maximum Aerobic Capacity	
Durey 2022	None
Jabbour 2022	None
Bellicha 2021	None
6-Minute Walking Test Distance (6MWT)	
Jabbour 2022	None
Herrera-Santelices 2022	High level of evidence with not serious risk of bias and no reported concerns about publication bias
Schurmans 2022	None
Bellicha 2021	None
Muscle Strength and Functional Capacity	
Jabbour 2022	None
Bellicha 2021	None
Resting Heart Rate (RHR)	
Schurmans 2022	None
Marshall 2020	Pooled pre and postoperative results: Very low level of evidence with serious risk of bias but no reported concerns about publication bias
Blood Pressure (BP)	
Schurmans 2022	None
Jabbour 2022	None
Bellicha 2021	None
Marshall 2020	For systolic and diastolic blood pressure, pooled pre and postoperative results: Very low level of evidence with serious risk of bias but no reported concerns about publication bias

Quality of Life (QoL)	
Herrera-Santelices 2022	Moderate level of evidence with serious risk of bias but no reported concerns about publication bias
Lodewijks 2022	None
Schurmans 2022	None
Bellicha 2021	None
Glucose and Lipid Metabolism	
Jabbour 2022	None
Bellicha 2021	None
Physical Activity	
Lodewijks 2022	None
Bellicha 2021	None
Adverse Events	
Durey 2022	None
Length of Hospital Stay	
Durey 2022	None
Jabbour 2022	None

Table S19: Post-MBS Outcomes

Systematic Review/ Meta-analysis	Author Comments on Publication Bias
Weight Loss (WL) \geq 12 Months Post MBS	
Bond 2023	Based on the tests introduced by Begg and Mazumdar ($P = .81$) and Egger et al. ($P = .52$), we did not observe any publication or small sample bias, whereas the funnel plots suggested publication or other reporting bias
Body Weight (BW) and Body Mass Index (BMI)	
Gasmi 2022	For the analysis on BMI and pooled BMI, the funnel plots showed a few outliers, but in both directions, suggesting true heterogeneity rather than publication bias
Schurmans 2022	None
Boppre 2021	There was no significant publication bias as demonstrated by the funnel plot symmetry and the Egger's test result adjusted to body weight. Bias coefficient is -3.00 (intercept) and p-value is higher ($p = 0.708$)
Bellicha 2021	Visual inspection of the funnel plot suggested little evidence of publication bias, which was suggested by Egger's test ($P = 0.22$)
Morales-Marroquin 2020	None
Carretero-Ruiz 2019	As seen in the funnel plot and once the Egger test was performed, there was no evidence of significant publication bias risk ($p = 0.208$) The funnel plot did not suggest publication bias for physical exercise with respect to body weight and the p value for publication bias was 0.44.
Ren 2018	For both body weight and BMI: Moderate level of evidence and not a high risk of publication bias
Waist Circumference (WC)	
Gasmi 2022	Funnel plots displayed a symmetrical appearance with no major outliers despite the small number of included studies

Boppre 2021	There was no significant publication bias as demonstrated by the funnel plot symmetry and the Egger's test result adjusted to body weight. Bias coefficient is -3.00 (intercept) and p-value is higher ($p = 0.708$)
Ren 2018	Low level of evidence and not a high risk of publication bias
Fat Mass (FM)	
Gasmi 2022	Funnel plots displayed a symmetrical appearance with no major outliers despite the small number of included studies
Boppre 2021	There was no significant publication bias as demonstrated by the funnel plot symmetry and the Egger's test result adjusted to body weight. Bias coefficient is -3.00 (intercept) and p-value is higher ($p = 0.708$)
Bellicha 2021	Visual inspection of the funnel plot suggested little evidence of publication bias
Morales-Marroquin 2020	None
Ren 2018	Low level of evidence and not a high risk of publication bias
Fat-Free Mass (FFM) and Lean Body Mass (LBM)	
Roth 2022	For fat-free mass: Very low level of evidence * Potential publication bias could not be assessed using funnel plots or statistical tests, such as Egger's test, because these methods do not possess enough power to distinguish chance from real asymmetry when fewer than 10 studies are involved in a pairwise metaanalysis
Gasmi 2022	For fat-free mass: Funnel plots displayed a symmetrical appearance with no major outliers despite the small number of included studies
Schurmans 2022	None
Boppre 2021	unclear
Bellicha 2021	Visual inspection of the funnel plot suggested little evidence of publication bias, which was suggested by Egger's test ($P = 0.40$ for lean mass loss outcomes)
Morales-Marroquin 2020	None
Ren 2018	For fat-free mass: Very low level of evidence and not a high risk of publication bias
Bone Mineral Density (BMD)	
Roth 2022	Moderate level of evidence * Potential publication bias could not be assessed using funnel plots or statistical tests, such as Egger's test, because these methods do not possess enough power to distinguish chance from real asymmetry when fewer than 10 studies are involved in a pairwise metaanalysis
Diniz-Souza 2022	* Publication bias assessment was not performed because such analysis is not recommended in meta-analysis with less than 10 studies
Bellicha 2021	None
Morales-Marroquin 2020	None
Vo₂max/Peak	
Boppre 2022	* Publication bias assessment was not performed, because outcomes analyses had less than 10 studies included
Schurmans 2022	None
Bellicha 2021	Visual inspection of the funnel plot suggested little evidence of publication bias

Carretero-Ruiz 2021	No evidence of publication bias. In all the funnel plots performed, a symmetrical study distribution was observed, leaving all the studies within the limits of the funnel plot. In all cases, the Egger's statistics were not significant.
da Silva 2019	The p value for Egger's test was 0.25, suggesting no risk of publication bias
6-Minute Walking Test Distance (6MWT)	
Schurmans 2022	None
Bellicha 2021	Visual inspection of the funnel plot suggested little evidence of publication bias
Ren 2018	Low level of evidence and not a high risk of publication bias
Muscle Strength	
Vieira 2022	For 1 rep maximum upper and lower muscle, sit to stand, dynamometer, and handgrip tests of muscle strength: Very low level of evidence * As none of the meta-analyses included more than 10 studies, Egger's test could not be used to assess publication bias. Therefore, we assessed publication bias by evaluating the search strategy and use of industry funding; the results indicated that none of the meta-analyses were affected by publication bias
Schurmans 2022	None
Bellicha 2021	Visual inspection of the funnel plot suggested little evidence of publication bias
Morales-Marroquin 2020	None
Resting Heart Rate (RHR)	
Boppre 2022	* Publication bias assessment was not performed, because outcomes analyses had less than 10 studies included
Schurmans 2022	None
Carretero-Ruiz 2021	No evidence of publication bias. In all the funnel plots performed, a symmetrical study distribution was observed, leaving all the studies within the limits of the funnel plot. In all cases, the Egger's statistics were not significant.
Marshall 2020	Pooled pre and postoperative results: Very low level of evidence with serious risk of bias but no reported concerns about publication bias
Ren 2018	Low level of evidence and high risk of publication bias
Blood Pressure (BP)	
Boppre 2022	* Publication bias assessment was not performed, because outcomes analyses had less than 10 studies included
Schurmans 2022	None
Carretero-Ruiz 2021	No evidence of publication bias. In all the funnel plots performed, a symmetrical study distribution was observed, leaving all the studies within the limits of the funnel plot. In all cases, the Egger's statistics were not significant.
Bellicha 2021	None
Marshall 2020	For systolic and diastolic blood pressure, pooled pre and postoperative results: Very low level of evidence with serious risk of bias but no reported concerns about publication bias
Ren 2018	For systolic blood pressure: Low level of evidence and not a high risk of publication bias For diastolic blood pressure: Very low level of evidence and high risk of publication bias
Quality of Life (QoL)	
Schurmans 2022	None

Bellicha 2021	None
Glucose Metabolism	
Boppre 2022	* Publication bias assessment was not performed, because outcomes analyses had less than 10 studies included
Schurmans 2022	None
Bellicha 2021	None
Marshall 2020	For fasting blood glucose and insulin: Low level of evidence with not serious risk of bias and no reported concerns about publication bias
Lipid Metabolism	
Boppre 2022	* Publication bias assessment was not performed, because outcomes analyses had less than 10 studies included
Schurmans 2022	None
Carretero-Ruiz 2021	No evidence of publication bias. In all the funnel plots performed, a symmetrical study distribution was observed, leaving all the studies within the limits of the funnel plot. In all cases, the Egger's statistics were not significant.
Bellicha 2021	None
Marshall 2020	For triglycerides, high-density lipoprotein, low-density lipoprotein, and total cholesterol: Low level of evidence with not serious risk of bias and no reported concerns about publication bias

Table S20: Post-MBS Sub Analysis Outcomes

Systematic Review/ Meta-analysis	Sub Analysis and Outcome	Author Comments on Publication Bias
Intervention Type		
Boppre 2022	Combined endurance/ resistance on systolic blood pressure	Moderate certainty of evidence * Publication bias assessment was not performed, because outcomes analyses had less than 10 studies included
	Combined endurance/ resistance on triglycerides	Low certainty of evidence * Publication bias assessment was not performed, because outcomes analyses had less than 10 studies included
Boppre 2021	Combined endurance/ resistance on body weight	None
	Combined endurance/ resistance on BMI	None
Ren 2018	Combined endurance/ resistance on body weight	None
Intervention Start Time		
Boppre 2022	> 6 months post MBS on systolic blood pressure	High certainty of evidence * Publication bias assessment was not performed, because outcomes analyses had less than 10 studies included
Boppre 2021	> 6 months post MBS on body weight	None
	> 6 months post MBS on BMI	None
Ren 2018	< 6 months post MBS on body weight	None
	> 12 months post MBS on body weight	None
Intervention Duration		

Boppre 2022	> 12 weeks on systolic blood pressure	High certainty of evidence * Publication bias assessment was not performed, because outcomes analyses had less than 10 studies included
-------------	---------------------------------------	--

Table S21: Baillot et al., 2022: Feasibility and Acceptability Outcomes. Egger's Test of Funnel Plot Asymmetry

Outcome	Author Comments on Publication Bias
Attendance rate (exercise arm)	$t = 0.8555, df = 8, p = 0.4172$
Dropout rate (exercise arm)	$t = 0.3489, df = 17, p = 0.7315$
Enrollment rate (exercise and control arms)	$t = -0.6026, df = 16, p = 0.5552$
Refusal rate (exercise and control arms)	$t = -0.4638, df = 14, p = 0.6499$
Retention rate (exercise arm)	$t = 0.9979, df = 24, p = 0.3283$

Note. The risk of publication bias was examined with funnel plots and tested using the Egger's test ($p < 0.10$ indicating a publication bias). A trim and fill analysis was also carried out to examine the impact of missing studies by adjusting the meta-analysis to take into account the theoretically missing studies. After the trim and fill, no risk of publication bias was found for any of the included outcomes.

Appendix O – Certainty of Evidence Summary (RCT-Only Meta-Analyses)

Table S22: Certainty of Evidence Downgrades (DG) and Final Calculations

Variable	RCT only SRMA	Imprecision		ROB; Trial Quality)		Inconsistency/ Heterogeneity		ROB (Review Quality)		Final COE		SRMA Reported COE
		Sample Size (<i>n</i>)	DG(s) ^a	% Low ROB	DG(s) ^b	I ²	DG(s) ^c	# Critical Absences	DG(s) ^d	Total DGs	Calculated COE	
PRE-MBS ^e												
Body Mass Index	Herrera 2022	115	1	0%	1	76%	1	3	2	5	very low	very low
Weight Loss	Durey 2022	142	1	0%	1	70%	0	4	2	4	low	NR
Fat Mass	Herrera 2022	75	2	0%	1	0%	0	3	2	5	very low	moderate
Fat-Free Mass	Herrera 2022	46	2	0%	1	0%	0	3	2	5	very low	moderate
VO ₂ max (PreMBS change)	Durey 2022	79	2	0%	1	62%	0	4	2	5	very low	NR
VO ₂ max (Change at max follow-up)	Durey 2022	131	1	0%	1	0%	0	4	2	4	low	NR
6-Minute Walking Test Distance	Herrera 2022	61	2	0%	1	0%	0	3	2	5	very low	high
Quality of Life	Herrera 2022	53	2	0%	1	67%	0	3	2	5	very low	moderate
Adverse Events	Durey 2022	22	2	0%	1	n/a	1	4	2	6	very low	NR
Length of Hospital Stay	Durey 2022	22	2	0%	1	n/a	1	4	2	6	very low	NR
POST-MBS ^f												
Weight Loss ≥ 12 months post-MBS	Bond 2023	189	1	13%	1	0%	0	1	1	3	low	NR
Body Weight	Boppre 2021	496	0	32%	1	0%	0	2	2	3	low	NR
	Ren 2018	347	0	ROB/n NR	1	51%	0	1	1	2	moderate	moderate
Body Mass Index	Gasmi 2022	341	0	ROB NR	1	85%	1	3	2	4	low	NR
	Boppre 2021	401	0	36%	1	0%	0	2	2	3	low	NR
	Ren 2018	259	0	ROB/k/n NR	1	44%	0	1	1	2	moderate	moderate
Waist Circumference	Boppre 2021	201	0	12%	1	9%	0	2	2	3	low	NR
	Ren 2018	198	1	ROB/k/n NR	1	94%	1	1	1	4	low	low

Fat Mass	Gasmi 2022	74	2	ROB NR	1	0%	0	3	2	5	very low	NR
	Boppre 2021	173	1	26%	1	0%	0	2	2	4	low	NR
	Ren 2018	186	1	ROB/k/n NR	1	95%	1	1	1	4	low	low
Fat-Free Mass	Gasmi 2022	54	2	ROB NR	1	0%	0	3	2	5	very low	NR
	Ren 2018	58	2	ROB/k/n NR	1	71%	0	1	1	4	low	very low
Lean-Body Mass	Boppre 2021	201	0	22%	1	0%	0	2	2	3	low	NR
Bone Density (Exercise v Control)	Roth 2022	63	2	100%	0	n/a	1	1 half	1	4	low	moderate
VO ₂ max	Boppre 2022	NR	1 ^g	total n NR	1	0%	0	3	2	4	low	NR
6-Minute Walking Test Distance	Ren 2018	65	2	ROB/n NR	1	0%	0	1	1	4	low	low
Resting Heart Rate	Boppre 2022	NR	1 ^g	total n NR	1	0%	0	3	2	4	low	NR
	Ren 2018	94	2	ROB/k/n NR	1	0%	0	1	1	4	low	low
Systolic Blood Pressure	Boppre 2022	314	0	74%	1	0%	0	3	2	3	low	moderate
	Ren 2018	229	0	ROB/k/n NR	1	6%	0	1	1	2	moderate	low
Diastolic Blood Pressure	Boppre 2022	NR	0 ^h	total n NR	1	59%	0	3	2	3	low	NR
	Ren 2018	229	0	ROB/k/n NR	1	83%	1	1	1	3	low	very low
Quality of Life	Bellicha 2021	NR	2 ⁱ	total n NR	1	0%	0	2 + 2 halves	2	5	very low	NR
Glucose Metabolism												
Fasting Insulin	Boppre 2022	NR	0 ^h	total n NR	1	71%	0	3	2	3	low	NR
	Marshall 2020	180	1	ROB NR	1	0%	0	1 + half	2	4	low	low
Fasting Glucose	Boppre 2022	NR	0 ^h	total n NR	1	0%	0	3	2	3	low	NR

	Marshall 2020	180	1	ROB NR	1	0%	0	1 + half	2	4	low	low
HOMA-IR	Boppre 2022	NR	2 ^j	total n NR	1	89%	1	3	2	6	very low	NR
	Bellicha 2021	NR	1 ^g	total n NR	1	0%	0	2 + 2 halves	2	4	low	NR
HbA1c	Boppre 2022	NR	2 ^j	total n NR	1	0%	0	3	2	5	very low	NR
Lipid Metabolism												
Total Cholesterol	Boppre 2022	NR	0 ^h	total n NR	1	0%	0	3	2	3	low	NR
	Marshall 2020	180	1	ROB NR	1	0%	0	1 + half	2	4	low	low
High-Density Lipoprotein	Boppre 2022	NR	0 ^h	total n NR	1	26%	0	3	2	3	low	NR
	Marshall 2020	180	1	ROB NR	1	0%	0	1 + half	2	4	low	low
Low-Density Lipoprotein	Boppre 2022	NR	0 ^h	total n NR	1	57%	0	3	2	3	low	NR
	Marshall 2020	180	1	ROB NR	1	0%	0	1 + half	2	4	low	low
Triglycerides	Boppre 2022	NR	0 ^h	total n NR	1	0%	0	3	2	3	low	NR
	Marshall 2020	180	1	ROB NR	1	0%	0	1 + half	2	4	low	low
POST-MBS Significant Subanalyses												
Combined Endurance/ Resistance on Systolic Blood Pressure	Boppre 2022	137	1	61%	1	0%	0	3	2	4	low	moderate
Combined Endurance/ Resistance on Triglycerides	Boppre 2022	171	1	35%	1	0%	0	3	2	4	low	low
Combined Endurance/ Resistance on Body Weight	Boppre 2021	221	0	28%	1	0%	0	2	2	3	low	NR
	Ren 2018	NR	1 ^g	total n NR	1	32%	0	1	1	3	low	NR

Combined Endurance/ Resistance on Body Mass Index	Boppre 2021	170	1	36%	1	0%	0	2	2	4	low	NR
Start time <6months on Body Weight	Boppre 2021	361	0	37%	1	0%	0	2	2	3	low	NR
	Ren 2018 ^k	NR	0	total n NR	1	32%	0	1	1	2	moderate	NR
Start time <6months on Body Mass Index	Boppre 2021	317	0	38%	1	0%	0	2	2	3	low	NR
Start time >6months on Systolic Blood Pressure	Boppre 2022	84	2	100%	0	0%	0	3	2	4	low	high
Start time >6months on Body Weight	Boppre 2021	135	1	18%	1	0%	0	2	2	4	low	NR
Start time >6months on Body Mass Index	Boppre 2021	84	2	29%	1	0%	0	2	2	5	very low	NR
Start time >12months on Body Weight	Ren 2018	NR	2 ⁱ	total n NR	1	0%	0	1	1	4	low	NR
Duration >12weeks on Systolic Blood Pressure	Boppre 2022	212	0	100%	0	0%	0	3	2	2	moderate	high

Note. MBS = metabolic and bariatric surgery; RCT = randomized controlled trial; SRMA = systematic review with meta-analysis; ROB = risk of bias; COE = certainty of evidence; DG = downgrade; NR = not reported; n = sample size; k = number of studies.

^a ≥ 200 no downgrade, 100-199 one downgrade, 1-99 two downgrade; ^b $\geq 75\%$ of participants have low ROB no downgrade, $< 75\%$ have low ROB one downgrade; ^c $I^2 \leq 75\%$ no downgrade, $I^2 > 75\%$ one downgrade; ^d 0 critical absences no downgrade, 1 absence one downgrade, 2+ absences two downgrades; ^e PreMBS: No RCT-only meta-analysis available for lean-body mass, muscle strength, resting heart rate, blood pressure, glucose/lipid metabolism, and physical activity; ^f PostMBS: No RCT-only meta-analysis available for muscle strength; ^g didn't report n but calculated to be between 100-199; ^h didn't report n but calculated to be ≥ 200 ; ⁱ didn't report n but calculated to be between 1-99; ^j studies not reported so n could not be calculated; ^k NOT a significant positive effect but included for comparison to Boppre 2021 which did find a significant positive effect.


Table S23: Data Missing from Certainty of Evidence Assessment for Trial Quality Risk of Bias

	PreMBS <i>n</i> = 10	PostMBS <i>n</i> = 39	PostMBS Subanalyses <i>n</i> = 13
% Low Risk of Bias Calculated [<i>n</i> (%)]	10 (100%)	8 (21%)	10 (77%)
% Low Risk of Bias NOT Calculated [<i>n</i> (%)]		31 (79%)	3 (23%)
<i>Risk of Bias Not Reported</i> [<i>n</i> (%)]		9 (23%)	
<i>Risk of Bias and Sample Size Not Reported</i> [<i>n</i> (%)]		2 (5%)	
<i>Risk of Bias, Sample Size, and Included Studies Not Reported</i> [<i>n</i> (%)]		7 (18%)	
<i>Total Sample Size Not Reported</i> [<i>n</i> (%)]		13 (33%)	3 (23%)

Note. *n* = the number of total entries (i.e., number of rows in Table S22).

Appendix P – Outcome Conclusions Flow Diagram

Figure S2: Outcome Conclusions Flow Diagram



Appendix Q – Downgrade Rules

1. Outcomes categorized into “what we still don’t know” cannot be downgraded further.
2. A maximum of one downgrade (i.e. one step) should be applied to an outcome finding.
3. Certainty of evidence (COE):
 - a. Outcomes categorized into “what we currently know” must be associated with a moderate or high certainty of evidence (COE). If COE is previously reported (by the review authors) or calculated (by the current authors) to be low/very low, outcome findings should be downgraded by one step (i.e., “what we currently know” → “what we think we know”).
 - b. Outcomes categorized into “what we think we know” must be associated with a low COE or higher. If COE is previously reported or calculated to be very low, outcome findings should be downgraded by one step (i.e., “what we think we know” → “what we still don’t know”).
 - c. If the previously reported and calculated COE are inconsistent, the COE reported by the review authors should be used to determine if a downgrade is necessary (in accordance with rules 3 and 4).
 - d. If multiple RCT-only meta-analyses and their respective COEs are presented for a single outcome, the COE of the most recent and comprehensive review should be used to determine if a downgrade is necessary (in accordance with rules 3 and 4).
 - e. If there is no RCT-only meta-analysis (i.e., no COE calculated by current authors), but there is a mixed (i.e., RCTs + NRCTs) meta-analysis with a previously reported COE, that COE should be used to determine if a downgrade is necessary (in accordance with rules 3 and 4).
4. If results vary between RCT only and mixed meta-analyses, the outcome should be downgraded by one step.
5. If all the meta-analyses for an outcome include data that is not confirmed to be final trial data (e.g., conference abstracts, posters), the outcome should be downgraded by one step.
6. Across reviews for a single outcome, if the total sample size is small (i.e., < 100) or if the number of total original studies is small (i.e., ≤ 3) and sample size is not reported, the outcome should be downgraded by one step.
7. In “what we think we know”, if an outcome is supported by multiple systematic reviews but no meta-analyses, if possible, results should be reviewed across the studies in the included systematic reviews (i.e., n_+ vs n_{NS}) to determine whether results favor the same direction (+/NS) of the original conclusion; if not, results should be downgraded by one step.
8. If all the meta-analyses for an outcome include multiple publications from a single original study, the outcome should be downgraded by one step.
9. If there is variability in the measures or domains used across systematic reviews for an outcome, it should be downgraded by one step.
10. If the effect of exercise training alone (i.e., without physical activity counselling) cannot be clearly determined based on the studies considered in the included systematic reviews, the outcome should be downgraded by one step.
11. If author conclusions caution interpretation of their results, outcome finding should be downgraded by one step.

Appendix R – Summary of Outcome Conclusions









Colours have been coordinated to Figure S2 in Appendix P for ease of interpretation.			
L		Concordant	 What we currently know
E		Discordant with potential reason	 What we think we know
G		Discordant without clear reason	 What we still don't know
E			
N		Inconclusive	* Downgraded outcome
D		N/A	

Table S24: Summary of PreMBS Exercise Training Outcome Conclusions and Categorizations

Outcome	# MAs # SRs	1 st Conclusion	Meta with $k=3+?$	2 nd Conclusion	Effect Sizes [95% confidence interval), I^2 , n , k with (<i>reported COE</i>) and/or [calculated COE] for RCT only MA(s)	Other considerations/ reason(s) for downgrade	Final Category
Pre1: BW/ BMI/ WL	1 MA (Durey 2022) 4 SR (Jabbour 2022, Lodewijks 2022, Schurmans 2022, Bellicha 2021) ** removed Herrera- Santelices 2022 (MA which includes 2 publications from 1 study)**	Discordance with potential explanation related to the inclusion of PAC Removing Lodejwiks and Durey (include PAC) → potential for concordance for + effect	No	What we think we know - ET (not PAC) likely has a + effect on BMI	RCT only MAs were removed		What we think we know - ET likely has a + effect on BMI
Pre2: FM	3 SR (Jabbour 2022, Lodewijks 2022, Bellicha 2021)	Inconclusive - No review with conclusive results		What we still don't know -			

	<i>**removed Herrera-Santelices 2022 (meta which includes 2 publications from 1 study)**</i>			Impact of ET on FM			
Pre3a: FFM	1 MA FFM (Herrera Santelices 2022)	N/A only 1 review	No	What we think we know - ET likely has a NS effect on FFM	Herrera 2022 SMD: - 0.41[-1.00; 0.18], I ² =0%; n=46, k=2, (Moderate) [Very low]	- 2 studies in total and small sample size (< 100)	* What we still don't know - Impact of ET on FFM
Pre3b: LBM	1 SR (Bellicha 2021)	N/A only 1 review and its findings are inconclusive		What we still don't know - Impact of ET on LBM			
Pre4: VO ₂ max	1 MA (Durey 2022) 2 SR (Jabbour 2022, Bellicha 2021)	Discordance without clear reason Prioritize RCT only MA (Durey) with + effect	Yes	What we currently know - Exercise Intervention has a + effect on VO ₂ max	Durey 2022 Pre-MBS VO ₂ max change: MD: 0.73 mL/kg/min [0.61; 0.86], I ² =62%; n=79, k=3, [Very low] VO ₂ max change at maximal follow up: MD: 0.98 mL/kg/min [0.05; 1.90], I ² =0%; n=131, k=3, [Low]	- COE - Includes 2 conference abstracts that may not be final data - Creel 2016 in MA may include pre and post effects	* What we think we know - Exercise Intervention likely has a + effect on VO ₂ max
Pre5: 6MWT	1 MA (Herrera Santelices 2022) 3 SR (Jabbour 2022,	Concordance for + effect	No	What we think we know -	Herrera Santelices 2022 SMD: 2.59 [1.89; 3.30], I ² =0%; n=61, k=2, (High) [Very low]		What we think we know -

	<i>Schurmans 2022, Bellicha 2021)</i>			ET likely has a + effect on 6MWTD			ET likely has a + effect on 6MWTD
Pre6: Strength	2 SR (Jabbour 2022, Bellicha 2021)	Discordance without clear reason		What we still don't know - Impact of ET on muscle strength			
Pre7: RHR	1 SR (<i>Schurmans 2022</i>) <i>**removed Marshall 2020 (Combined pre and post MBS results)**</i>	N/A only 1 review and its findings are inconclusive		What we still don't know - Impact of ET on RHR			
Pre8: BP	3 SR (<i>Schurmans 2022, Jabbour 2022, Bellicha 2021</i>) <i>**removed Marshall 2020 (Combined pre and post MBS results)**</i>	Concordance for NS effect	No	What we think we know - ET likely has NS effect on BP	N/A no MA(s)	- 4 studies total across the reviews; sample size not reported and 2 found + effect while 2 found NS effect	* What we still don't know - Impact of ET on BP
Pre9: QoL	1 MA (<i>Herrera Santelices 2022</i>) 3 SR (<i>Lodewijks 2022, Schurmans 2022, Bellicha 2021</i>)	Discordance without clear reason Prioritize Herrera-Santelices MA with NS effect (RCT)	Yes	What we currently know - ET has a NS effect on QoL	Herrera 2022 SMD: 0.88 [-0.23; 1.99], I ² =67%; n=53, k=3, (Moderate) [Very low]	- small sample size (< 100) - variability in QOL domains/measurements	* What we think we know - ET likely has a NS effect on QoL

Pre10: Glucose/ Lipid	2 SR (<i>Jabbour 2021, Bellicha 2021</i>)	N/A (Different outcome measures)		What we still don't know - Impact of ET on glucose/lipid metabolism			
Pre11: PA	1 SR (<i>Lodewijks 2022</i>)	N/A only 1 review and it includes PAC (conclusive findings)	No	What we think we know - Exercise intervention likely has a + effect on PA	N/A no MA(s)	- 4 studies but two are from the same original study - includes 2 PAC interventions	* What we still don't know - Impact of ET on PA
Pre12: Adverse (Surgical) Events	1 MA (<i>Durey 2022</i>)	N/A only 1 review and its findings are inconclusive due to inclusion of only 1 article		What we still don't know - Impact of ET on surgical adverse events			
Pre13: Hospital Stay	1 MA (<i>Durey 2022</i>) 1 SR (<i>Jabbour 2022</i>)	Inconclusive - Both the MA and SR only included 1 study		What we still don't know - Impact of ET on length of hospital stay			

Note. MBS= metabolic and bariatric surgery, MA=meta-analysis, SR=systematic review, I^2 =measure of heterogeneity, n =sample size, k =number of studies, COE=certainty of evidence, RCT=randomized controlled trial, BW=body weight, BMI=body mass index, WL=weight loss, FM=fat mass, FFM=fat-free mass, LBM=lean body mass, 6MWTD=6-minute walking distance, RHR=resting heart rate, BP=blood pressure, QoL=quality of life, PA=physical activity, N/A=not applicable, ET=exercise training, PAC=physical activity counselling, MD=mean difference, SMD=standardized mean difference.

Table S25: Summary of PostMBS Exercise Training Outcome Conclusions and Categorizations

Outcome	# MAs # SRs	1 st Conclusion	Meta with $k=3+?$	2 nd Conclusion	Effect Sizes [95% confidence interval), I^2 , n , k with (reported COE) and/or [calculated COE] for RCT only MA(s)	Other considerations/ reason(s) for downgrade	Final Category
Post1: WL \geq 12 months	1 MA (Bond 2023)	N/A only 1 review	Yes	What we currently know - ET has NS effect on WL \geq 12 months post-MBS	Bond 2023 SMD: - 2.26 [-2.07; 1.55], $I^2=0\%$; $n=189$, $k=5$, [Low]	- COE - authors list a need for more studies that specifically aim to improve weight loss maintenance	* What we think we know - ET likely has NS effect on WL \geq 12 months post-MBS
Post2: BW/ BMI/ WL	5 MA (Gasmi 2022, Boppre 2021, Bellicha 2021, Carretero Ruiz 2019, Ren 2018) 2 SR (Schurmans 2022, Morales Marroquin 2020)	Discordance with potential explanation Removing Schurmans (multiple same studies), Morales Marroquin (only resistance) and Carretero Ruiz (respiratory, physio and PAC) → concordance for a + effect	Yes	What we currently know - ET has a + effect on BW and BMI	Gasmi 2022 BMI: SMD: -0.93 [-1.65; -0.20], $I^2=85\%$; $n=341$, $k=5$, [Low] --- Boppre 2021 BMI: MD: -0.84 kg/m2 [- 1.60; -0.08], $I^2=0\%$; $n=401$, $k=7$, [Low] BW: MD: -2.51 kg [-4.74; - 0.27], $I^2=0\%$; $n=496$, $k=10$, [Low] --- Ren 2018 BMI: WMD: -0.40 kg/m2 [- 0.81; 0.00], $I^2=44\%$; $n=259$, $k=5$, (Moderate) [Moderate]	- COE (Boppre 2021 more comprehensive than Ren 2018)	* What we think we know - ET likely has a + effect on BW and BMI

					BW: WMD: -1.94 kg [-3.18; -0.69], $I^2=51\%$; $n=347$, $k=8$, (Moderate) [Moderate]		
Post3: WC	2 MA (Boppre 2021, Ren 2018) ** removed Gasmi 2022 as uses different measure**	Concordance for + effect	Yes	What we currently know - ET has a + effect on WC	Boppre 2021 MD: -4.14 cm [-8.16; -0.12], $I^2=9\%$; $n=201$, $k=4$, [Low] --- Ren 2018 WMD: -5.25 cm [-10.48; -0.03], $I^2=94\%$; $n=198$, $k=4$, (Low) [Low]	- COE	* What we think we know - ET likely has a + effect on WC
Post4: FM	4 MA (Gasmi 2022, Boppre 2021, Bellicha 2021, Ren 2018) 1 SR (Morales Marroquin 2020)	Discordance with potential explanation Bellicha (RCT/NRCT) → + effect, Morales (RCT/NRCT) which requires a resistance component → NS, and gold standard FM assessment (DXA and RCT only by Boppre) → NS	Yes	What we currently know - ET has a NS effect on FM	Gasmi 2022 SMD: -0.08 [-0.54; 0.38], $I^2=0\%$; $n=74$, $k=3$, [Very low] --- Boppre 2021 MD: -0.49 kg [-1.71; 2.69], $I^2=0\%$; $n=173$, $k=2$, [Low] --- Ren 2018 WMD: -3.35 kg [-7.99; 1.29], $I^2=95\%$; $n=186$, $k=3$, (Low) [Low]	- COE - different results when NRCTs are included	* What we think we know - ET likely has a NS effect on FM
Post5a: FFM	3 MA (Roth 2022, Gasmi 2022, Ren 2018) 2 SR (Schurmans 2022, Morales Marroquin 2020)	Discordance with potential explanation Schurmans has + ($k=2$) however one was only for	Yes	What we currently know - ET has NS effect on FFM	Gasmi 2022 SMD: 0.23 [-0.31; 0.77], $I^2=0\%$; $n=54$, $k=2$, [Very low] --- Ren 2018	- COE - supporting mixed RCT (Roth 2022) reports low/very low COE	* What we think we know - ET likely has NS

		combined E/R group and other was only at 24 weeks. Excluding Schurmans → concordance for a NS effect			WMD: 0.53 kg [-1.88; 2.94], $I^2=71\%$; $n=58$, $k=2$, (<i>Very low</i>) [Low]		effect on FFM
Post5b: LBM	2 MA (Boppre 2021, Bellicha 2021) 1 SR (Schurmans 2022)	Concordance for NS effect	Yes	What we currently know - ET has NS effect on LBM	Boppre 2021 MD: 0.87 [-0.65; 2.40], $I^2=0\%$; $n=201$, $k=3$, [Low]	- COE	* What we think we know - ET likely has NS effect on LBM
Post6: BMD	3 MA (Roth 2022, Diniz Sousa 2022, Bellicha 2021) 1 SR (Morales Marroquin 2020)	Concordance for + effect	Yes	What we currently know - ET has a + effect on BMD	Roth 2022 Ex vs. C: SMD: 0.51 [0.01; 1.01], $I^2=N/A$; $n=63$, $k=1$, (<i>Moderate</i>) [Low]	- RCT only MA only included 1 study; mixed (RCT+NRCT) MA supporting results reports low/very low COE	* What we think we know - ET likely has a + effect on BMD
Post7: VO2max	3 MA (Boppre 2022, Bellicha 2021, Carretero Ruiz 2021) 1 SR (Schurmans 2022) **removed da Silva (MA with 2 publications from same study)**	Discordance with potential explanation related to study design Remove Schurman (4 articles from 1 study)	Yes	What we currently know - ET has a NS effect on Vo2max	Boppre 2022 MD: 0.26 L/min [-0.11; 0.63], $I^2=0\%$; $n=NR$, $k=3$, [Low]	- COE - different results when NRCTs are included	* What we think we know - ET likely has a NS effect on Vo2max

		Boppre (only RCT) → NS effect Inclusion of NRCTs → + effect					
Post8: 6MWTD	2 MA (Bellicha 2021, Ren 2018) 1 SR (Schurmans 2022)	Discordance with potential explanation Removal of Schurmans (difficult interpretation of text in SR and in original articles) → concordance for a + effect	Yes	What we currently know - ET has a + effect on 6MWTD	Ren 2018 WMD: 29.67 m [25.97; 33.37], $I^2=0\%$; $n=65$, $k=2$, (Low) [Low]	- COE	* What we think we know - ET likely has a + effect on 6MWTD
Post9: Strength	2 MA (Vieira 2022, Bellicha 2021) 2 SR (Schurmans 2022, Morales Marroquin 2020)	Concordance for + effect	Yes	What we currently know - ET has a + effect on muscle strength	N/A no RCT only MA(s)	- mixed (RCT+NRCT) MA reported very low COE	* What we think we know - ET likely has a + effect on muscle strength
Post10: RHR	3 MA (Boppre 2022, Carretero Ruiz 2021, Ren 2018) 1 SR (Schurmans 2022) **removed Marshall 2020 (Combined pre and post MBS results)**	Discordance without clear reason Prioritize Boppre (only RCT) → NS effect and Carretero	Yes	What we currently know - ET has a NS effect on RHR	Boppre 2022 MD: -2.05 bpm [-6.64; 2.54], $I^2=0\%$; $n=NR$, $k=3$, [Low] --- Ren 2018 WMD: -4.39 bpm [-8.11; -0.68], $I^2=0\%$; $n=94$, $k=3$, (Low) [Low]	- COE - different results when NRCTs are included	* What we think we know - ET likely has a NS effect on RHR

		(RCT/NRCT) → + effect					
Post11a: SBP	4 MA (<i>Boppre 2022, Carretero Ruiz 2021, Bellicha 2021, Ren 2018</i>) 1 SR (<i>Schurmans 2022</i>) <i>**removed Marshall 2020 (Combined pre and post MBS results)**</i>	Discordance without clear reason Potential reason related to study inclusion. MAs that include only RCT (Boppre and Ren) → + effect, MA that include (RCT/NRCT) → NS effect	Yes	What we currently know - ET has a + effect on SBP	Boppre 2022 SBP: MD: - 5.33 mmHg [- 8.99; -1.66], $I^2=0\%$; $n=314$, $k=6$, (<i>Moderate</i>) [Low] --- Ren 2018 SBP: WMD: -4.12 mmHg [- 6.68; -1.55], $I^2=6\%$; $n=229$, $k=4$, (<i>Low</i>) [Moderate]	- no COE downgrade as Boppre 2021 is more comprehensive than Ren 2018 - different results when NRCTs are included	* What we think we know - ET likely has a + effect on SBP
Post11b: DBP	4 MA (<i>Boppre 2022, Carretero Ruiz 2021, Bellicha 2021, Ren 2018</i>) 1 SR (<i>Schurmans 2022</i>) <i>**removed Marshall 2020 (Combined pre and post MBS results)**</i>	Concordance for NS effect	Yes	What we currently know - ET has a NS effect on DBP	Boppre 2022 MD: -2.66 mmHg [-6.72; 1.40], $I^2=59\%$; $n=NR$, $k=6$, [Low] --- Ren 2018 WMD: -3.56 mmHg [-8.61; 1.48], $I^2=83\%$; $n=229$, $k=4$, (<i>Very low</i>) [Low]	- COE	* What we think we know - ET likely has NS effect on DBP
Post12: QoL	1 MA (<i>Bellicha 2021</i>) 1 SR (<i>Schurmans 2022</i>)	Concordance for NS effect	No	What we think we know - ET likely has a NS effect on QoL***	Bellicha 2021 Physical.: MD: -2.5 [-5.1; 0.2], $I^2=0\%$; $n=NR$, $k=2$, [Very low] Mental: MD: 3.9 [-0.5; 8.3], $I^2=0\%$; $n=NR$, $k=2$, [Very low]	- COE - unclear domains of QoL evaluated and sample size not reported - only 3 studies across reviews and sample size not reported	* What we still don't know - Impact of ET on QoL

Post13a: Glucose: Fasting insulin and glucose	2 MA (<i>Boppre 2022, Marshall 2020</i>)	Concordance for a NS effect	Yes Insulin and glucose	What we currently know - ET has a NS effect on fasting glucose and insulin	Boppre 2022 Fasting insulin: MD: -1.58 μ IU/mL [-5.14; 1.98], $I^2=71\%$; $n=NR$, $k=4$, [Low] Fasting glucose: MD: 0.94 mg/dL [-3.31; 5.19], $I^2=0\%$; $n=NR$, $k=4$, [Low] --- Marshall 2020 Fasting insulin: MD: 4.88 pmol/L [-2.09; 11.84], $I^2=0\%$; $n=180$, $k=2$, (Low) [Low] Fasting glucose MD: 0.05 mmol/L [-0.14; 0.24], $I^2=0\%$; $n=180$, $k=2$, (Low) [Low]	- COE	* What we think we know - ET likely has a NS effect on fasting glucose and insulin
Post13b: Glucose: HOMA-IR	2 MA (<i>Boppre 2022, Bellicha 2021</i>) 1 SR (<i>Schurmans 2022</i>)	Concordance for a NS effect	No	What we think we know - ET likely has a NS effect on HOMA-IR	Boppre 2022 MD: 1.39 [-1.30; 4.08], $I^2=71\%$; $n=NR$, $k=4$, [Very low] --- Bellicha 2021 SMD: 0.14 [-0.10; 0.38], $I^2=0\%$; $n=NR$, $k=2$, [Low]	- COE (Boppre 2022 more comprehensive than Bellicha 2021)	* What we still don't know - Impact of ET on HOMA-IR
Post13c: Glucose: HbA1c	1 MA (<i>Boppre 2022</i>)	N/A only 1 review	No	What we think we know - ET likely has a NS effect on HbA1c	Boppre 2022 MD: -0.65 mmol/mol [-2.22; 0.93], $I^2=0\%$; $n=NR$, $k=2$, [Very low]	- COE - only 2 studies and sample size not reported	* What we still don't know - Impact of ET on HbA1c

Post13d: Glucose	1 SR (<i>Schurmans 2022</i>)	N/A only 1 review and its findings are inconclusive		What we still don't know - Impact of ET on insulin sensitivity, AIRg, Di, SPISE and glucose effectiveness			
Post14a: Lipid: TG, LDL, and TC	4 MA (<i>Boppre 2022, Carretero Ruiz 2021, Bellicha 2021, Marshall 2020</i>) 1 SR (<i>Schurmans 2022</i>)	Concordance for a NS effect	Yes	What we currently know - ET has a NS effect on TG, LDL or TC	<p>Boppre 2022 TG: MD: -8.38 mg/dL [-19.81; 3.04], $I^2=0\%$; $n=NR$, $k=5$, [Low] LDL: MD: -8.17 mg/dL [-20.35; 4.00], $I^2=57\%$; $n=NR$, $k=5$, [Low] TC: MD: -3.08 mg/dL [-12.04; 5.87], $I^2=0\%$; $n=NR$, $k=5$, [Low] ---</p> <p>Marshall 2020 TG: MD: 0.01 mmol/L [-0.15; 0.16], $I^2=0\%$; $n=180$, $k=2$, (<i>Low</i>) [Low] LDL: MD: -0.06 mmol/L [-0.21; 0.09], $I^2=0\%$; $n=180$, $k=2$, (<i>Low</i>) [Low] TC: MD: -0.08 mmol/L [-0.26; 0.11], $I^2=0\%$; $n=180$, $k=2$, (<i>Low</i>) [Low]</p>	- COE	* What we think we know - ET likely has a NS effect on TG, LDL or TC

Post14b: Lipid: HDL	4 MA (<i>Boppre 2022, Carretero Ruiz 2021, Bellicha 2021, Marshall 2020</i>) 1 SR (<i>Schurmans 2022</i>)	Discordance with potential explanation related to study design Only RCT (Boppre and Marshall) → NS effect. Carretero and Bellicha both include NRCTs but Carretero is more comprehensive thus inclusion of NRCTs → + effect	Yes	What we currently know - ET has a NS effect on HDL	Boppre 2022 MD: 0.61 mg/dL [-3.05; 4.28], $I^2=26\%$; $n=NR$, $k=5$, [Low] --- Marshall 2020 MD: -0.00 mmol/L [-0.01; 0.01], $I^2=0\%$; $n=180$, $k=2$, (Low) [Low]	- COE - different results when NRCTs are included	* What we think we know - ET likely has a NS effect on HDL
--------------------------------------	--	--	-----	---	---	--	---

Note. MBS= metabolic and bariatric surgery, MA=meta-analysis, SR=systematic review, I^2 =measure of heterogeneity, n =sample size, k =number of studies, COE=certainty of evidence, RCT=randomized controlled trial, NRCT=non randomized controlled trial, WL=weight loss, BW=body weight, BMI=body mass index, WC=waist circumference, FM=fat mass, DXA=dual x-ray absorptiometry, FFM=fat-free mass, LBM=lean body mass, BMD=bone mineral density, 6MWT=6-minute walking distance, RHR=resting heart rate, SBP=systolic blood pressure, DBP=diastolic blood pressure, QoL=quality of life, HOMA-IR= homeostatic model assessment for insulin resistance, HbA1c=hemoglobin A1C, AIRg=acute insulin response to glucose, Di=disposition index, SPISE=single-point insulin sensitivity estimator, TG=triglycerides, LDL=low density lipoprotein, HDL=high density lipoprotein, TC=total cholesterol, N/A=not applicable, ET=exercise training, PAC=physical activity counselling, MD=mean difference, SMD=standardized mean difference.

Table S26: Summary of PostMBS Exercise Training Sub Analysis Outcome Conclusions and Categorizations

Outcome	# MAs # SRs	1 st Conclusion	Meta with $k=3+$?	2 nd Conclusion	Effect Sizes [95% confidence interval), I^2 , n , k with (<i>reported COE</i>) and/or [calculated COE] for RCT only MA(s)	Other considerations/ reason(s) for downgrade	Final Category
Sub1: Combined E/R on BW/BMI/ WL	3 MA (<i>Boppre 2021,</i> <i>Carretero</i> <i>Ruiz 2019,</i> <i>Ren 2018</i>)	Discordance with potential explanation related to study design Only RCT → + effect Inclusion of NRCTs and alternative interventions (physiotherapy and respiratory) → NS effect	Yes	What we currently know - Combined E/R has a + effect on BW/BMI/WL	Boppre 2021 BW: MD: -5.02 kg [-8.13; - 1.90], $I^2=0\%$; $n=221$, $k=5$, [Low] BMI: MD: -1.62 kg/m ² [- 2.72; -0.59], $I^2=0\%$; $n=170$, $k=4$, [Low] --- Ren 2018 BW: MD: -7.71 mmHg [- 13.12; -2.31], $I^2=0\%$; $n=84$, $k=2$, [Low]	- COE - different results when NRCTs are included	* What we think we know - Combined E/R likely has a + effect on BW/BMI
Sub2: Combined E/R on SBP and TG	1 MA (<i>Boppre</i> <i>2022</i>)	N/A only 1 review	Yes	What we currently know - Combined E/R has a + effect on SBP and TG	Boppre 2022 SBP: MD: -7.18 mmHg [- 12.42; -1.94], $I^2=0\%$; $n=137$, $k=3$, (<i>Moderate</i>) [Low] TG: MD: -17.56 mg/dL [- 34.15; -0.96], $I^2=0\%$; $n=171$, $k=3$, (<i>Low</i>) [Low]	- COE for TG	What we currently know - Combined E/R has a + effect on SBP * What we think we know - Combined E/R likely

							has a + effect on TG
Sub3: Exercise start time < 6 months post-MBS on BW/BMI	2 MA (Boppre 2021, Ren 2018)	Discordance without clear reason Boppre newer and more comprehensive than Ren	Yes	What we currently know - Exercise start time < 6 months has a NS effect on BW/BMI	Boppre 2021 BW: MD: -0.12 kg [-3.17; 2.93], I ² =0%; n=361, k=7, [Low] BMI: MD: -0.16 kg/m ² [- 1.15; 0.82], I ² =0%; n=317, k=5, [Low]	- COE	* What we think we know - Exercise start time < 6 months likely has NS effect on BW/BMI
Sub4: Exercise start time > 6 months post-MBS on SBP	1 MA (Boppre 2022)	N/A only 1 review	No	What we think we know - Exercise start time > 6 months likely has a + effect on SBP	Boppre 2022 MD: -7.71 mmHg [-13.12; -2.31], I ² =0%; n=84, k=2 (High) [Low]	- only 2 studies and small sample size (<100)	* What we still don't know - Impact of exercise start time > 6 months on SBP
Sub5: Exercise start time > 6 months post-MBS on BW/BMI	1 MA (Boppre 2021)	N/A only 1 review	Yes	What we currently know - Exercise start time > 6 months has a + effect on BW/BMI	Boppre 2021 BW: MD: -5.25 kg [-8.52; - 1.97], I ² =0%; n=135, k=3, [Low] BMI: MD: -1.84 kg/m ² [- 3.04; -0.64], I ² =0%; n=84, k=2, [Very low]	- COE	* What we think we know - Exercise start time > 6 months likely has a + effect on BW/BMI

Sub6: Exercise start time > 12 months post-MBS on BW	1 MA (Ren 2018)	N/A only 1 review	No	What we think we know - Exercise start time > 12 months likely has a + effect on BW	Ren 2018 BW: WMD: -3.63 kg [-5.35; -1.91], I ² =0%; n=NR, k=2, [Low]	- only 2 studies and small sample not reported	* What we still don't know - Impact of exercise start time > 12 months on BW
Sub7: Intervention duration > 12 weeks on SBP	1 MA (Boppre 2022)	N/A only 1 review	Yes	What we currently know - Intervention duration > 12 weeks has a + effect on SBP	Boppre 2022 MD: -5.78 mmHg [-9.91; -1.66], I ² =0%; n=212, k=3 (High) [Moderate]		What we currently know - Intervention duration > 12 weeks has a + effect on SBP

Note. MBS= metabolic and bariatric surgery, MA=meta-analysis, SR=systematic review, I²=measure of heterogeneity, n=sample size, k=number of studies, COE=certainty of evidence, RCT=randomized controlled trial, NRCT=non randomized controlled trial, BW=body weight, BMI=body mass index, WL=weight loss, SBP=systolic blood pressure, TG=triglycerides, E/R=endurance and resistance, N/A=not applicable, MD=mean difference, WMD=weighted mean difference.

Table S27: Summary of Feasibility and Acceptability Outcome Conclusions and Categorizations

Outcome	# MAs # SRs	1 st Conclusion	Meta with $k=3+$?	2 nd Conclusion	Effect Sizes [95% confidence interval), I^2 , n , k with (<i>reported COE</i>) and/or [calculated COE] for RCT only MA(s)	Other considerations/ reason(s) for downgrade	Final Category
Feasibility and Acceptability 1: Attendance, drop out, and retention rates for the exercise arm	1 MA (Baillot 2022)	N/A only 1 review	Yes	What we currently know - Exercise training has high attendance and retention rates, and low drop out rates	All analyses were conducted on mixed (RCT and NRCT) samples	- only 1 review and authors conclude that results should be interpreted with caution as outcomes are rarely reported in the primary articles.	* What we think we know - Exercise training likely has high attendance and retention rates, and low drop out rates

Note. MA=meta-analysis, SR=systematic review, I^2 =measure of heterogeneity, n =sample size, k =number of studies, COE=certainty of evidence, RCT=randomized controlled trial, NRCT=non randomized controlled trial, N/A=not applicable.