



# Sex differences in sit-and-reach flexibility in children and adolescents: a meta-analysis

Supplementary materials:

<https://osf.io/yjexh/>

For correspondence:

[j.nuzzo@ecu.edu.au](mailto:j.nuzzo@ecu.edu.au)

X: [@JamesLNuzzo](#)

James L. Nuzzo

School of Medical and Health Sciences, Edith Cowan University, Joondalup, Australia

*Please cite as:* Nuzzo JL. (2024). Sex differences in sit-and-reach flexibility in children and adolescents: a meta-analysis. *SportRxiv*.

## ABSTRACT

The sit-and-reach, which primarily assesses hamstrings extensibility, is arguably the most frequently performed test of flexibility in history. Thus, the sit-and-reach can be used to explore, with high statistical power, sexual dimorphism of a proposed fitness attribute throughout development. Here, the aim was to use meta-analysis to examine sex differences in sit-and-reach flexibility in children and adolescents. The analysis included 408 effects from 96 studies conducted in 38 countries between 1983–2023. The total sample was 944,420 (484,380 boys, 460,040 girls). At all ages (3-16 years old), girls had greater sit-and-reach flexibility than boys. Between 4–10 years old, the effect size was fairly consistent and moderate in size ( $g = -0.30$  to  $-0.50$ ). The sex difference widened at age 11 ( $g = -0.59$ , 95% confidence intervals (CIs)  $[-0.75, -0.44]$ ), peaked at age 12 ( $g = -0.78$ , 95% CIs  $[-0.96, -0.61]$ ), then returned to pre-pubertal levels by age 16 ( $g = -0.42$ , 95% CIs  $[-0.61, -0.52]$ ). Secondary analyses revealed the sex difference in sit-and-reach flexibility is similar between countries and has remained relatively stable since the 1980s. Results confirm that sexual dimorphism in flexibility occurs early in development, is impacted by puberty, and is broadly consistent across time and place.

## INTRODUCTION

Flexibility refers to the intrinsic properties of body tissues that determine maximal joint range of motion (ROM) without causing injury (Holt et al., 1996; Knudson et al., 2000). A common test of flexibility is the sit-and-reach. The sit-and-reach was created in 1952 (Wells & Dillon, 1952). Since its creation, the sit-and-reach has been included in various physical fitness test batteries around the world (Marques et al., 2021).

The sit-and-reach assesses hamstrings extensibility, and to a lesser extent, lumbar extensibility (Castro-Piñero et al., 2009; Chillón et al., 2010; Hartman & Looney, 2003; Mayorga-Vega et al., 2014; Muyor et al., 2014; Patterson et al., 1996). Concerns exist about the concurrent validity between sit-and-reach performance and meaningful health and performance outcomes. Consequently, some researchers have suggested that the sit-and-reach, and perhaps flexibility more broadly, be de-emphasized or altogether removed from fitness test batteries (Institute of Medicine, 2012; Nindl et al., 2015; Nuzzo, 2020; Reuben et al., 2013). Nevertheless, the sit-and-reach is arguably the most frequently performed test of flexibility in history. Thus, it provides, with high statistical power, a unique opportunity to explore sexual dimorphism of a proposed fitness attribute.

In adults, sex differences in sit-and-reach flexibility are clear, with women significantly more flexible than men (Davis et al., 2008; Jackson & Langford, 1989; Katzmarzyk et al., 2000; Liemohn et al., 1994; Shephard et al., 1990). The development of this sex difference during childhood and adolescents is less clear. For example, it is unclear at what age the sex difference in sit-and-reach flexibility emerges and to what extent the size of the difference changes throughout development. Meta-analysis of all existing data can be used to clarify these questions.

In 1985, Thomas and French (1985) used meta-analysis to examine sex differences in flexibility and other physical attributes in boys and girls. They found that girls had greater sit-and-reach flexibility than boys ( $d = -0.29$ ). Nevertheless, their analysis, which is now 40 years old, had limitations. First, it included only 13 effects from 2,069 boys and girls (Thomas & French, 1985). Originally, the researchers reported that these 13 effects came from five studies (Thomas & French, 1985), but they later revealed that these effects came from only four studies - one of which was not published in a journal (Thomas & French, 1987). Second, the ages of the participants who were included in the analysis of the sit-and-reach were not stated. Thus, how

the sex difference in sit-and-reach flexibility might modulate throughout development was not examined.

A few years later, the same researchers conducted a more powerful analysis (Thomas et al., 1991). They generated effect sizes of the sex difference in sit-and-reach flexibility in a sample of over 12,000 boys and girls aged 6-18 years (Thomas et al., 1991). The effect size during early childhood was approximately  $d = -0.50$ . During puberty, it increased to approximately  $d = -1.00$ . At ages 17 and 18, it returned to pre-puberty levels ( $d = -0.60$ ). Thus, Thomas et al. (1991) found that girls had greater sit-and-reach flexibility than boys at all ages assessed, but the magnitude of the difference changed throughout development.

These two seminal studies enhanced knowledge of sex differences in sit-and-reach flexibility in children and adolescents (Thomas & French, 1985; Thomas et al., 1991). Nevertheless, both analyses are now decades old and can be expanded in various ways. First, significantly more studies and effects are now available. Second, neither analysis included confidence intervals (CI) of the effect sizes, and such information is important for establishing confidence in any sex differences observed. Third, the two studies included sit-and-reach flexibility data only from children in the United States of America (USA). Sit-and-reach flexibility data are now available from boys and girls who reside in various countries, allowing for examination of how sex differences in sit-and-reach flexibility might differ between countries. Finally, given that the sit-and-reach test was created over 70 years ago (Wells & Dillon, 1952), secular changes in any sex differences in flexibility can now be explored.

Therefore, the primary aim of the current study was to use meta-analysis to provide an updated examination of sex differences in sit-and-reach flexibility in children and adolescents. The study also had two secondary, exploratory aims. The first exploratory aim was to examine if sex differences in sit-and-reach flexibility in children and adolescents have changed over time (i.e., secular analysis). The second exploratory aim was to examine if the size of the sex difference in sit-and-reach flexibility in children and adolescents differs between countries. Results from these analyses have potential to inform discussions about sex differences in proposed fitness attributes and the impact of such differences on male and female sports performances and policies (Brown et al., 2024; Hamilton et al., 2024; Hilton & Lundberg, 2021; Lundberg et al., 2024; Nokoff et al., 2023; Nuzzo, 2023a; Tucker et al., 2024).

## **METHOD**

### **Literature search**

The literature search for this paper was performed in August of 2024. Many papers on the topic were already known to me based on searches conducted for other reviews (Nuzzo, 2020, 2024). The search strategy was similar to that described by Greenhalgh and Peacock (2005). The first component of the search involved checking personal digital files associated with previous research (Nuzzo, 2020). The second component involved performing relevant keyword searches in PubMed and Google Scholar. Example keyword searches included various combinations of words such as "children," "boys," "girls," "youth," "adolescents," "flexibility," "sit-and-reach," "sit-reach," "fitness," and "physical fitness." The third component involved "snowballing" strategies, such as reference and citation tracking. The literature search, article screening, and data extraction were conducted by a single author. I have used these strategies successfully in previous reviews and meta-analyses, which have identified more papers on their relevant topics than other previous attempts (Nuzzo, 2023b; Nuzzo, Pinto, & Nosaka, 2023; Nuzzo, Pinto, Nosaka, et al., 2023; Nuzzo et al., 2024). Nevertheless, these approaches have limitations. These limitations are described in the Discussion. The current project was not pre-registered.

### **Eligibility**

For a study to be included in the current meta-analysis, it needed to meet the following criteria: (a) published in an academic journal in 2023 or earlier; (b) published in English; (c) included male and female participants who were between the ages of birth to 16 years old, healthy, and not competitive athletes; (d) included sample sizes and sex-segregated means and standard deviations (SD) of sit-and-reach scores that were not statistically adjusted for covariates. Common reasons for exclusion were: (a) no sample sizes reported; (b) no means or SDs reported; (c) a test other than the sit-and-reach was performed (e.g., stand-and-reach); and (d) the age range of the cohort was beyond the established criteria (described below).

### **Data extraction and organization**

Extracted data from eligible studies included year of publication, year of data collection (if provided), sample size, sample age, and means and SDs of sit-and-reach scores. Researchers

reported their data in various ways. Consequently, I established the following criteria for data extraction and organization.

***Year of data collection.*** Year of data collection is more scientifically valid for examining secular changes in sit-and-reach flexibility. Year of data collection represents the true time at which a measurement was taken. Year of publication, on the other hand, could be years after the measurement was taken. When researchers listed a single year for date of data collection, that year was used to represent that study in the current analysis. Some researchers reported dates of data collection that spanned two or more years. Often, their data were not reported by each year, but, instead, were aggregated as a single mean. When the duration of data collection spanned two years (e.g., 2007-2008), the first of the two years was used to represent that study in the current analysis. When the duration of data collection spanned more than two years (e.g., 2007-2009), the year in the middle (i.e., 2008) was used to represent that study in the current analysis. When year of data collection was not reported by researchers, the year of publication was used to represent that study in the current analysis.

***Age category.*** Researchers presented information on participant age in different ways. Given these differences, I adopted eligibility and data extraction criteria that were liberal enough to allow for as much data as possible to be included in the analysis, while not compromising the validity of the current study's aims.

I treated age as a categorical variable (e.g., 6-year-olds) rather than a continuous variable (e.g., 6.2 years). This strategy prioritized data that were presented for categorical age groups that spanned only one year (e.g., "5-year-olds"). However, under certain conditions, the strategy also permitted sit-and-reach scores from studies in which age cohorts spanned multiple years. First, if the researcher reported sit-and-reach scores for a cohort whose age spanned four or more years (e.g., 10- to 15-year-olds), the study was ineligible for inclusion in the current analysis. Second, if the researcher reported sit-and-reach scores for a cohort whose age spanned three years (e.g., 10- to 12-year-olds), the middle age (i.e., 11 years old) was typically used to represent these data in the current analysis. Often, this middle age corresponded with the rounded down mean age of the cohort (e.g., 11.3 years in a cohort of 10- to 12-year-olds whose ages could have ranged from 10.0 to 12.9 years). Third, if the researcher reported sit-and-reach scores for a cohort whose age spanned two years (e.g., 10- to 11-year-olds), the younger of the two ages (i.e., 10 years old) was used to represent the study in the current analysis. Often, in such scenarios the rounded down mean corresponded more closely with the younger of the two ages (e.g., mean age of 10.3 years in a cohort of 10- and 11-year-olds whose ages could have ranged from

10.0 to 11.9 years). Finally, if any participants in a cohort were 17 years of age or older, the study was ineligible for the current analysis.

**Study type.** Sit-and-reach flexibility data were reported in studies of various designs. Common study designs included longitudinal studies on development, reliability studies on consistency of sit-and-reach scores, cross-sectional studies comparing cohorts of children who differed on some characteristic (e.g., body weight status), and intervention studies that compared sit-and-reach scores before and after interventions (e.g., exercise program).

For longitudinal studies, grip strength data from each year of development were included in the current analysis. For reliability studies, only grip strength data from the first trial or first day of testing were included in the current analysis. For cross-sectional studies that compared sit-and-reach scores in healthy children (“controls”) and children with health conditions, only data from the healthy children were included in the current analysis. For intervention studies that involved baseline and post-intervention assessments of sit-and-reach flexibility, only data from the baseline assessments were included in the current analysis.

**Data extraction from graphs.** Researchers typically presented sit-and-reach scores in tables. When scores were presented in graphs, I used a graph digitizer (WebPlotDigitizer, <https://apps.automeris.io/wpd/>) to estimate them (Aydin & Yassikaya, 2022; Nuzzo, 2023b; Nuzzo, Pinto, Nosaka, et al., 2023; Nuzzo et al., 2024). With the digitizer, I calibrated the y-axis, inputting the sit-and-reach scores associated with the bottom and top of the y-axis. I then clicked each symbol on the graph that represented a mean and SD of interest. The software generated a spreadsheet of the means and SDs calibrated against the y-axis. Standard errors, whether presented in graphs or tables, were converted to SDs by multiplying the standard error by the square root of the sample size.

## Statistical analysis

The data spreadsheet and statistical results are available at the Open Science Framework (<https://osf.io/yjexh/>). Version 29 of the Statistical Software Package for the Social Sciences (SPSS, IBM Corporation, Armonk, NY, USA) was used to complete the statistical analyses. Frequency counts were generated for the number of effects by age, country, and year of data collection. Random effects meta-analyses were used to generate effect sizes (Hedges  $g$ ) with 95% confidence intervals (CI) and prediction intervals. Forest plots were used to graphically display the effect sizes for each study. Effect sizes equal to 0.2, 0.5, and 0.8 are often considered small, moderate, and large, respectively, but such benchmarks are arbitrary and should be interpreted cautiously (Lakens, 2013). Confidence intervals that do not cross zero indicate effects that are

statistically significant (i.e.,  $p \leq 0.05$ ) (Cumming, 2009). Funnel plots of tests of heterogeneity were also generated.

For the analysis of secular trends, cumulative effect sizes were generated for two large age cohorts (5-10- and 14-16-years-olds) and across each of the six decades from which data were available from these two cohorts (1980s, 1990s, 2000s, 2010s, and 2020s [i.e., 2020-2023]). Ages 11-13 are most closely aligned with puberty onset in both sexes. Thus, removing participants in this age range from the secular and between-countries analyses allowed for examination of one cohort who is mostly pre-pubescent (5-10-year-olds) and another cohort who is mostly post-pubescent (14-16-year-olds). This segregation helped to minimize confounding of pubertal status on the effect of time. For example, if one decade or country included proportionally more samples of 5-10-year-olds, whereas other decades or countries included proportionally more samples of 14-16-year-olds, any observed difference over time or between countries would be confounded by pubertal status. Moreover, this segregation helped minimize a potentially confounding influence of secular changes in age at menarche (Gomula & Koziel, 2018; Lei et al., 2021; Wang et al., 2024) and differences in age at menarche between countries (Lei et al., 2021; Saczuk et al., 2018; Wang et al., 2024).

For the analysis that compared sex differences in sit-and-reach flexibility between countries, cumulative effect sizes were generated for 5-10-years-olds and 14-16-years-olds for each country. For some countries, few effects were available, rendering their inclusion into the comparison uninformative. Thus, only countries in which sufficient effects were available are discussed.

## **Results**

### **Study characteristics**

A total of 96 studies met the eligibility criteria and were included in the meta-analysis (Amado-Pacheco et al., 2019; Anselma et al., 2020; Armstrong et al., 2011; Ayán Pérez et al., 2020; Bae et al., 2015; Baquet et al., 2006; Bartkowiak et al., 2021; Batez et al., 2021; Benzo et al., 2023; Casajús et al., 2007; Chagas & Barnett, 2023; Chen et al., 2022; Cieśła et al., 2014; Cieśła et al., 2017; De la Cruz-Sánchez & Pino-Ortega, 2010; Deforche et al., 2003; Docherty & Bell, 1985; Eather et al., 2016; Estivaleti et al., 2022; Fang et al., 2017; Fiori et al., 2021; Flanagan et al., 2015; Fogelholm et al., 2008; Fortier et al., 2001; Godoy-Cumillaf et al., 2020; Golle et al., 2014; Gontarev et al., 2018; Gulías-González et al., 2014; Haapala et al., 2015; Haapala et al., 2016; Hands et al., 2009; Haugen et al., 2014; He et al., 2019; Henriques-Neto et al., 2022; Hong & Hamlin, 2005; Hsu et al., 2021; Huang & Malina, 2002; Jones et al., 2005; Jürimäe & Saar, 2003;

Jürimäe & Volbekiene, 1998; Kamandulis et al., 2013; Karppanen et al., 2012; Katzmarzyk et al., 2000; Kidokoro et al., 2016; Kim & Park, 2017; Kondric et al., 2013; Koslow, 1987; Lehnhard et al., 1992; Lintu et al., 2016; Lo et al., 2017; Lovecchio, Giuriato, et al., 2019; Lovecchio et al., 2015; Lovecchio, Novak, et al., 2019; Mačak et al., 2022; Marshall et al., 1998; Martinho et al., 2022; McMillan & Erdmann, 2010; Moliner-Urdiales et al., 2010; Monyeki et al., 2005; Oja & Jürimäe, 1997, 2002; Örjan et al., 2005; Ortega et al., 2008; Panczyk et al., 2014; Pelicer et al., 2016; Pissanos et al., 1983; Podstawski & Boryślawski, 2012; Puszczalowska-Lizis et al., 2023; Richards et al., 2022; Riddoch et al., 1991; Runhaar et al., 2010; Ruzbarsky et al., 2022; Ryu et al., 2021; Sacchetti et al., 2012; Safrit & Wood, 1987; Santos et al., 2023; Sasayama & Adachi, 2019; Sember et al., 2022; Siegel et al., 1989; Smith & Miller, 1985; Sokolowski & Chrzanowska, 2012; Tishukaj et al., 2017; True et al., 2021; Tsimeas et al., 2005; Tsoukos & Bogdanis, 2021; Vanhelst et al., 2017; Veraksa et al., 2021; Vitali et al., 2019; Vlahov et al., 2014; Volbekiene & Gričiūte, 2007; Welk et al., 2015; Weston et al., 2019; Xu et al., 2020; Zhang et al., 2021)

The studies included a total of 408 effects from 944,420 children and adolescents (484,380 boys, 460,040 girls). The number of effects available by age and country are listed in Table 1.

Study publication dates ranged from 1983 to 2023. The earliest year of data collection was 1980. Numbers of effect available by decade of data collection and across all ages were as follows: 1980s (60 effects, 14.7%), 1990s (35 effects, 8.6%), 2000s (95 effects, 23.3%), and 2010s (217 effects, 53.3%).

The number of effects available by country and across all ages are listed in Table 2. Data were available from 38 countries. Study samples from Poland contributed the greatest number of effects to the meta-analysis (63 effects, 15.4%), followed by the USA (42 effects, 10.3%), South Africa (32 effects, 7.8%), Italy (27 effects, 6.6%), China (23 effects, 5.6%) and Spain (22 effects, 5.4%).

### Sit-and-reach flexibility

**Age.** From age 3, sit-and-reach flexibility was greater in girls than boys (Fig. 1, Table 1). At age 3, only one effect was available. Consequently, the 95% CIs were much wider at age 3 than other ages. From age 4 to 10 years old, effect sizes were small-to-moderate in magnitude, typically ranging between  $g = -0.30$  –  $-0.50$ . The effect size then increased at age 11 ( $g = -0.59$ , 95% CIs  $[-0.75, -0.44]$ ) and peaked at age 12 ( $g = -0.78$ , 95% CIs  $[-0.96, -0.61]$ ). Starting at age 13, the effect size began to decrease, and by age 16, it returned to pre-puberty levels ( $g = -0.42$ , 95% CIs  $[-0.61, -0.52]$ ).

**Secular changes.** For 5-10-year-olds, the mean effect size has remained broadly similar since the 1980s, as evident by the overlap in the 95% CIs across decades (Fig. 2, Table 3).



Similarly, for 14-16-year-olds, the mean effect size has remained broadly similar since the 1980s, as evident by the overlap in the 95% CIs across decades (Fig. 3, Table 3).

**Country.** Effect sizes of the difference in sit-and-reach flexibility between boys and girls by age and country are presented in Table 4. The size of the sex difference in flexibility differed somewhat by country, but numbers of effects available from most countries were limited. For 5-10-year-olds, the effect sizes from China, Italy, Spain, South Africa, and the USA were broadly similar ( $g = -0.32$  to  $-0.62$ ) with much overlap in their 95% CIs. In Peru, however, the sex difference in sit-and-reach flexibility was smaller than in other countries and did not reach statistical significance ( $g = -0.12$ , 95% CIs  $[-0.31, 0.06]$ ).

For the 14-16-year-olds, few countries had enough effects for robust between-country comparisons. Nevertheless, significant overlap in the 95% CIs existed between the USA, Lithuania, and Poland. China exhibited different results. The mean effect size from China was smaller than these three countries, such that the sex difference in sit-and-reach flexibility among 14-16-year-olds in China was not statistically significant ( $g = -0.73$ , 95% CIs  $[-0.51, 0.03]$ ).

## Discussion

Girls have greater sit-and-reach flexibility than boys at all ages. This sex difference in flexibility is small-to-moderate in size in early childhood, it increases with onset of female puberty, and it returns to levels observed in early childhood by age 16. The size of the sex difference in sit-and-reach flexibility is similar between countries and has remained mostly stable since the 1980s.

### During development

The main novel aspect of the current meta-analysis was the large aggregate sample used to update the question of sex differences in sit-and-reach flexibility in children and adolescents. A total of 96 studies provided 408 effects from over 900,000 boys and girls. This represents a substantial enhancement in statistical power compared to the only other meta-analyses on the topic, which are now 30-40 years old (Thomas & French, 1985; Thomas et al., 1991).

Sit-and-reach data are available from children as young as 3 years old. At this age, girls have greater sit-and-reach flexibility than boys ( $g = -1.35$ ). Nevertheless, this finding should be interpreted cautiously because only one effect was available (Koslow, 1987).

At older ages, confidence in the size of the sex difference in sit-and-reach flexibility is enhanced due to the large number of effects available. Between ages 4-10, girls have a small-to-moderate advantage in sit-and-reach flexibility ( $g = -0.30$  –  $-0.50$ ). Then, at age 11, the effect size

increases ( $g = -0.59$ ), likely due to onset of female puberty (Brix et al., 2019; Euling et al., 2008; Tanner, 1971). The effect size then peaks at age 12 ( $g = -0.78$ ) before narrowing. By age 16, the sex difference in sit-and-reach flexibility returns to pre-puberty levels ( $g = -0.42$ ).

These results are broadly consistent with the results from the two previous meta-analyses (Thomas & French, 1985; Thomas et al., 1991). In their sample of 12,000 boys and girls aged 6-18, Thomas et al. (1991) found greater sit-and-reach flexibility in girls than boys at all ages. They reported an effect size of about  $d = -0.50$  during early childhood, a slight increase in the sex difference at age 11, a peak in the difference at ages 12-13, and a near-return to pre-puberty levels by age 17 (Thomas et al., 1991). However, the peak difference was smaller in the current analysis ( $g = -0.78$ ) than the previous analysis ( $d = -1.1$ ), and though both analyses revealed a narrowing of the difference from ages 15 to 16, the current study findings suggest a return to pre-puberty levels at age 16, whereas the previous analysis found a return to pre-puberty levels at age 17 (Thomas et al., 1991). Another difference between the two studies is that the former study included 6-year-olds as the youngest cohort (Thomas et al., 1991), whereas the current study included samples as young as 3 years old and was thus able to reveal that sex differences in sit-and-reach flexibility occur earlier than previously known.

### **Across time**

A second novel aspect of the current work was the secular analysis. This analysis revealed that the sex difference in sit-and-reach flexibility has remained fairly constant since the 1980s, as evident by the overlap in the 95% CIs across decades. This was true for both 5-10-year-olds and 14-16-year-olds. The overall lack of change in the sex difference in sit-and-reach flexibility over time suggests little or no secular change in the underlying biological or social/environmental factors that impact performance on the test.

Female and male puberty impact the sex difference in sit-and-reach flexibility (Fig. 1). Consequently, in future research examining changes in sex differences in sit-and-reach flexibility across generations, secular changes in puberty onset warrant consideration. Secular decreases in the age of onset of some pubertal stages have been observed in girls, with results in boys less clear (Euling et al., 2008). If boys and girls experience dissimilar secular changes in puberty onset, this could impact the peaks and troughs of sex differences in sit-and-reach flexibility throughout development.

### **Between countries**

A third novel aspect of the current work was the between-countries analysis. This analysis revealed that the sex difference in sit-and-reach flexibility is similar between countries. For 5-10-year-olds, effect sizes from China, Italy, Spain, and South Africa were broadly similar. One exception is Peru, where the sex difference in sit-and-reach flexibility in 5-10-year-olds is not statistically different. Why a sex difference is not present in Peru is unclear.

For 14-16-year-olds, fewer effects are available. Substantial overlap in the 95% CIs exists between the USA, Lithuania, and Poland. China is an exception, as its sex difference in sit-and-reach flexibility in 14-16-year-olds is not statistically different. Why a sex difference is not present in China is unclear. Moving forward, larger datasets of sit-and-reach flexibility will allow for more precise conclusions regarding between-countries sex differences in sit-and-reach flexibility, particularly among 14-16-year-olds. As the average age at menarche is not the same in all countries (Lei et al., 2021; Saczuk et al., 2018; Wang et al., 2024), differences in puberty patterns between countries warrant consideration in future research that explores sex differences in flexibility around the world.

### **Role of physical activity**

What causes the sex difference in sit-and-reach flexibility? One possible cause is a sex difference in physical activity participation, as participation in some sports and exercise can increase flexibility (Afonso et al., 2021; Bennell et al., 2001; Wang et al., 2023).

Throughout childhood and adolescents, girls report greater interest and more frequent participation in yoga, gymnastics, and dancing – activities known to target or rely heavily on flexibility (Anderson et al., 2017; Centers for Disease Control and Prevention, 2023; Deaner et al., 2012; Harrell et al., 2003; National Federation of State High School Associations, 2024; O'Neill et al., 2011; Woods & U.S. Bureau of Labor Statistics, 2017). Flexibility is also the only component of fitness in which girls rate their self-confidence equal to boys' self-confidence (Klomsten et al., 2004). Girls' greater self-confidence and interest in flexibility-based activities compared to boys likely stems from girls' greater flexibility and joint laxity at early ages (Haley et al., 1986; Jansson et al., 2004; Kondratek et al., 2007; Remvig et al., 2007; Rikken-Bultman et al., 1997). This sex difference in flexibility often appears prior to an age when rigorous exercise training is likely. Moreover, even when boys and girls participate in the same sport, girls still exhibit greater flexibility (Aedo-Munoz et al., 2019; Eisenmann & Malina, 2003; Khan et al., 1997). Thus, sex differences in physical activity participation do not seem to fully explain differences in flexibility between boys and girls.

Instead, girls appear to have a biological propensity for flexibility that occurs early in development. This predisposition, coupled with the fact that activities which require high flexibility levels are viewed by boys and girls as being feminine (Schmalz & Kerstetter, 2006), likely leads girls to participate in flexibility-based activities more often than boys. In adults, interest to have one's flexibility measured and to participate in flexibility programs is greater in women than men (Nuzzo & Deaner, 2023). Women's greater interest in being measured on flexibility and wanting to enhance flexibility likely reflects a desire to display, affirm, accentuate, or receive feedback on an aspect of femininity (Nuzzo & Deaner, 2023).

Though greater female interest and participation in flexibility activities has potential to exacerbate baseline sex differences in flexibility, the sex difference in sit-and-reach flexibility is consistent between ages of 4-10, and the difference modulates with key biological events (e.g., female puberty). Such findings, in combination with results that sex differences in sit-and-reach flexibility are broadly similar across time and place (Fig. 2, 3, Table 4), suggest that sex differences in sit-and-reach flexibility are primarily rooted in biology rather than social or environmental factors. Biological factors that warrant consideration are body anthropometrics and muscle stiffness.

### **Role of body height and limb length**

Body height increases with age in boys and girls. Boys are taller than girls until about age 11, which is when girls become taller than boys (USA data) (Kuczmarski et al., 2002). At age 14, girls' body height begins to plateau, while boys' body height continues to increase (Kuczmarski et al., 2002). At age 16, boys' median body height is significantly greater than girls' median body height (Kuczmarski et al., 2002).

Sex differences in body height probably do not contribute significantly to sex differences in sit-and-reach flexibility because body height (in adults) either does not correlate with sit-and-reach flexibility (Mathews et al., 1957; Mathews et al., 1959) or it correlates weakly in the negative direction (Simoneau, 1998). Leg length (Mathews et al., 1957; Mathews et al., 1959; Wear, 1963) and arm length (Simoneau, 1998) also do not correlate strongly with sit-and-reach flexibility, though individuals with long legs and short arms and trunk are disadvantaged in the traditional sit-and-reach (Hoeger & Hopkins, 1992; Hoeger et al., 1990; Wear, 1963). Nonetheless, sex differences in arm-to-leg ratios, should they exist, are unlikely to fully explain greater sit-and-reach performance in females than males. This is because (a) the finger-to-box distance in the sit-and-reach test accounts for only 13% of the variance in sit-and-reach performance (Hoeger & Hopkins, 1992; Hoeger et al., 1990), (b) girls maintain their flexibility advantage over boys even

when the modified version of the sit-and-reach, which accounts for finger-to-box distance, is performed (Castro-Piñero et al., 2009), (c) the modified and traditional versions of the sit-and-reach correlate similarly with hamstrings extensibility in children and adolescents (Castro-Piñero et al., 2009), and (d) females often have greater ROM than males during joint actions that are not confounded by limb length (Grimston et al., 1993; Kawaguchi et al., 2024; McKay et al., 2017).

### **Roles of body mass and fat**

Body mass increases with age in boys and girls. Girls have lower body masses than boys until about age 11 (Kuczmarski et al., 2002). At age 11, girls begin to weigh more than boys (Kuczmarski et al., 2002). However, at age 14, boys regain their body mass advantage, and by age 16, body mass is significantly greater among boys than girls (Kuczmarski et al., 2002).

Sex differences in body mass and fat probably do not contribute to sex differences in sit-and-reach flexibility. Children and adolescents who are overweight and obese have sit-and-reach flexibility scores equal to their youth counterparts who are not overweight or obese (Aijsafe et al., 2018; Deforche et al., 2003; Woll et al., 2013). Also, though girls and women have higher body fat percentages than boys and men (Garnett et al., 2004; Nelson & Barondess, 1997; Taylor et al., 1997; Wells, 2007), body fatness does not correlate with sit-and-reach performance (Minck et al., 2000). In adult women, body mass also does not significantly correlate with toe touch test scores (Broer & Galles, 1958).

### **Roles of muscle mass and stiffness**

The sit-and-reach test primarily examines hamstrings extensibility, as evident by the moderate correlations between sit-and-reach and passive straight leg test scores (Castro-Piñero et al., 2009; Hartman & Looney, 2003; Mayorga-Vega et al., 2014; Muyor et al., 2014; Patterson et al., 1996). Two determinants of hamstrings extensibility include hamstrings muscle stiffness and pain tolerance to hamstrings stretch. During the passive straight leg test, men exhibit more muscle stiffness and less hip flexion ROM, and men report higher levels of pain from the stretch compared to women (Marshall & Siegler, 2014). Men also appear to report slightly higher levels of discomfort during the sit-and-reach compared to women, though this was not formally assessed with a test of statistical significance (Hui & Yuen, 2000).

Thus, in adults, lower sit-and-reach flexibility in males than females can be explained partly by greater hamstrings muscle stiffness and lower pain tolerance to hamstrings stretch in males (Blackburn et al., 2009; Blackburn et al., 2004; Gajdosik et al., 1990; Granata et al., 2002; Marshall & Siegler, 2014; McHugh et al., 1992). Lesser hip ROM on the passive straight leg test

among boys than girls (Castro-Piñero et al., 2009; Hartman & Looney, 2003; Muyor et al., 2014; Patterson et al., 1996) suggests that sex differences in hamstrings muscle stiffness and pain tolerance probably also play a role in explaining sex differences in sit-and-reach flexibility in children and adolescents.

In adults, the cause of the sex difference in hamstrings muscle stiffness is believed to be greater hamstrings muscle cross-sectional area in men than women (Blackburn et al., 2004; Gajdosik et al., 1990; Nuzzo, 2023a). Blackburn et al. (2004) speculated that greater muscle mass in men than women might be due to (a) men having a greater amount of connective tissue that would be resistant to stretch, and (b) men having a greater number of cross-bridges available for spontaneous re-attachment during stretch.

Compared to pre-pubertal girls, pre-pubertal boys have more fat-free or lean mass, less fat mass, and lower body fat percentages (Arfaei et al., 2002; Garnett et al., 2004; He et al., 2002; Leppänen et al., 2017; McCarthy et al., 2014; Nelson & Barondess, 1997; Soininen et al., 2018; Taylor et al., 1997). Thus, prior to puberty, greater hamstrings muscle mass in boys could play a role in their lower flexibility. During male puberty, boys experience a 20-30-fold increase in testosterone and a substantial increase in muscle mass (Courant et al., 2010; Elmlinger et al., 2005; Handelsman et al., 2018; Khairullah et al., 2014; Round et al., 1999).

Because of the disproportionate increase in muscle mass in boys during and after puberty, one might expect the sex difference in sit-and-reach flexibility to widen. Yet, the sex difference in sit-and-reach flexibility narrows after male puberty (Fig. 1). One possible explanation for the return of the sex difference in sit-and-reach flexibility to pre-puberty levels by age 16 is the sex-specific change in lumbar flexion that occurs during development. Hamstrings extensibility is largely responsible for sit-and-reach performance, but sit-and-reach performance is also weakly correlated with lumbar extensibility (Hartman & Looney, 2003; Mayorga-Vega et al., 2014; Patterson et al., 1996). Studies on spinal mobility in boys and girls have reported somewhat mixed results, but overall, they appear to show greater lumbar flexion ROM among girls than boys in early childhood (Haley et al., 1986; Kondratek et al., 2007; Penha et al., 2008), followed by greater lumbar flexion among boys than girls after about age 14 (Oksanen & Salminen, 1996; Sullivan et al., 1994; van Adrichem & van der Korst, 1973). Kondratek et al. (2007) speculated that girls' decline in lumbar flexion ROM at a younger age compared to boys could be due to earlier maturation in girls than boys. Thus, greater male than female trunk flexion ROM after puberty might help explain why the sex difference in sit-and-reach flexibility decreases from peak levels observed at age 12.

To the extent that girls have greater spinal mobility than boys in early development, this may be due, in part, to smaller vertebral cross-sectional areas in girls. Newborn girls have significantly smaller vertebral cross-sectional areas than newborn boys (Gilsanz et al., 2018; Ponrartana et al., 2015). This reduced vertebral size is thought to be “programmed” and representative of a female adaptation to the necessity of fetal load (Gilsanz et al., 2018; Ponrartana et al., 2015). Smaller vertebral cross-sectional areas in children and adolescents correlate with greater lumbar lordosis (Wren et al., 2017), and pre-pubertal girls have smaller vertebra and less paraspinous muscles than boys, with 22% of the difference in vertebral cross-sectional area explained by sex-related differences in paraspinous musculature (Arfaei et al., 2002). Thus, smaller vertebral cross-sectional areas and paraspinous musculature in girls than boys might be other biological factors that contribute to sex differences in sit-and-reach flexibility.

### Limitations

The current study has limitations. First, the literature search did not follow a formal flow diagram. Consequently, study replication will be challenging. However, the aim of the research was to update two meta-analyses that are now 30-40 years old (Thomas & French, 1985; Thomas et al., 1991). This aim was accomplished, as the current analysis includes 408 effects from 96 studies with over 900,000 participants.

Second, only studies published in English were included in the analysis. Consequently, sit-and-reach flexibility data acquired in non-English speaking countries and published in non-English journals were not accounted for in this analysis. Such data could have improved the confidence in effects observed in non-English speaking countries where only a small number of effects were available in English.

Third, the current study did not explicitly follow PRISMA guidelines (Page et al., 2021). For example, the literature search, article screening, and data extraction were conducted by a single author. This may have increased the likelihood of error or bias in these processes. Nevertheless, I have used these strategies successfully in previous reviews and meta-analyses, including papers that have summarized results from more papers than previous identified for certain topics (Nuzzo, 2023b; Nuzzo, Pinto, & Nosaka, 2023; Nuzzo, Pinto, Nosaka, et al., 2023; Nuzzo et al., 2024). Here, the number of relevant studies and effects identified is substantially greater than in the two most relevant previous attempts (Thomas & French, 1985; Thomas et al., 1991). Any papers that might have been missed in the search, or any errors that might have been made in data extraction or processing due to the single-author approach, are unlikely to impact the

overall results, in part, because the large number effects identified have provided effect sizes of high statistical confidence. The data spreadsheet and output have been made publicly available to other researchers who wish to build on this work or re-examine it.

Fourth, the current study was only concerned with sit-and-reach flexibility. Therefore, the results are specific to sex differences in hamstrings extensibility, and to a lesser extent, lumbar extensibility (Castro-Piñero et al., 2009; Chillón et al., 2010; Hartman & Looney, 2003; Mayorga-Vega et al., 2014; Muyor et al., 2014; Patterson et al., 1996). The results are not directly applicable to other muscles or joint actions.

Fifth, the cause of the sex difference in sit-and-reach flexibility was not assessed. Converging lines of evidence suggest that the sex difference in sit-and-reach flexibility before, during, and after puberty is caused primarily by biological factors. Nevertheless, longitudinal studies that examine factors such as muscle stiffness, tolerance to stretch, muscle and vertebral cross-sectional areas, and physical activity participation will be required to identify the causes of the sex difference in sit-and-reach flexibility in children and adolescents.

Sixth, the current study says nothing about the validity of sit-and-reach flexibility as a health and fitness outcome. In 2012, the Institute of Medicine recommended that tests of flexibility not be included in youth fitness testing because of a “lack of evidence for an association between flexibility tests and health outcomes...” (Institute of Medicine, 2012). Similarly, in 2020, a review on flexibility and the sit-and-reach test concluded that flexibility should be de-emphasized as a major component of fitness due to a lack of robust, consistent evidence of an association between sit-and-reach flexibility and important health and performance outcomes in adults (Nuzzo, 2020). The current meta-analysis does not resolve this debate. However, sit-and-reach flexibility appears to be the only proposed fitness attribute in which girls score higher than boys (Thomas & French, 1985; Thomas et al., 1991), which suggests that flexibility is not tightly linked with important fitness attributes (e.g. muscle strength, cardiovascular endurance), and evidence of this lack of an association in adults has been summarized previously (Nuzzo, 2020).

## Conclusion

Girls have greater sit-and-reach flexibility than boys at all ages. Between 3-10 years old, the sex difference in sit-and-reach flexibility is fairly consistent and moderate in size. This difference widens at age 11, peaks at age 12, and then narrows such that by age 16 it equals pre-puberty levels. The sex difference in sit-and-reach flexibility is similar between countries and has remained largely unchanged since the 1980s. Overall, the results confirm that sexual



dimorphism in flexibility occurs early in human development, is impacted by puberty, and is broadly consistent across time and place.

## **Data availability statement**

The data spreadsheet and statistical results associated with this study are available at the Open Science Framework (<https://osf.io/yjexh/>).

## **Discloser and funding statement**

I have no conflicts of interest to report.

## **Funding statement**

No funding was received for this research.

## **Ethics approval**

Ethical approval is not required for a meta-analysis of published data.

## **ORCID**

James L. Nuzzo: <https://orcid.org/0000-0001-9081-0522>

## REFERENCES

- Aedo-Munoz, E., Araya-Ibacache, M., Miarka, B., Moya-Jofre, C., Cancino-Lopez, J., Mozer, R. L., & Brito, C. J. (2019). Effect of sex differences in sports groups on hamstring flexibility based on the sit- reach test: new parameters for Chilean athletes. *Journal of Physical Education and Sport*, 19(4), 2374-2378.
- Afonso, J., Ramirez-Campillo, R., Moscão, J., Rocha, T., Zacca, R., Martins, A., Milheiro, A. A., Ferreira, J., Sarmiento, H., & Clemente, F. M. (2021). Strength Training versus Stretching for Improving Range of Motion: A Systematic Review and Meta-Analysis. *Healthcare*, 9(4). <https://doi.org/10.3390/healthcare9040427>
- Aijsafe, T., Garcia, T., & Fanchiang, H. (2018). Musculoskeletal fitness measures are not created equal: an assessment of school children in Corpus Christi, Texas. *Frontiers in Public Health*, 6(14).
- Amado-Pacheco, J. C., Prieto-Benavides, D. H., Correa-Bautista, J. E., García-Hermoso, A., Agostinis-Sobrinho, C., Alonso-Martínez, A. M., Izquierdo, M., & Ramírez-Vélez, R. (2019). Feasibility and Reliability of Physical Fitness Tests among Colombian Preschool Children. *International Journal of Environmental Research and Public Health*, 16(17). <https://doi.org/10.3390/ijerph16173069>
- Anderson, S. D., Darkings Leyland, S., & Ling, J. (2017). Gender differences in motivation for participation in extra-curricular dance: application of the Theory of Planned Behaviour. *Research in Dance Education*, 18(2), 150-160.
- Anselma, M., Collard, D. C. M., van Berkum, A., Twisk, J. W. R., Chinapaw, M. J. M., & Altenburg, T. M. (2020). Trends in Neuromotor Fitness in 10-to-12-Year-Old Dutch Children: A Comparison Between 2006 and 2015/2017. *Frontiers in Public Health*, 8, 559485. <https://doi.org/10.3389/fpubh.2020.559485>
- Arfai, K., Pitukcheewanont, P. D., Goran, M. I., Tavaré, C. J., Heller, L., & Gilsanz, V. (2002). Bone, muscle, and fat: sex-related differences in prepubertal children. *Radiology*, 224(2), 338-344. <https://doi.org/10.1148/radiol.2242011369>
- Armstrong, M. E., Lambert, E. V., & Lambert, M. I. (2011). Physical fitness of South African primary school children, 6 to 13 years of age: discovery vitality health of the nation study. *Perceptual and Motor Skills*, 113(3), 999-1016. <https://doi.org/10.2466/06.10.13.Pms.113.6.999-1016>
- Ayán Pérez, C., Álvarez Pérez, S., González Baamonde, S., & Martínez de Quel, Ó. (2020). Influence of the Box Dimensions on the Reliability and Validity of the Sit and Reach in Preschoolers. *Journal of Strength and Conditioning Research*, 34(9), 2683-2692. <https://doi.org/10.1519/jsc.0000000000002886>
- Aydin, O., & Yassikaya, M. Y. (2022). Validity and Reliability Analysis of the PlotDigitizer Software Program for Data Extraction from Single-Case Graphs. *Perspectives on*

- Behavior Science*, 45, 239-257. <https://doi.org/https://doi.org/10.1007/s40614-021-00284-0>
- Bae, J. Y., Jang, K. S., Kang, S., Han, D. H., Yang, W., & Shin, K. O. (2015). Correlation between basic physical fitness and pulmonary function in Korean children and adolescents: a cross-sectional survey. *Journal of Physical Therapy Science*, 27(9), 2687-2692. <https://doi.org/10.1589/jpts.27.2687>
- Baquet, G., Twisk, J. W., Kemper, H. C., Van Praagh, E., & Berthoin, S. (2006). Longitudinal follow-up of fitness during childhood: interaction with physical activity. *American Journal of Human Biology*, 18(1), 51-58. <https://doi.org/10.1002/ajhb.20466>
- Bartkowiak, S., Konarski, J. M., Strzelczyk, R., Janowski, J., Karpowicz, M., & Malina, R. M. (2021). Physical Fitness of Rural Polish School Youth: Trends Between 1986 and 2016. *Journal of Physical Activity and Health*, 18(7), 789-800. <https://doi.org/10.1123/jpah.2020-0712>
- Batez, M., Milošević, Ž., Mikulić, I., Sporiš, G., Mačak, D., & Trajković, N. (2021). Relationship between Motor Competence, Physical Fitness, and Academic Achievement in Young School-Aged Children. *BioMed Research International*, 2021, 6631365. <https://doi.org/10.1155/2021/6631365>
- Bennell, K. L., Khan, K. M., Matthews, B. L., & Singleton, C. (2001). Changes in hip and ankle range of motion and hip muscle strength in 8-11 year old novice female ballet dancers and controls: a 12 month follow up study. *British Journal of Sports Medicine*, 35(1), 54-59. <https://doi.org/10.1136/bjsm.35.1.54>
- Benzo, R. M., St George, S. M., Messiah, S. E., Lovan, P., Leite, R., Patel, A., Lee, T. K., & Prado, G. J. (2023). Physical Fitness Among Adolescents Who are Hispanic With Overweight or Obesity. *Pediatric Physical Therapy*, 35(2), 252-258. <https://doi.org/10.1097/pep.0000000000000997>
- Blackburn, J. T., Bell, D. R., Norcross, M. F., Hudson, J. D., & Kimsey, M. H. (2009). Sex comparison of hamstring structural and material properties. *Clinical Biomechanics*, 24(1), 65-70. <https://doi.org/10.1016/j.clinbiomech.2008.10.001>
- Blackburn, J. T., Riemann, B. L., Padua, D. A., & Guskiewicz, K. M. (2004). Sex comparison of extensibility, passive, and active stiffness of the knee flexors. *Clinical Biomechanics*, 19(1), 36-43. <https://doi.org/10.1016/j.clinbiomech.2003.09.003>
- Brix, N., Ernst, A., Lauridsen, L. L. B., Parner, E., Støvring, H., Olsen, J., Henriksen, T. B., & Ramlau-Hansen, C. H. (2019). Timing of puberty in boys and girls: A population-based study. *Pediatric and Perinatal Epidemiology*, 33(1), 70-78. <https://doi.org/10.1111/ppe.12507>
- Broer, M. R., & Galles, N. R. G. (1958). Importance of Relationship Between Various Body Measurements in Performance of the Toe-Touch Test. *Research Quarterly*, 29(3), 253-263. <https://doi.org/10.1080/10671188.1958.10762018>

- Brown, G. A., O'Connor, M. I., & Parker, M. G. (2024). Comments on Sports Participation and Transgender Youths. *JAMA Pediatrics*, 178(3), 315. <https://doi.org/10.1001/jamapediatrics.2023.5960>
- Casajús, J. A., Leiva, M. T., Villarroya, A., Legaz, A., & Moreno, L. A. (2007). Physical performance and school physical education in overweight Spanish children. *Annals of Nutrition & Metabolism*, 51(3), 288-296. <https://doi.org/10.1159/000105459>
- Castro-Piñero, J., Chillón, P., Ortega, F. B., Montesinos, J. L., Sjöström, M., & Ruiz, J. R. (2009). Criterion-related validity of sit-and-reach and modified sit-and-reach test for estimating hamstring flexibility in children and adolescents aged 6-17 years. *International Journal of Sports Medicine*, 30(9), 658-662. <https://doi.org/10.1055/s-0029-1224175>
- Centers for Disease Control and Prevention. (2023). QuickStats: Percentage\* of Children and Adolescents Aged 4-17 Years Who Practiced Yoga During the Past 12 Months,(†) by Sex and Age Group - National Health Interview Survey,(§) United States, 2022. *MMWR. Morbidity and Mortality Weekly Report*, 72(46), 1279. <https://doi.org/10.15585/mmwr.mm7246a6>
- Chagas, D. D. V., & Barnett, L. M. (2023). Adolescents' Flexibility Can Affect Motor Competence: The Pathway from Health Related Physical Fitness to Motor Competence. *Perceptual and Motor Skills*, 130(1), 94-111. <https://doi.org/10.1177/00315125221128638>
- Chen, G., Chen, J., Liu, J., Hu, Y., & Liu, Y. (2022). Relationship between body mass index and physical fitness of children and adolescents in Xinjiang, China: a cross-sectional study. *BMC Public Health*, 22(1), 1680. <https://doi.org/10.1186/s12889-022-14089-6>
- Chillón, P., Castro-Piñero, J., Ruiz, J. R., Soto, V. M., Carbonell-Baeza, A., Dafos, J., Vicente-Rodríguez, G., Castillo, M. J., & Ortega, F. B. (2010). Hip flexibility is the main determinant of the back-saver sit-and-reach test in adolescents. *Journal of Sports Sciences*, 28(6), 641-648.
- Cieśła, E., Mleczko, E., Bergier, J., Markowska, M., & Nowak-Starz, G. (2014). Health-Related Physical Fitness, BMI, physical activity and time spent at a computer screen in 6 and 7-year-old children from rural areas in Poland. *Annals of Agricultural and Environmental Medicine*, 21(3), 617-621. <https://doi.org/10.5604/12321966.1120613>
- Cieśła, E., Zareba, M., & Koziel, S. (2017). The level of physical fitness in children aged 6–7 years with low birthweight. *Early Human Development*, 11, 23-39.
- Courant, F., Aksglaede, L., Antignac, J. P., Monteau, F., Sorensen, K., Andersson, A. M., Skakkebaek, N. E., Juul, A., & Bizec, B. L. (2010). Assessment of circulating sex steroid levels in prepubertal and pubertal boys and girls by a novel ultrasensitive gas chromatography-tandem mass spectrometry method. *Journal of Clinical Endocrinology and Metabolism*, 95(1), 82-92. <https://doi.org/10.1210/jc.2009-1140>

- Davis, D. S., Quinn, R. O., Whiteman, C. T., Williams, J. D., & Young, C. R. (2008). Concurrent validity of four clinical tests used to measure hamstring flexibility. *Journal of Strength and Conditioning Research*, 22(2), 583-588.
- De la Cruz-Sánchez, E., & Pino-Ortega, J. (2010). An active lifestyle explains sex differences in physical performance in children before puberty. *Collegium Antropologicum*, 34(2), 487-491.
- Deaner, R. O., Geary, D. C., Puts, D. A., Ham, S. A., Kruger, J., Fles, E., Winegard, B., & Grandis, T. (2012). A sex difference in the predisposition for physical competition: males play sports much more than females even in the contemporary U.S. *PLoS One*, 7(11), e49168. <https://doi.org/10.1371/journal.pone.0049168>
- Deforche, B., Lefevre, J., De Bourdeaudhuij, I., Hills, A. P., Duquet, W., & Bouckaert, J. (2003). Physical fitness and physical activity in obese and nonobese Flemish youth. *Obesity Research*, 11(3), 434-441. <https://doi.org/10.1038/oby.2003.59>
- Docherty, D., & Bell, R. D. (1985). The relationships between flexibility and linearity measures in boys and girls 6-15 years of age. *Journal of Human Movement Studies*, 11, 279-288.
- Eather, N., Morgan, P. J., & Lubans, D. R. (2016). Improving health-related fitness in adolescents: the CrossFit Teens™ randomised controlled trial. *Journal of Sports Sciences*, 34(3), 209-223. <https://doi.org/10.1080/02640414.2015.1045925>
- Eisenmann, J. C., & Malina, R. M. (2003). Age- and sex-associated variation in neuromuscular capacities of adolescent distance runners. *Journal of Sports Sciences*, 21(7), 551-557. <https://doi.org/10.1080/0264041031000101845>
- Elmlinger, M. W., Kühnel, W., Wormstall, H., & Döller, P. C. (2005). Reference intervals for testosterone, androstenedione and SHBG levels in healthy females and males from birth until old age. *Clinical Laboratory*, 51(11-12), 625-632.
- Estivaleti, J. M. O., Bergamo, R. R., Oliveira, L. C., Beltran, D. C. G., Silva Junior, J. P. D., Santos, M. D., & Matsudo, V. K. R. (2022). Physical activity level measured by accelerometry and physical fitness of schoolchildren. *Rev Paul Pediatr*, 41, e2021230. <https://doi.org/10.1590/1984-0462/2023/41/2021230>
- Euling, S. Y., Herman-Giddens, M. E., Lee, P. A., Selevan, S. G., Juul, A., Sørensen, T. I., Dunkel, L., Himes, J. H., Teilmann, G., & Swan, S. H. (2008). Examination of US puberty-timing data from 1940 to 1994 for secular trends: panel findings. *Pediatrics*, 121 Suppl 3, S172-191. <https://doi.org/10.1542/peds.2007-1813D>
- Fang, H., Quan, M., Zhou, T., Sun, S., Zhang, J., Zhang, H., Cao, Z., Zhao, G., Wang, R., & Chen, P. (2017). Relationship between Physical Activity and Physical Fitness in Preschool Children: A Cross-Sectional Study. *BioMed Research International*, 2017, 9314026. <https://doi.org/10.1155/2017/9314026>
- Fiori, F., Bravo, G., Parpinel, M., Messina, G., Malavolta, R., & Lazzer, S. (2021). A 3-year school-based intervention improved physical fitness and reduced the prevalence of

- overweight and obesity in Italian prepubertal children. *Journal of Sports Medicine and Physical Fitness*, 61(12), 1682-1689. <https://doi.org/10.23736/s0022-4707.21.12011-0>
- Flanagan, S. D., Dunn-Lewis, C., Hatfield, D. L., Distefano, L. J., Fragala, M. S., Shoap, M., Gotwald, M., Trail, J., Gomez, A. L., Volek, J. S., Cortis, C., Comstock, B. A., Hooper, D. R., Szivak, T. K., Looney, D. P., DuPont, W. H., McDermott, D. M., Gaudiose, M. C., & Kraemer, W. J. (2015). Developmental differences between boys and girls result in sex-specific physical fitness changes from fourth to fifth grade. *Journal of Strength and Conditioning Research*, 29(1), 175-180. <https://doi.org/10.1519/jsc.0000000000000623>
- Fogelholm, M., Stigman, S., Huisman, T., & Metsämuuronen, J. (2008). Physical fitness in adolescents with normal weight and overweight. *Scandinavian Journal of Medicine and Science in Sports*, 18(2), 162-170. <https://doi.org/10.1111/j.1600-0838.2007.00685.x>
- Fortier, M. D., Katzmarzyk, P. T., Malina, R. M., & Bouchard, C. (2001). Seven-year stability of physical activity and musculoskeletal fitness in the Canadian population. *Medicine and Science in Sports and Exercise*, 33(11), 1905-1911. <https://doi.org/10.1097/00005768-200111000-00016>
- Gajdosik, R. L., Giuliani, C. A., & Bohannon, R. W. (1990). Passive compliance and length of the hamstring muscles of healthy men and women. *Clinical Biomechanics*, 5(1), 23-29. [https://doi.org/10.1016/0268-0033\(90\)90028-5](https://doi.org/10.1016/0268-0033(90)90028-5)
- Garnett, S. P., Högler, W., Blades, B., Baur, L. A., Peat, J., Lee, J., & Cowell, C. T. (2004). Relation between hormones and body composition, including bone, in prepubertal children. *American Journal of Clinical Nutrition*, 80(4), 966-972. <https://doi.org/10.1093/ajcn/80.4.966>
- Gilsanz, V., Wren, T. A. L., Ponrartana, S., Mora, S., & Rosen, C. J. (2018). Sexual Dimorphism and the Origins of Human Spinal Health. *Endocrine Reviews*, 39(2), 221-239. <https://doi.org/10.1210/er.2017-00147>
- Godoy-Cumillaf, A., Bruneau-Chávez, J., Fuentes-Merino, P., Vásquez-Gómez, J., Sánchez-López, M., Álvarez-Bueno, C., & Cervero-Redondo, I. (2020). Reference Values for Fitness Level and Gross Motor Skills of 4-6-Year-Old Chilean Children. *International Journal of Environmental Research and Public Health*, 17(3). <https://doi.org/10.3390/ijerph17030797>
- Golle, K., Granacher, U., Hoffmann, M., Wick, D., & Muehlbauer, T. (2014). Effect of living area and sports club participation on physical fitness in children: a 4 year longitudinal study. *BMC Public Health*, 14, 499. <https://doi.org/10.1186/1471-2458-14-499>
- Gomula, A., & Koziel, S. (2018). Secular trend and social variation in age at menarche among Polish schoolgirls before and after the political transformation. *American Journal of Human Biology*, 30(1). <https://doi.org/10.1002/ajhb.23048>

- Gontarev, S., Kalac, R., Velickovska, L. A., & Zivkovic, V. (2018). Physical fitness reference standards in Macedonian children and adolescents: the MAKFIT study. *Nutricion Hospitalaria*, 35(6), 1275-1286. <https://doi.org/10.20960/nh.1881> (Estándares de referencia de aptitud física en niños y adolescentes de Macedonia: el estudio MAKFIT.)
- Granata, K. P., Wilson, S. E., & Padua, D. A. (2002). Gender differences in active musculoskeletal stiffness. Part I.: Quantification in controlled measurements of knee joint dynamics. *Journal of Electromyography and Kinesiology*, 12(2), 119-126. [https://doi.org/https://doi.org/10.1016/S1050-6411\(02\)00002-0](https://doi.org/https://doi.org/10.1016/S1050-6411(02)00002-0)
- Greenhalgh, T., & Peacock, R. (2005). Effectiveness and efficiency of search methods in systematic reviews of complex evidence: audit of primary sources. *British Medical Journal*, 331(7524), 1064-1065. <https://doi.org/10.1136/bmj.38636.593461.68>
- Grimston, S. K., Nigg, B. M., Hanley, D. A., & Engsberg, J. R. (1993). Differences in ankle joint complex range of motion as a function of age. *Foot Ankle*, 14(4), 215-222. <https://doi.org/10.1177/107110079301400407>
- Gulías-González, R., Sánchez-López, M., Olivas-Bravo, Á., Solera-Martínez, M., & Martínez-Vizcaíno, V. (2014). Physical fitness in Spanish schoolchildren aged 6-12 years: reference values of the battery EUROFIT and associated cardiovascular risk. *Journal of School Health*, 84(10), 625-635. <https://doi.org/10.1111/josh.12192>
- Haapala, E. A., Lintu, N., Väistö, J., Robinson, L. E., Viitasalo, A., Lindi, V., & Lakka, T. A. (2015). Associations of Physical Performance and Adiposity with Cognition in Children. *Medicine and Science in Sports and Exercise*, 47(10), 2166-2174. <https://doi.org/10.1249/mss.0000000000000652>
- Haapala, E. A., Väistö, J., Lintu, N., Tompuri, T., Brage, S., Westgate, K., Ekelund, U., Lampinen, E. K., Sääkslahti, A., Lindi, V., & Lakka, T. A. (2016). Adiposity, physical activity and neuromuscular performance in children. *Journal of Sports Sciences*, 34(18), 1699-1706. <https://doi.org/10.1080/02640414.2015.1134805>
- Haley, S. M., Tada, W. L., & Carmichael, E. M. (1986). Spinal mobility in young children. A normative study. *Physical Therapy*, 66(11), 1697-1703. <https://doi.org/10.1093/ptj/66.11.1697>
- Hamilton, B., Brown, A., Montagner-Moraes, S., Comeras-Chueca, C., Bush, P. G., Guppy, F. M., & Pitsiladis, Y. P. (2024). Strength, power and aerobic capacity of transgender athletes: a cross-sectional study. *British Journal of Sports Medicine*. <https://doi.org/10.1136/bjsports-2023-108029>
- Handelsman, D. J., Hirschberg, A. L., & Bermon, S. (2018). Circulating testosterone as the hormonal basis of sex differences in athletic performance. *Endocrine Reviews*, 39(5), 803-829. <https://doi.org/10.1210/er.2018-00020>
- Hands, B., Larkin, D., Parker, H., Straker, L., & Perry, M. (2009). The relationship among physical activity, motor competence and health-related fitness in 14-year-old



- adolescents. *Scandinavian Journal of Medicine and Science in Sports*, 19(5), 655-663. <https://doi.org/10.1111/j.1600-0838.2008.00847.x>
- Harrell, J. S., Pearce, P. F., Markland, E. T., Wilson, K., Bradley, C. B., & McMurray, R. G. (2003). Assessing physical activity in adolescents: common activities of children in 6ht-8th grades. *Journal of the American Academy of Nurse Practitioners*, 15(4), 170-178.
- Hartman, J. G., & Looney, M. (2003). Norm-Referenced and Criterion-Referenced Reliability and Validity of the Back-Saver Sit-and- Reach. *Measurement in Physical Education and Exercise Science*, 7(2), 71-87.
- Haugen, T., Høigaard, R., & Seiler, S. (2014). Normative data of BMI and physical fitness in a Norwegian sample of early adolescents. *Scandinavian Journal of Public Health*, 42(1), 67-73. <https://doi.org/10.1177/1403494813504502>
- He, H., Pan, L., Du, J., Liu, F., Jin, Y., Ma, J., Wang, L., Jia, P., Hu, Z., & Shan, G. (2019). Muscle fitness and its association with body mass index in children and adolescents aged 7-18 years in China: a cross-sectional study. *BMC Pediatrics*, 19(1), 101. <https://doi.org/10.1186/s12887-019-1477-8>
- He, Q., Horlick, M., Thornton, J., Wang, J., Pierson, R. N., Jr., Heshka, S., & Gallagher, D. (2002). Sex and race differences in fat distribution among Asian, African-American, and Caucasian prepubertal children. *Journal of Clinical Endocrinology and Metabolism*, 87(5), 2164-2170. <https://doi.org/10.1210/jcem.87.5.8452>
- Henriques-Neto, D., Hetherington-Rauth, M., Magalhães, J. P., Correia, I., Júdice, P. B., & Sardinha, L. B. (2022). Physical fitness tests as an indicator of potential athletes in a large sample of youth. *Clinical Physiology and Functional Imaging*, 42(2), 88-95. <https://doi.org/10.1111/cpf.12735>
- Hilton, E. N., & Lundberg, T. R. (2021). Transgender women in the female category of sport: perspectives on testosterone suppression and performance advantage. *Sports Medicine*, 51(2), 199-214. <https://doi.org/10.1007/s40279-020-01389-3>
- Hoeger, W. W. K., & Hopkins, D. R. (1992). A comparison of the sit and reach and the modified sit and reach in the measurement of flexibility in women. *Research Quarterly for Exercise and Sport*, 63(2), 191-195.
- Hoeger, W. W. K., Hopkins, D. R., Button, S., & Palmer, T. A. (1990). Comparing the Sit and Reach with the Modified Sit and Reach in Measuring Flexibility in Adolescents. *Pediatric Exercise Science*, 2(2), 156-162. <https://doi.org/10.1123/pes.2.2.156>
- Holt, J., Holt, L. E., & Pelham, T. W. (1996). Flexibility redefined. In T. Bauer (Ed.), *Biomechanics in Sports XIII* (pp. 170-174). Lakehead University.
- Hong, S. W., & Hamlin, M. J. (2005). Secular trends and contemporary differences in physique and health-related fitness levels of 11-12 year-old South Korean and New Zealand children. *Southeast Asian Journal of Tropical Medicine and Public Health*, 36(5), 1339-1345.

- Hsu, C. Y., Chen, L. S., Chang, I. J., Fang, W. C., Huang, S. W., Lin, R. H., Ueng, S. W., & Chuang, H. H. (2021). Can Anthropometry and Body Composition Explain Physical Fitness Levels in School-Aged Children? *Children*, 8(6).  
<https://doi.org/10.3390/children8060460>
- Huang, Y. C., & Malina, R. M. (2002). Physical activity and health-related physical fitness in Taiwanese adolescents. *Journal of Physiological Anthropology and Applied Human Science*, 21(1), 11-19. <https://doi.org/10.2114/jpa.21.11>
- Hui, S. S., & Yuen, P. Y. (2000). Validity of the modified back-saver sit-and-reach test: a comparison with other protocols. *Medicine and Science in Sports and Exercise*, 32(9), 1655-1659.
- Institute of Medicine. (2012). *Fitness measures and health outcomes in youth*. The National Academic Press.
- Jackson, A., & Langford, N. J. (1989). The criterion-related validity of the sit and reach test: replication and extension of previous findings. *Research Quarterly for Exercise and Sport*, 60(4), 384-387.
- Jansson, A., Saartok, T., Werner, S., & Renström, P. (2004). General joint laxity in 1845 Swedish school children of different ages: age- and gender-specific distributions. *Acta Paediatrica*, 93(9), 1202-1206. <https://doi.org/10.1080/08035250410023971>
- Jones, M. A., Stratton, G., Reilly, T., & Unnithan, V. B. (2005). Biological risk indicators for recurrent non-specific low back pain in adolescents. *British Journal of Sports Medicine*, 39(3), 137-140. <https://doi.org/10.1136/bjism.2003.009951>
- Jürimäe, T., & Saar, M. (2003). Self-perceived and actual indicators of motor abilities in children and adolescents. *Perceptual and Motor Skills*, 97(3 Pt 1), 862-866.  
<https://doi.org/10.2466/pms.2003.97.3.862>
- Jürimäe, T., & Volbekiene, V. (1998). Eurofit Test Results in Estonian and Lithuanian 11 to 17-Year-Old Children: A Comparative Study. *European Journal of Physical Education*, 3(2), 178-184.
- Kamandulis, S., Emeljanovas, A., & Skurvydas, A. (2013). Stretching exercise volume for flexibility enhancement in secondary school children. *Journal of Sports Medicine and Physical Fitness*, 53(6), 687-692.
- Karppanen, A. K., Ahonen, S. M., Tammelin, T., Vanhala, M., & Korpelainen, R. (2012). Physical activity and fitness in 8-year-old overweight and normal weight children and their parents. *International Journal of Circumpolar Health*, 71, 17621.  
<https://doi.org/10.3402/ijch.v71i0.17621>
- Katzmarzyk, P. T., Pérusse, L., Rao, D. C., & Bouchard, C. (2000). Familial risk ratios for high and low physical fitness levels in the Canadian population. *Medicine and Science in Sports and Exercise*, 32(3), 614-619. <https://doi.org/10.1097/00005768-200003000-00010>

- Kawaguchi, D., Ito, T., Yamamoto, Y., Fukaya, Y., Mizusawa, J., Ochi, N., Tomita, H., & Noritake, K. (2024). Developmental changes in lower limb joint ranges of motion in Japanese children aged 6–15 years. *Journal of Orthopaedics*.  
<https://doi.org/https://doi.org/10.1016/j.jor.2024.07.004>
- Khairullah, A., Klein, L. C., Ingle, S. M., May, M. T., Whetzel, C. A., Susman, E. J., & Paus, T. (2014). Testosterone trajectories and reference ranges in a large longitudinal sample of male adolescents. *PLoS One*, 9(9), e108838.  
<https://doi.org/10.1371/journal.pone.0108838>
- Khan, K., Roberts, P., Nattrass, C., Bennell, K., Mayes, S., Way, S., Brown, J., McMeeken, J., & Wark, J. (1997). Hip and ankle range of motion in elite classical ballet dancers and controls. *Clinical Journal of Sports Medicine*, 7(3), 174-179.
- Kidokoro, T., Tanaka, H., Naoi, K., Ueno, K., Yanaoka, T., Kashiwabara, K., & Miyashita, M. (2016). Sex-specific associations of moderate and vigorous physical activity with physical fitness in adolescents. *European Journal of Sport Science*, 16(8), 1159-1166.  
<https://doi.org/10.1080/17461391.2016.1183050>
- Kim, O. H., & Park, J. K. (2017). Effects of an Integrated Health Care Program for Children. *Journal of Exercise Nutrition and Biochemistry*, 21(1), 7-12.  
<https://doi.org/10.20463/jenb.2016.0037>
- Klomsten, A. T., Skaalvik, E. M., & Espnes, G. A. (2004). Physical self-concept and sports: do gender differences still exist? *Sex Roles*, 50(1/2), 119-127.
- Knudson, D. V., Magnusson, P., & McHugh, M. (2000). Current issues in flexibility fitness. *President's Council on Physical Fitness and Sports Research Digest*, 3(19).
- Kondratak, M., Krauss, J., Stiller, C., & Olson, R. (2007). Normative values for active lumbar range of motion in children. *Pediatric Physical Therapy*, 19(3), 236-244.  
<https://doi.org/10.1097/PEP.0b013e3180f61c43>
- Kondric, M., Trajkovski, B., Strbad, M., Foretić, N., & Zenić, N. (2013). Anthropometric influence on physical fitness among preschool children: gender-specific linear and curvilinear regression models. *Collegium Antropologicum*, 37(4), 1245-1252.
- Koslow, R. E. (1987). Sit and reach flexibility measures for boys and girls aged three through eight years. *Perceptual and Motor Skills*, 64, 1103-1106.
- Kuczarski, R. J., Ogden, C. L., & Guo, S. S. (2002). 2000 CDC growth charts for the United States: Methods and development. *Vital Health Statistics*, 11(246).
- Lehnhard, H. R., Lehnhard, R. A., Butterfield, S. A., Beckwith, D. M., & Marion, S. F. (1992). Health-related physical fitness levels of elementary school children ages 5-9. *Perceptual and Motor Skills*, 75(3 Pt 1), 819-826.  
<https://doi.org/10.2466/pms.1992.75.3.819>
- Lei, Y., Luo, D., Yan, X., Zhang, J., Hu, P., Ma, J., Song, Y., & Lau, P. W. C. (2021). The mean age of menarche among Chinese schoolgirls declined by 6 months from 2005 to 2014. *Acta Paediatrica*, 110(2), 549-555. <https://doi.org/10.1111/apa.15441>

- Leppänen, M. H., Henriksson, P., Delisle Nyström, C., Henriksson, H., Ortega, F. B., Pomeroy, J., Ruiz, J. R., Cadenas-Sanchez, C., & Löf, M. (2017). Longitudinal Physical Activity, Body Composition, and Physical Fitness in Preschoolers. *Medicine and Science in Sports and Exercise*, 49(10), 2078-2085.  
<https://doi.org/10.1249/mss.0000000000001313>
- Liemohn, W., Sharpe, G. L., & Wassermann, J. F. (1994). Criterion related validity of the sit-and-reach test. *Journal of Strength and Conditioning Research*, 8(2), 91-94.
- Lintu, N., Savonen, K., Viitasalo, A., Tompuri, T., Paananen, J., Tarvainen, M. P., & Lakka, T. (2016). Determinants of Cardiorespiratory Fitness in a Population Sample of Girls and Boys Aged 6 to 8 Years. *Journal of Physical Activity and Health*, 13(11), 1149-1155.  
<https://doi.org/10.1123/jpah.2015-0644>
- Lo, K. Y., Wu, M. C., Tung, S. C., Hsieh, C. C., Yao, H. H., & Ho, C. C. (2017). Association of School Environment and After-School Physical Activity with Health-Related Physical Fitness among Junior High School Students in Taiwan. *International Journal of Environmental Research and Public Health*, 14(1).  
<https://doi.org/10.3390/ijerph14010083>
- Lovecchio, N., Giuriato, M., Zago, M., & Nevill, A. (2019). Identifying the optimal body shape and composition associated with strength outcomes in children and adolescent according to place of residence: An allometric approach. *Journal of Sports Sciences*, 37(12), 1434-1441. <https://doi.org/10.1080/02640414.2018.1562615>
- Lovecchio, N., Novak, D., Eid, L., Casolo, F., & Podnar, H. (2015). Urban and rural fitness level: comparison between Italian and Croatian students. *Perceptual and Motor Skills*, 120(2), 367-380. <https://doi.org/10.2466/06.PMS.120v11x8>
- Lovecchio, N., Novak, D., Sedlacek, J., Hamar, P., Milanovic, I., Radisavljevic-Janic, S., Emeljanovas, A., Eid, L., & Zago, M. (2019). Physical fitness for sedentary students: a common trend from six European countries. *Journal of Sports Medicine and Physical Fitness*, 59(8), 1389-1396. <https://doi.org/10.23736/s0022-4707.18.08926-0>
- Lundberg, T. R., Tucker, R., McGawley, K., Williams, A. G., Millet, G. P., Sandbakk, Ø., Howatson, G., Brown, G. A., Carlson, L. A., Chantler, S., Chen, M. A., Heffernan, S. M., Heron, N., Kirk, C., Murphy, M. H., Pollock, N., Pringle, J., Richardson, A., Santos-Concejero, J., Stebbings, G. K., Christiansen, A. V., Phillips, S. M., Devine, C., Jones, C., Pike, J., & Hilton, E. N. (2024). The International Olympic Committee framework on fairness, inclusion and nondiscrimination on the basis of gender identity and sex variations does not protect fairness for female athletes. *Scandinavian Journal of Medicine and Science in Sports*, 34(3), e14581. <https://doi.org/10.1111/sms.14581>
- Mačak, D., Popović, B., Babić, N., Cadenas-Sanchez, C., Madić, D. M., & Trajković, N. (2022). The effects of daily physical activity intervention on physical fitness in preschool children. *Journal of Sports Sciences*, 40(2), 146-155.  
<https://doi.org/10.1080/02640414.2021.1978250>

- Marques, A., Henriques-Neto, D., Peralta, M., Martins, J., Gomes, F., Popovic, S., Masanovic, B., Demetriou, Y., Schlund, A., & Ihle, A. (2021). Field-Based Health-Related Physical Fitness Tests in Children and Adolescents: A Systematic Review. *Frontiers in Pediatrics*, 9, 640028. <https://doi.org/10.3389/fped.2021.640028>
- Marshall, P. W., & Siegler, J. C. (2014). Lower hamstring extensibility in men compared to women is explained by differences in stretch tolerance. *BMC Musculoskeletal Disorders*, 15, 223. <https://doi.org/10.1186/1471-2474-15-223>
- Marshall, S. J., Sarkin, J. A., Sallis, J. F., & McKenzie, T. L. (1998). Tracking of health-related fitness components in youth ages 9 to 12. *Medicine and Science in Sports and Exercise*, 30(6), 910-916. <https://doi.org/10.1097/00005768-199806000-00021>
- Martinho, D. V., Gouveia É, R., França, C., Lopes, H., Ihle, A., Marques, A., Rodrigues, A., Alves, R., & Sarmiento, H. (2022). Body Composition and Physical Fitness in Madeira Youth. *Children*, 9(12). <https://doi.org/10.3390/children9121833>
- Mathews, D. K., Shaw, V., & Bohnen, M. (1957). Hip Flexibility of College Women as Related to Length of Body Segments. *Research Quarterly*, 28(4), 352-356. <https://doi.org/10.1080/10671188.1957.10612943>
- Mathews, D. K., Shaw, V., & Woods, J. B. (1959). Hip Flexibility of Elementary School Boys as Related to Body Segments. *Research Quarterly*, 30(3), 297-302. <https://doi.org/10.1080/10671188.1959.10613039>
- Mayorga-Vega, D., Merino-Marban, R., & Viciano, J. (2014). Criterion-related validity of sit-and-reach tests for estimating hamstring and lumbar extensibility: a meta-analysis. *Journal of Sports Science and Medicine*, 13(1), 1-14.
- McCarthy, H. D., Samani-Radia, D., Jebb, S. A., & Prentice, A. M. (2014). Skeletal muscle mass reference curves for children and adolescents. *Pediatric Obesity*, 9(4), 249-259. <https://doi.org/10.1111/j.2047-6310.2013.00168.x>
- McHugh, M. P., Magnusson, S. P., Gleim, G. W., & Nicholas, J. A. (1992). Viscoelastic stress relaxation in human skeletal muscle. *Medicine and Science in Sports and Exercise*, 24(12), 1375-1382.
- McKay, M. J., Baldwin, J. N., Ferreira, P., Simic, M., Vanicek, N., & Burns, J. (2017). Normative reference values for strength and flexibility of 1,000 children and adults. *Neurology*, 88(1), 36-43.
- McMillan, C. S., & Erdmann, L. D. (2010). Tracking adiposity and health-related physical fitness test performances from early childhood through elementary school. *Pediatric Exercise Science*, 22(2), 231-244. <https://doi.org/10.1123/pes.22.2.231>
- Minck, M. R., Ruiters, L. M., Van Mechelen, W., Kemper, H. C., & Twisk, J. W. (2000). Physical fitness, body fatness, and physical activity: The Amsterdam Growth and Health Study. *American Journal of Human Biology*, 12(5), 593-599. [https://doi.org/10.1002/1520-6300\(200009/10\)12:5<593::Aid-ajhb3>3.0.Co;2-u](https://doi.org/10.1002/1520-6300(200009/10)12:5<593::Aid-ajhb3>3.0.Co;2-u)

- Moliner-Urdiales, D., Ortega, F. B., Vicente-Rodriguez, G., Rey-Lopez, J. P., Gracia-Marco, L., Widhalm, K., Sjöström, M., Moreno, L. A., Castillo, M. J., & Ruiz, J. R. (2010). Association of physical activity with muscular strength and fat-free mass in adolescents: the HELENA study. *European Journal of Applied Physiology*, 109(6), 1119-1127. <https://doi.org/10.1007/s00421-010-1457-z>
- Monyeki, M. A., Koppes, L. L., Kemper, H. C., Monyeki, K. D., Toriola, A. L., Pienaar, A. E., & Twisk, J. W. (2005). Body composition and physical fitness of undernourished South African rural primary school children. *European Journal of Clinical Nutrition*, 59(7), 877-883. <https://doi.org/10.1038/sj.ejcn.1602153>
- Muyor, J. M., Zemková, E., Štefániková, G., & Kotyra, M. (2014). Concurrent validity of clinical tests for measuring hamstring flexibility in school age children. *International Journal of Sports Medicine*, 35(8), 664-669.
- National Federation of State High School Associations. (2024). *2022-23 High School Athletics Participation Survey*. <http://www.nfhs.org>
- Nelson, D. A., & Barondess, D. A. (1997). Whole body bone, fat and lean mass in children: comparison of three ethnic groups. *American Journal of Physical Anthropology*, 103(2), 157-162. [https://doi.org/10.1002/\(sici\)1096-8644\(199706\)103:2<157::Aid-ajpa2>3.0.Co;2-r](https://doi.org/10.1002/(sici)1096-8644(199706)103:2<157::Aid-ajpa2>3.0.Co;2-r)
- Nindl, B. C., Alvar, B. A., J. R. D., Favre, M. W., Martin, G. J., Sharp, M. A., Warr, B. J., Stephenson, M. D., & Kraemer, W. J. (2015). Executive Summary From the National Strength and Conditioning Association's Second Blue Ribbon Panel on Military Physical Readiness: Military Physical Performance Testing. *Journal of Strength and Conditioning Research*, 29 Suppl 11, S216-S220. <https://doi.org/10.1519/jsc.0000000000001037>
- Nokoff, N. J., Senefeld, J., Krausz, C., Hunter, S., & Joyner, M. (2023). Sex differences in athletic performance: perspectives on transgender athletes. *Exercise and Sport Sciences Reviews*. <https://doi.org/10.1249/jes.0000000000000317>
- Nuzzo, J. L. (2020). The case for retiring flexibility as a major component of physical fitness. *Sports Medicine*, 50(5), 853-870. <https://doi.org/10.1007/s40279-019-01248-w>
- Nuzzo, J. L. (2023a). Narrative review of sex differences in muscle strength, endurance, activation, size, fiber type, and strength training participation rates, preferences, motivations, injuries, and neuromuscular adaptations. *Journal of Strength and Conditioning Research*, 37(2), 494-536. <https://doi.org/10.1519/jsc.0000000000004329>
- Nuzzo, J. L. (2023b). Sex differences in skeletal muscle fiber types: a meta-analysis. *Clinical Anatomy*. <https://doi.org/10.1002/ca.24091>
- Nuzzo, J. L. (2024). PRE-PRINT - Sex differences in grip strength from birth to age 16: a meta-analysis. *SportRxiv*. <https://doi.org/https://doi.org/10.51224/SRXIV.441>

- Nuzzo, J. L., & Deaner, R. O. (2023). Men and women differ in their interest and willingness to participate in exercise and sports science research. *Scandinavian Journal of Medicine and Science in Sports*, 33(9), 1850-1865. <https://doi.org/10.1111/sms.14404>
- Nuzzo, J. L., Pinto, M. D., & Nosaka, K. (2023). Muscle fatigue during maximal eccentric-only, concentric-only, and eccentric-concentric bicep curl exercise with automated drop setting. *Scandinavian Journal of Medicine and Science in Sports*. <https://doi.org/10.1111/sms.14330>
- Nuzzo, J. L., Pinto, M. D., Nosaka, K., & Steele, J. (2023). The eccentric:concentric strength ratio of human skeletal muscle *in vivo*: meta-analysis of the influences of sex, age, joint action, and velocity. *Sports Medicine*, 53(6), 1125-1136. <https://doi.org/10.1007/s40279-023-01851-y>
- Nuzzo, J. L., Pinto, M. D., Nosaka, K., & Steele, J. (2024). Maximal number of repetitions at percentages of the one repetition maximum: a meta-regression and moderator analysis of sex, age, training status, and exercise. *Sports Medicine*, 54(2), 303-321. <https://doi.org/10.1007/s40279-023-01937-7>
- O'Neill, J. R., Pate, R. R., & Liese, A. D. (2011). Descriptive epidemiology of dance participation in adolescents. *Research Quarterly for Exercise and Sport*, 82(3), 373-380. <https://doi.org/10.1080/02701367.2011.10599769>
- Oja, L., & Jürimäe, T. (1997). Assessment of motor ability of 4- and 5-year-old children. *American Journal of Human Biology*, 9(5), 659-664. [https://doi.org/10.1002/\(sici\)1520-6300\(1997\)9:5<659::Aid-ajhb12>3.0.Co;2-l](https://doi.org/10.1002/(sici)1520-6300(1997)9:5<659::Aid-ajhb12>3.0.Co;2-l)
- Oja, L., & Jürimäe, T. (2002). Physical activity, motor ability, and school readiness of 6-yr.-old children. *Perceptual and Motor Skills*, 95(2), 407-415. <https://doi.org/10.2466/pms.2002.95.2.407>
- Oksanen, A., & Salminen, J. J. (1996). Tests of spinal mobility and muscle strength in the young: Reliability and normative values. *Physiotherapy Theory and Practice*, 12(3), 151-160.
- Örjan, E., Kristjan, O., & Björn, E. (2005). Physical performance and body mass index in Swedish children and adolescents. *Scandinavian Journal of Nutrition*, 49(4), 172-179. <https://doi.org/10.1080/11026480500441275>
- Ortega, F. B., Ruiz, J. R., Castillo, M. J., Moreno, L. A., Urzánqui, A., González-Gross, M., Sjörström, M., & Gutiérrez, A. (2008). Health-related physical fitness according to chronological and biological age in adolescents. The AVENA study. *Journal of Sports Medicine and Physical Fitness*, 48(3), 371-379.
- Page, M. J., McKenzie, J. E., Bossuyt, P. M., Boutron, I., Hoffmann, T. C., Mulrow, C. D., Shamseer, L., Tetzlaff, J. M., Akl, E. A., Brennan, S. E., Chou, R., Glanville, J., Grimshaw, J. M., Hróbjartsson, A., Lalu, M. M., Li, T., Loder, E. W., Mayo-Wilson, E., McDonald, S., McGuinness, L. A., Stewart, L. A., Thomas, J., Tricco, A. C., Welch, V. A., Whiting, P., &

- Moher, D. (2021). The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *British Medical Journal*, 372, n71. <https://doi.org/10.1136/bmj.n71>
- Panczyk, W., Drabik, J., & Shapiro, J. L. (2014). Physical fitness levels of Polish children compared to the U.S. Presidential and National Physical Fitness Award standards. *Journal of Kinesiology and Exercise Sciences*, 66(24), 41-48.
- Patterson, P., Wiksten, D. L., Ray, L., Flanders, C., & Sanphy, D. (1996). The validity and reliability of the back saver sit-and-reach test in middle school girls and boys. *Research Quarterly for Exercise and Sport*, 67(4), 448-451.
- Pelicer, F. R., Nagamine, K. K., Faria, M. A., de Lima Freitas, V., Neiva, C. M., Pessôa Filho, D. M., Gomes Ciolac, E., & Lopes Verardi, C. E. (2016). Health-Related Physical Fitness in School Children and Adolescents. *International Journal of Sports Science* 61, 19-24.
- Penha, P. J., Casarotto, R. A., Sacco, I. C. N., Marques, A. P., & João, S. M. A. (2008). Qualitative postural analysis among boys and girls of seven to ten years of age. *Revista Brasileira de Fisioterapia*, 12(5), 386-391.
- Pissanos, B. W., Moore, J. B., & Reeve, T. G. (1983). Age, sex, and body composition as predictors of children's performance on basic motor abilities and health-related fitness items. *Perceptual and Motor Skills*, 56(1), 71-77. <https://doi.org/10.2466/pms.1983.56.1.71>
- Podstawski, R., & Boryslawski, K. (2012). Relationships between selected anthropometric features and motor abilities of children aged 7-9 [Clin Kinesiol]. *Clinical Kinesiology*, 66(4), 81-90.
- Ponrartana, S., Aggabao, P. C., Dharmavaram, N. L., Fisher, C. L., Friedlich, P., Devaskar, S. U., & Gilsanz, V. (2015). Sexual Dimorphism in Newborn Vertebrae and Its Potential Implications. *Journal of Pediatrics*, 167(2), 416-421. <https://doi.org/10.1016/j.jpeds.2015.04.078>
- Puszczalowska-Lizis, E., Kaźmierczak, M., Omorczyk, J., Lizis, S., Lenková, R., & Pieniędz, K. (2023). Sex-related differences in the physical fitness of adolescents aged 16 to 18 years. *Journal of Kinesiology and Exercise Sciences*, 103(33), 1-7. <https://doi.org/10.5604/01.3001.0053.7363>
- Remvig, L., Jensen, D. V., & Ward, R. C. (2007). Epidemiology of general joint hypermobility and basis for the proposed criteria for benign joint hypermobility syndrome: review of the literature. *Journal of Rheumatology*, 34(4), 804-809.
- Reuben, D. B., Magasi, S., McCreath, H. E., Bohannon, R. W., Wang, Y. C., Bubela, D. J., Rymer, W. Z., Beaumont, J., Rine, R. M., Lai, J. S., & Gershon, R. C. (2013). Motor assessment using the NIH Toolbox. *Neurology*, 80, S65-S75.
- Richards, A. B., Klos, L., Swindell, N., Griffiths, L. J., De Martelaer, K., Edwards, L. C., Brophy, S., & Stratton, G. (2022). Associations between swimming & cycling abilities and fitness in 9-11 year old boys and girls. *Journal of Sports Sciences*, 40(6), 658-666. <https://doi.org/10.1080/02640414.2021.2013616>



- Riddoch, C., Savage, J. M., Murphy, N., Cran, G. W., & Boreham, C. (1991). Long term health implications of fitness and physical activity patterns. *Archives of Disease in Childhood*, 66(12), 1426-1433. <https://doi.org/10.1136/adc.66.12.1426>
- Rikken-Bultman, D. G. A., Wellink, L., & van Dongen, P. W. J. (1997). Hypermobility in two Dutch school populations. *European Journal of Obstetrics & Gynecology*, 73, 189-192.
- Round, J. M., Jones, D. A., Honour, J. W., & Nevill, A. M. (1999). Hormonal factors in the development of differences in strength between boys and girls during adolescence: a longitudinal study. *Annals of Human Biology*, 26(1), 49-62. <https://doi.org/10.1080/030144699282976>
- Runhaar, J., Collard, D. C., Singh, A. S., Kemper, H. C., van Mechelen, W., & Chinapaw, M. (2010). Motor fitness in Dutch youth: differences over a 26-year period (1980-2006). *Journal of Science and Medicine in Sport* 13(3), 323-328. <https://doi.org/10.1016/j.jsams.2009.04.006>
- Ruzbarsky, P., Zvonar, M., Kacur, P., Kokinda, M., & Anton, V. (2022). Health-Related Fitness in Slovak High School Students in Prešov Region. *Sustainability*(14). <https://doi.org/10.3390/su14063606>
- Ryu, J. S., Chung, H. R., Meador, B. M., Seo, Y., & Kim, K. O. (2021). The Associations between Physical Fitness, Complex vs Simple Movement, and Academic Achievement in a Cohort of Fourth Graders. *International Journal of Environmental Research and Public Health*, 18(5). <https://doi.org/10.3390/ijerph18052293>
- Sacchetti, R., Ceciliani, A., Garulli, A., Masotti, A., Poletti, G., Beltrami, P., & Leoni, E. (2012). Physical fitness of primary school children in relation to overweight prevalence and physical activity habits. *Journal of Sports Sciences*, 20(7), 633-640.
- Saczuk, J., Wasiluk, A., & Wilczewski, A. (2018). Body height and age at menarche of girls from eastern Poland in the period of political transformation. *Anthropological Review*, 81(2), 130-145. <https://doi.org/10.2478/anre-2018-0010>
- Safrit, M. J., & Wood, T. M. (1987). The Test Battery Reliability of the Health-Related Physical Fitness Test. *Research Quarterly for Exercise and Sport*, 58(2), 160-167. <https://doi.org/10.1080/02701367.1987.10605439>
- Santos, C., Pereira, S., Vasconcelos, O., Tani, G., Hedeker, D., Katzmarzyk, P. T., Maia, J., & Bustamante, A. (2023). Secular Trends in Physical Fitness of Peruvian Children Living at High-Altitude. *International Journal of Environmental Research and Public Health*, 20(4). <https://doi.org/10.3390/ijerph20043236>
- Sasayama, K., & Adachi, M. (2019). Tracking of objective physical activity and physical fitness in Japanese children. *BMC Research Notes*, 12(1), 252. <https://doi.org/10.1186/s13104-019-4288-y>
- Schmalz, D. L., & Kerstetter, D. L. (2006). Girlie Girls and Manly Men: Children's Stigma Consciousness of Gender in Sports and Physical Activities. *Journal of Leisure Research*, 38(4), 536-557.

- Sember, V., Jurak, G., Starc, G., & Morrison, S. A. (2022). Can Primary School Mathematics Performance Be Predicted by Longitudinal Changes in Physical Fitness and Activity Indicators? *Frontiers in Psychology, 13*, 796838. <https://doi.org/10.3389/fpsyg.2022.796838>
- Shephard, R. J., Berridge, M., & Montelpare, W. (1990). On the generality of the "sit and reach" test: an analysis of flexibility data for an aging population. *Research Quarterly for Exercise and Sport, 61*(4), 326-330. <https://doi.org/10.1080/02701367.1990.10607495>
- Siegel, J. A., Camaione, D. N., & Manfredi, T. G. (1989). The Effects of Upper Body Resistance Training on Prepubescent Children. *Pediatric Exercise Science, 1*(2), 145-154. <https://doi.org/10.1123/pes.1.2.145>
- Simoneau, G. G. (1998). The impact of various anthropometric and flexibility measurements on the sit-and-reach test. *Journal of Strength and Conditioning Research, 12*(4), 232-237.
- Smith, J. F., & Miller, C. V. (1985). The effect of head position on sit and reach performance. *Research Quarterly for Exercise and Sport, 56*(1), 84-85.
- Soininen, S., Sidoroff, V., Lindi, V., Mahonen, A., Kröger, L., Kröger, H., Jääskeläinen, J., Atalay, M., Laaksonen, D. E., Laitinen, T., & Lakka, T. A. (2018). Body fat mass, lean body mass and associated biomarkers as determinants of bone mineral density in children 6-8years of age - The Physical Activity and Nutrition in Children (PANIC) study. *Bone, 108*, 106-114. <https://doi.org/10.1016/j.bone.2018.01.003>
- Sokolowski, B., & Chrzanowska, M. (2012). Development of selected motor skills in boys and girls in relation to their rate of maturation - a longitudinal study. *Human Movement, 13*(2), 132-138. <https://doi.org/10.2478/v10038-012-0014-5>
- Sullivan, M. S., Dickinson, C. E., & Troup, J. D. (1994). The influence of age and gender on lumbar spine sagittal plane range of motion. A study of 1126 healthy subjects. *Spine, 19*(6), 682-686. <https://doi.org/10.1097/00007632-199403001-00007>
- Tanner, J. M. (1971). Sequence, Tempo, and Individual Variation in the Growth and Development of Boys and Girls Aged Twelve to Sixteen. *Daedalus, 100*(4), 907-930.
- Taylor, R. W., Gold, E., Manning, P., & Goulding, A. (1997). Gender differences in body fat content are present well before puberty. *International Journal of Obesity and Related Metabolic Disorders, 21*(11), 1082-1084. <https://doi.org/10.1038/sj.ijo.0800522>
- Thomas, J. R., & French, K. E. (1985). Gender differences across age in motor performance a meta-analysis. *Psychological Bulletin, 98*(2), 260-282. <https://doi.org/10.1037/0033-2909.98.2.260>
- Thomas, J. R., & French, K. E. (1987). References for motor tasks--gender differences across age in motor performance: a meta-analysis. *Perceptual and Motor Skills, 64*(2), 503-506. <https://doi.org/10.2466/pms.1987.64.2.503>

- Thomas, J. R., Nelson, J. K., & Church, G. (1991). A developmental analysis of gender differences in health related physical fitness. *Pediatric Exercise Science*, 3, 28-42. <https://doi.org/10.1123/pes.3.1.28>
- Tishukaj, F., Shalaj, I., Gjaka, M., Ademi, B., Ahmetxhekaj, R., Bachl, N., Tschan, H., & Wessner, B. (2017). Physical fitness and anthropometric characteristics among adolescents living in urban or rural areas of Kosovo. *BMC Public Health*, 17(1), 711. <https://doi.org/10.1186/s12889-017-4727-4>
- True, L., Martin, E. M., Pfeiffer, K. A., Siegel, S. R., Branta, C. F., Haubenstricker, J., & Seefeldt, V. (2021). Tracking of Physical Fitness Components from Childhood to Adolescence: A Longitudinal Study. *Measurement in Physical Education and Exercise Science*, 25(1), 22-34.
- Tsimeas, P. D., Tsiokanos, A. L., Koutedakis, Y., Tsigilis, N., & Kellis, S. (2005). Does living in urban or rural settings affect aspects of physical fitness in children? An allometric approach. *British Journal of Sports Medicine*, 39(9), 671-674. <https://doi.org/10.1136/bjism.2004.017384>
- Tsoukos, A., & Bogdanis, G. C. (2021). The Effects of a Five-Month Lockdown Due to COVID-19 on Physical Fitness Parameters in Adolescent Students: A Comparison between Cohorts. *International Journal of Environmental Research and Public Health*, 19(1). <https://doi.org/10.3390/ijerph19010326>
- Tucker, R., Hilton, E. N., McGawley, K., Pollock, N., Millet, G. P., Sandbakk, Ø., Howatson, G., Brown, G. A., Carlson, L. A., Chen, M. A., Heron, N., Kirk, C., Murphy, M. H., Pringle, J., Richardson, A., Santos-Concejero, J., Christiansen, A. V., Jones, C., Alonso, J. M., Robinson, R., Jones, N., Wilson, M., Parker, M. G., Chintoh, A., Hunter, S., Senefeld, J. W., O'Connor, M. I., Joyner, M., Carneiro, E. M., Devine, C., Pike, J., & Lundberg, T. R. (2024). Fair and Safe Eligibility Criteria for Women's Sport. *Scandinavian Journal of Medicine and Science in Sports*, 34(8), e14715. <https://doi.org/10.1111/sms.14715>
- van Adrichem, J. A., & van der Korst, J. K. (1973). Assessment of the flexibility of the lumbar spine. A pilot study in children and adolescents. *Scandinavian Journal of Rheumatology*, 2(2), 87-91. <https://doi.org/10.3109/03009747309098823>
- Vanhelst, J., Labreuche, J., Béghin, L., Drumez, E., Fardy, P. S., Chapelot, D., Mikulovic, J., & Ulmer, Z. (2017). Physical Fitness Reference Standards in French Youth: The BOUGE Program. *Journal of Strength and Conditioning Research*, 31(6), 1709-1718. <https://doi.org/10.1519/jsc.0000000000001640>
- Veraksa, A., Tvardovskaya, A., Gavrilo, M., Yakupova, V., & Musálek, M. (2021). Associations Between Executive Functions and Physical Fitness in Preschool Children. *Frontiers in Psychology*, 12, 674746. <https://doi.org/10.3389/fpsyg.2021.674746>

- Vitali, F., Robazza, C., Bortoli, L., Bertinato, L., Schena, F., & Lanza, M. (2019). Enhancing fitness, enjoyment, and physical self-efficacy in primary school children: a DEDIPAC naturalistic study. *PeerJ*, 7, e6436. <https://doi.org/10.7717/peerj.6436>
- Vlahov, E., Baghurst, T. M., & Mwavita, M. (2014). Preschool motor development predicting high school health-related physical fitness: a prospective study. *Perceptual and Motor Skills*, 119(1), 279-291. <https://doi.org/10.2466/10.25.PMS.119c16z8>
- Volbekiene, V., & Griciūte, A. (2007). Health-related physical fitness among schoolchildren in Lithuania: a comparison from 1992 to 2002. *Scandinavian Journal of Public Health*, 35(3), 235-242. <https://doi.org/10.1080/14034940601160649>
- Wang, X., Soh, K. G., Samsudin, S., Deng, N., Liu, X., Zhao, Y., & Akbar, S. (2023). Effects of high-intensity functional training on physical fitness and sport-specific performance among the athletes: A systematic review with meta-analysis. *PLoS One*, 18(12), e0295531. <https://doi.org/10.1371/journal.pone.0295531>
- Wang, Z., Asokan, G., Onnela, J. P., Baird, D. D., Jukic, A. M. Z., Wilcox, A. J., Curry, C. L., Fischer-Colbrie, T., Williams, M. A., Hauser, R., Coull, B. A., & Mahalingaiah, S. (2024). Menarche and Time to Cycle Regularity Among Individuals Born Between 1950 and 2005 in the US. *JAMA Network Open*, 7(5), e2412854. <https://doi.org/10.1001/jamanetworkopen.2024.12854>
- Wear, C. L. (1963). Relationship of Flexibility Measurements to Length of Body Segments. *Research Quarterly for Exercise and Sport*, 34(2), 234-238.
- Welk, G. J., Saint-Maurice, P. F., & Csányi, T. (2015). Health-Related Physical Fitness in Hungarian Youth: Age, Sex, and Regional Profiles. *Research Quarterly for Exercise and Sport*, 86 Suppl 1, S45-S57. <https://doi.org/10.1080/02701367.2015.1043231>
- Wells, J. C. (2007). Sexual dimorphism of body composition. *Best Practice & research. Clinical Endocrinology & Metabolism*, 21(3), 415-430. <https://doi.org/10.1016/j.beem.2007.04.007>
- Wells, K. F., & Dillon, E. K. (1952). The sit and reach - a test of back and leg flexibility. *Research Quarterly*, 23(1), 115-118. <https://doi.org/10.1080/10671188.1952.10761965>
- Weston, K. L., Pasecinic, N., & Basterfield, L. (2019). A Preliminary Study of Physical Fitness in 8- to 10-Year-Old Primary School Children From North East England in Comparison With National and International Data. *Pediatric Exercise Science*, 31(2), 229-237. <https://doi.org/10.1123/pes.2018-0135>
- Woll, A., Worth, A., Mündermann, A., Hölling, H., Jekauc, D., & Bös, K. (2013). Age- and sex-dependent disparity in physical fitness between obese and normal weight children and adolescents. *Journal of Sports Medicine and Physical Fitness*, 53(1), 48-55.
- Woods, R. A., & U.S. Bureau of Labor Statistics. (2017). *Sports and exercise*. U.S. Bureau of Labor Statistics. <https://www.bls.gov/spotlight/2017/sports-and-exercise/home.htm>
- Wren, T. A., Aggabao, P. C., Poorghasamians, E., Chavez, T. A., Ponrartana, S., & Gilsanz, V. (2017). Association between vertebral cross-sectional area and lumbar lordosis


angle in adolescents. *PLoS One*, 12(2), e0172844.

<https://doi.org/10.1371/journal.pone.0172844>

Xu, Y., Mei, M., Wang, H., Yan, Q., & He, G. (2020). Association between Weight Status and Physical Fitness in Chinese Mainland Children and Adolescents: A Cross-Sectional Study. *International Journal of Environmental Research and Public Health*, 17(7).

<https://doi.org/10.3390/ijerph17072468>

Zhang, F., Bi, C., Yin, X., Chen, Q., Li, Y., Liu, Y., Zhang, T., Li, M., Sun, Y., & Yang, X. (2021). Physical fitness reference standards for Chinese children and adolescents. *Sci Rep*, 11(1), 4991. <https://doi.org/10.1038/s41598-021-84634-7>



**Figure 1.** Effect sizes (Hedges  $g$ ) of differences in sit-and-reach flexibility between boys and girls from birth to age 16. Black circles represent cumulative effect sizes for each age. The number of effects making up each cumulative effect size are listed in Table 1. Dashed lines around the effect sizes represent the upper and lower limits of the 95% confidence intervals (CIs). Girls had greater sit-and-reach flexibility than boys at all ages. The large effect size at age 3 is due to one large effect being available at that age. From 4 to 10 years old, the effect size generally ranged between  $g = -0.30$  to  $-0.50$ , and it became more consistent between the ages of 8 to 10 years old due, in part, to the greater number of effects available at those ages. The “dip” in the effect size at 11 and 12 years old represents an increase in the sex difference in sit-and-reach flexibility. This dip is likely due to girls reaching puberty earlier than boys. After boys reach puberty, the sex difference in sit-and-reach flexibility narrows to pre-puberty levels..

### Youth aged 5-10 years old





Figure 2. Effect sizes (Hedges  $g$ ) of differences in sit-and-reach flexibility between 5-10-year-old boys and girls from the 1980s to today. The black circles are cumulative effect sizes for all relevant study effects. The number of effects making up each cumulative effect size are listed in Table 3. The dashed lines around the effect sizes represent the upper and lower limits of the 95% confidence intervals (CI). Girls had greater sit-and-reach flexibility than boys in all decades. This sex difference has remained mostly stable since the 1980s, as evident by the overlap of the 95% CIs for most decades.

### Youth aged 14-16 years old



**Figure 3.** Effect size (Hedges  $g$ ) of differences in sit-and-reach flexibility between 14-16-years-old boys and girls from the 1980s to today. The black circles are cumulative effect sizes for all relevant study effects. The number of effects making up each cumulative effect size are listed in Table 3. The dashed lines around the effect sizes represent the upper and lower limits of the 95% CIs. Girls had greater sit-and-reach flexibility than boys in all decades. This sex difference has remained mostly stable since the 1980s, as evident by the overlap of the 95% CIs for most decades.



Table 1. Effect sizes of the difference in sit-and-reach flexibility between boys and girls by age group.

Age group	No. effects	Hedges <i>g</i>	<i>p</i>	95% CI		95% PI	
				Lower	Upper	Lower	Upper
3 yr	1	-1.35	<.001	-1.78	-0.92	NA	NA
4 yr	6	-0.30	<.001	-0.47	-0.14	-0.76	0.15
5 yr	5	-0.52	<.001	-0.73	-0.31	-1.23	0.19
6 yr	21	-0.38	<.001	-0.47	-0.14	-0.83	0.07
7 yr	24	-0.50	<.001	-0.67	-0.33	-1.39	0.38
8 yr	37	-0.40	<.001	-0.49	-0.31	-0.89	0.08
9 yr	33	-0.43	<.001	-0.53	-0.35	-0.89	0.03
10 yr	41	-0.43	<.001	-0.52	-0.33	-0.94	0.09
11 yr	51	-0.59	<.001	-0.75	-0.44	-1.70	0.52
12 yr	53	-0.78	<.001	-0.96	-0.61	-2.07	-0.50
13 yr	37	-0.72	<.001	-0.95	-0.50	-2.11	-0.67
14 yr	46	-0.61	<.001	-0.71	-0.51	-1.26	0.03
15 yr	33	-0.67	<.001	-0.79	-0.56	-1.26	-0.09
16 yr	20	-0.42	<.001	-0.57	-0.27	-1.06	0.21
Overall	408	-0.57	<.001	-0.61	-0.52	-1.46	0.33

CI = confidence interval; NA = not applicable; PI = prediction interval.

Table 2. Number of effects in the meta-analysis by country or region of data collection.

Country or region	No. effects	Percent
Australia	2	0.5
Belgium	4	1.0
Brazil	12	2.9
Canada	12	2.9
China	23	5.6
Chile	3	0.7
Columbia	1	0.2
Croatia	9	2.2
England	2	0.5
Estonia	12	2.9
Europe (multiple countries)	3	0.7
Finland	6	1.5
France	7	1.7
Germany	4	1.0
Greece	5	1.2
Hungary	6	1.5
Ireland	6	1.5
Italy	27	6.6
Japan	6	1.5
Korea (South)	15	3.7
Kosovo	2	0.5
Lithuania	16	3.9
Macedonia	9	2.2
Netherlands	14	3.4
New Zealand	2	0.5
Norway	3	0.7
Peru	12	2.9
Poland	63	15.4
Portugal	10	2.5
Russia	1	0.2
South Africa	32	7.8
Serbia	3	0.7
Slovakia	4	1.0
Spain	22	5.4
Sweden	3	0.7
Taiwan	4	1.0
United States of America	42	10.3
Wales	1	0.2
Total	408	100.0

Table 3. Effect sizes of the difference in sit-and-reach flexibility between boys and girls by age and decade.

Age group, decade	No. effects	Hedges <i>g</i>	<i>p</i>	95% CI	
				Lower	Upper
5-10 yr old					
1980s	29	-0.52	<.001	-0.61	-0.44
1990s	8	-0.45	<.001	-0.54	-0.36
2000	44	-0.29	<.001	-0.38	-0.20
2010s	68	-0.46	<.001	-0.54	-0.39
2020s	12	-0.51	<.001	-0.68	-0.34
Overall	161	-0.43	<.001	-0.48	-0.38
14-16 yr old					
1980s	11	-0.72	<.001	-1.03	-0.41
1990s	12	-0.65	<.001	-0.80	-0.51
2000	18	-0.63	<.001	-0.74	-0.53
2010s	46	-0.54	<.001	-0.63	-0.41
2020s	12	-0.64	<.001	-0.85	-0.43
Overall	99	-0.60	<.001	-0.66	-0.53

CI = confidence interval.

Table 4. Effect sizes of the difference in sit-and-reach flexibility between boys and girls by age and country.

Age group, country	No. effects	Hedges <i>g</i>	<i>p</i>	95% CI	
				Lower	Upper
5-10 yr old					
China	8	-0.48	.005	-0.81	-0.14
Italy	9	-0.62	<.001	-0.68	-0.56
Peru	10	-0.12	.200	-0.31	0.06
Poland	24	-0.35	<.001	-0.43	-0.27
South Africa	19	-0.32	<.001	-0.46	-0.26
Spain	12	-0.44	<.001	-0.57	-0.32
United States of America	26	-0.58	<.001	-0.68	-0.47
14-16 yr old					
China	8	-0.24	0.082	-0.51	0.03
Lithuania	11	-0.55	<.001	-0.73	-0.38
Poland	20	-0.63	<.001	-0.80	-0.47
United States of America	4	-0.73	<.001	-0.91	-0.54

CI = confidence interval.